

MARIA DE LURDES SOARES DE ALMEIDA

Licenciada em Química (U. P.)

Assistente de Química da Faculdade de Ciências da Universidade do Porto

**SELECTIVE PROTECTION OF POLYAMINES:
SYNTHESIS OF SPERMIDINE DERIVATIVES**

DISSERTAÇÃO PARA DOUTORAMENTO EM QUÍMICA ORGÂNICA
NA FACULDADE DE CIÊNCIAS DA UNIVERSIDADE DO PORTO

Porto
1989

MARIA DE LURDES SOARES DE ALMEIDA

Licenciada em Química (U. P.)

Assistente de Química da Faculdade de Ciências da Universidade do Porto

**SELECTIVE PROTECTION OF POLYAMINES:
SYNTHESIS OF SPERMIDINE DERIVATIVES**

DISSERTAÇÃO PARA DOUTORAMENTO EM QUÍMICA ORGÂNICA
NA FACULDADE DE CIÊNCIAS DA UNIVERSIDADE DO PORTO

Porto

1989

Aos meus Pais

Aos meus Irmãos

Aos meus Sobrinhos

À memória da minha Irmã Maria de Fátima

ACKNOWLEDGEMENTS

The experimental work described here was carried out at the Institute of Biochemistry, Biomedical Center, University of Uppsala, Sweden, from September/1986 - May/1989, under the supervision of Doctor Ulf Ragnarsson.

I wish to express my sincere gratitude to **Doctor Ulf Ragnarsson**, who kindly accepted me in his Research Group, for his constant encouragement, support, invaluable insights and guidance throughout the course of this project. His friendship, great enthusiasm and availability for discussions at any time were a constant during my stay.

I am also deeply indebted to **Dr. Leif Grehn** for his collaboration, many stimulating discussions, constructive criticism and for helping me to overcome the initial obstacles.

I also wish to express my gratitude to **The Swedish Institute** for a scholarship from September/1986 to May/1989 and to **The University of Porto**, Portugal, for a leave of absence for a period of three years.

This work was also supported by grants from **The National Swedish Board for Technical Development** and **The Swedish Natural Science Research Council**.

In addition I wish to express my thanks to:

Prof. Olle Heby, University of Umea, Sweden, for stimulating discussion on biological aspects of polyamines;

Dr. Bengt Fransson for pleasant cooperation in the h.p.l.c. studies;

Prof. Dr. Maria Joaquina S.A.A. Trigo and Prof. Dr. Carlos Corrêa at the Chemistry Department, Faculty of Sciences of Porto, Portugal, for constant encouragement and friendship;

Fil. mag. Karin Myrberg, Vasteras, Sweden, for the linguistic correction of my manuscripts during a stressed period in the month of August 1989 and specially for her skilled and enthusiastic work;

Kerstin Gunnarsson and Dr. Gunnar Johansson for their constant support, kindness and friendship;

Elisabeth Ragnarsson for her support and friendship;

All my colleagues at the Institute of Biochemistry, Biomedical Center, University of Uppsala, namely Henry Franzén, Fredrik Degerbeck, Lu Ding, Jussi Kantele, and Jozsef Bodi for their pleasant company.

My Colleagues at the Chemistry Department, Faculty of Sciences of Porto, Portugal, for their kindness and friendship.

All my Friends who contributed in some way for the success of my work.

Natural polyamines and their derivatives are important compounds, involved in many processes on the cellular level. Recent progress in this field has shown their potential as antineoplastic agents and in the treatment of parasitic diseases.

From a synthetic point of view, the simultaneous presence of primary and secondary amino groups attracted our interest, particularly so, as in a triamine such as spermidine, with a secondary and two non-equivalent primary groups. Thus, chemically as well as biologically, polyamines are challenging compounds.

Although several methods have been developed for selective functionalization of polyamines, an alternative simple procedure for selective protection of mixed primary/secondary amines seemed desirable.

The novel chemistry based on the exhaustive tert-butoxy-carbonylation of amide type groups, which was developed in Doctor Ragnarsson's group, seemed a promising basis for the development of new methodologies for the synthesis and selective protection of polyamines. Thus, the present work, carried out during nearly three years at the Institute of Biochemistry, Biomedical Center, University of Uppsala, Sweden, under the supervision of Doctor Ulf Ragnarsson, had as its principal objective the study of selective protection of spermidine using the new approach, the DMAP-catalysed N-tert-butoxycarbonylation of urethane groups.^{1,8}

It was possible to synthesize the N¹,N⁸-bis(tert-butoxycarbonyl)spermidine and N¹-benzyloxycarbonyl-N⁸-tert-butoxycarbonylspermidine. Then was studied the applicability of these compounds to synthetic work with the preparation of some biologically interesting substrates, such as acetyl and ethyl spermidine derivatives.

In connection with the previous study and aiming at total synthesis of polyamines, several Gabriel type reagents, ZNHCOOR, were also prepared. It is expected that in the future these derivatives will be useful for the synthesis of natural polyamine-containing substrates.

The results obtained during this research project are the subject of this dissertation, which also includes an introductory section on the biological aspects of natural polyamines and a review of the methods commonly used in their synthesis.

Part of the results presented in this thesis is the basis of four original articles published in collaboration^{109a-d}.

CONTENTS

Abbreviations

1- <u>Polyamines</u>	1
1.1- <u>Biological aspects</u>	1
1.2- <u>Synthesis of polyamines and their analogues</u>	11
1.2.1- Introduction	11
1.2.2- Total synthesis of polyamines	11
1.2.2.1- Alkylating methods	12
A. Alkylation of tosylamides	12
B. Alkylation of amino groups	16
1.2.2.2- Reductive methods	18
A. Reduction of nitriles	18
B. Reduction of amides	24
C. Reduction of imines	26
D. Reduction of azides	28
1.2.3- Direct selective protection of polyamines ...	30
1.2.3.1- Regioselective acylation and alkoxy- carbonylation of primary amines	30
A. 2-Oxazolones	31
B. Thiazolidine-2-thiones	33
C. Imidazoles	35
D. Benzo- and naphthosultones	36
E. 1-Hydroxypiperidine	38
F. Nefkens's reagent	40
G. Nitriles	41
1.2.3.2- "Transiently protected" polyamines .	42
A. Cyclic ureas	43
B. Hexahydropyrimidines	45
1.2.4- Conclusions	49

2- <u>Background of the present work</u>	52
2.1- <u>Introduction</u>	52
2.2- <u>tert-Butoxycarbonylation of amide type groups</u>	55
2.3- <u>Selective cleavage of amides</u>	57
3- <u>Objectives and synthetic methodology</u>	60
4- <u>Results and conclusions</u>	67
4.1- <u>Selective protection of spermidine</u>	67
4.1.1- Synthesis of N ¹ ,N ⁸ -bis(<u>tert</u> -butoxycarbonyl)- spermidine	67
4.1.2- Synthesis of N ¹ -benzyloxycarbonyl-N ⁸ - <u>tert</u> - butoxycarbonylspermidine	70
4.1.3- Attempted synthesis of N ⁸ -benzyloxycarbonyl- N ¹ - <u>tert</u> -butoxycarbonylspermidine	74
4.2- <u>Synthesis of modified spermidines</u>	76
4.2.1- Synthesis of N ⁴ -acetylspermidine dioxalate...	76
4.2.2- Synthesis of N ¹ -acetylspermidine dihydro- chloride	79
4.2.3- Synthesis of N ⁸ -acetylspermidine dihydro- chloride	81
4.2.4- Synthesis of N ¹ ,N ⁸ -diethylspermidine tri- oxalate	83
4.2.5- Attempted syntheses of N ¹ -ethyl- and N ⁸ ethylspermidine	85
A. Alkylation experiments	85
B. Reduction experiments	90
4.3- <u>Synthesis of alkyl benzyl imidodicarbonates</u>	93
4.4- <u>Concluding comments</u>	96

5- <u>Experimental</u>	101
Materials. General methods	101
5.1- <u>Experiments with model compounds</u>	103
5.1.1- Synthesis of N ² -benzyloxycarbonyl-N ¹ - <u>tert</u> - butoxycarbonyl-N ¹ -ethylethylenediamine	103
5.1.2- Synthesis of N ¹ -benzyloxycarbonyl-N ² - <u>tert</u> - butoxycarbonyl-N ¹ ,N ² -diethylethylenediamine .	104
5.1.3- Synthesis of N ¹ - <u>tert</u> -butoxycarbonyl-N ¹ - ethylethylenediamine	106
5.1.4- Synthesis of N ² -acetyl-N ¹ - <u>tert</u> -butoxy- carbonyl-N ¹ -ethylethylenediamine	107
5.1.5- Synthesis of N ¹ - <u>tert</u> -butoxycarbonyl-N ¹ ,N ² -di- ethylethylenediamine	108
5.1.6- Synthesis of N ² -acetyl-N ¹ -ethylethylene- diamine	109
5.1.7- Synthesis of N ¹ ,N ² -diethylethylenediamine dihydrochloride	110
5.1.8- Synthesis of N ¹ -benzyloxycarbonyl-N ⁴ - <u>tert</u> - butoxycarbonylputrescine	112
5.2- <u>Experiments with spermidine</u>	113
5.2.1- Synthesis of N ¹ ,N ⁸ -bis(<u>tert</u> -butoxycarbonyl)- spermidine	113
N ¹ ,N ⁴ ,N ⁸ -Tribenzyloxycarbonylspermidine ...	113
N ¹ ,N ⁴ ,N ⁸ -Tribenzyloxycarbonyl-N ¹ ,N ⁸ -bis- (<u>tert</u> -butoxycarbonyl)spermidine	114
N ¹ ,N ⁸ -Bis(<u>tert</u> -butoxycarbonyl)spermidine ..	116
5.2.2- Synthesis of N ⁴ -benzyloxycarbonyl-N ¹ ,N ⁸ -bis- (<u>tert</u> -butoxycarbonyl)spermidine	117
5.2.3- Synthesis of N ¹ -benzyloxycarbonyl-N ⁸ - <u>tert</u> - butoxycarbonylspermidine	119
N ¹ ,N ⁴ -Methylenespermidine	119
N ¹ ,N ⁸ -Dibenzyloxycarbonyl-N ¹ ,N ⁴ -methylene- spermidine	119

	N^1, N^8 -Dibenzylloxycarbonyl- N^8 - <u>tert</u> -butoxy-carbonyl- N^1, N^4 -methylenespermidine	120
	N^1 -Benzylloxycarbonyl- N^8 - <u>tert</u> -butoxy-carbonylspermidine	121
	N^1 -Benzylloxycarbonyl- N^8 - <u>tert</u> -butoxy-carbonyl- N^1, N^4 -methylenespermidine	121
	N^1, N^8 -Dibenzylloxycarbonyl- N^8 - <u>tert</u> -butoxy-carbonylspermidine	123
	N^1 -(Benzylloxycarbonyl- β -alanyl)- N^4 - <u>tert</u> -butoxycarbonyltetramethylenediamine	124
5.2.4-	Attempted synthesis of N^8 -benzylloxycarbonyl- N^1 - <u>tert</u> -butoxycarbonylspermidine	127
	N^8 -Benzylloxycarbonyl- N^1 - <u>tert</u> -butoxy-carbonyl- N^1, N^4 -methylenespermidine	127
	N^8 -Benzylloxycarbonyl- N^1 - <u>tert</u> -butoxy-carbonylspermidine	129
5.2.5-	Synthesis of N^4 -acetylspermidine dioxalate...	130
	N^1, N^4, N^8 -Triacetylspermidine	130
	N^1, N^4, N^8 -Triacetyl- N^1, N^8 -bis(<u>tert</u> -butoxy-carbonyl)spermidine	131
	N^4 -Acetyl- N^1, N^8 -bis(<u>tert</u> -butoxycarbonyl)-spermidine	133
	N^4 -Acetylspermidine dioxalate	134
5.2.6-	Synthesis of N^1 -acetylspermidine dihydrochloride	135
	N^1 -Benzylloxycarbonyl- N^4, N^8 -bis(<u>tert</u> -butoxy-carbonyl)spermidine	135
	N^4, N^8 -Bis(<u>tert</u> -butoxycarbonyl)spermidine ..	136
	N^1 -Acetyl- N^4, N^8 -bis(<u>tert</u> -butoxycarbonyl)-spermidine	137
	N^1 -Acetylspermidine dihydrochloride	137
5.2.7-	Synthesis of N^8 -acetylspermidine dihydrochloride	138
	N^1, N^4 -Dibenzylloxycarbonyl- N^8 - <u>tert</u> -butoxy-	

carbonylspermidine	138
N ¹ ,N ⁴ -Dibenzoyloxycarbonylspermidine	140
N ⁸ -Acetyl-N ¹ ,N ⁴ -dibenzoyloxycarbonyl- spermidine	140
N ⁸ -Acetylspermidine dihydrochloride	141
5.2.8- Synthesis of N ¹ ,N ⁸ -diethylspermidine trioxalate	144
N ¹ ,N ⁴ ,N ⁸ -Tribenzoyloxycarbonyl-N ¹ ,N ⁸ -di- ethylspermidine	144
N ¹ ,N ⁸ -Diethylspermidine trioxalate	145
5.2.9- Attempted syntheses of N ¹ -ethyl- and N ⁸ -ethylspermidine	146
N ¹ -Ethylspermidine trioxalate	146
N ⁸ -Ethylspermidine trioxalate	148
5.3- <u>Synthesis of alkyl benzyl imidodicarbonates</u>	149
5.3.1- Benzoyloxycarbonyl isocyanate	149
5.3.2- Alkyl benzyl imidodicarbonates	150
General procedure	150
Table of compounds synthesized	156
References	160
Abstract	170
Resumo	172


ABBREVIATIONS

Ac	acetyl
Adoc	1-adamantylloxycarbonyl
Aloc	allyloxycarbonyl
aq.	aqueous
ATP	adenosine triphosphate
Bz	benzoyl
Bzl	benzyl
Boc	<u>tert</u> -butoxycarbonyl
Boc-ON	<u>tert</u> -butoxycarbonyloxyimino-2-phenylacetonitrile
Bu _t	<u>n</u> -butyl
Bu ^t	<u>tert</u> -butyl
CDI	N,N'-carbonyldiimidazole
DC	decarboxylase
DEAEA	2-diethylaminoethylamine
DFMO	α -difluoromethylornithine
DMAP	4-dimethylaminopyridine
DMF	N,N-dimethylformamide
DNA	deoxyribonucleic acid
DPP-ox	diphenyl-2-oxo-3-oxazolinyolphosphonate
eq.	equivalent
Et	ethyl
Fmoc	9-fluorenylmethoxycarbonyl
h	hour
Hox	2-oxazolone
h.p.l.c.	high-performance liquid chromatography
Me	methyl
m.p.	melting point
MPP	4-(4-methyl-1-piperidiny)pyridine
NMM	N-methylmorpholine
n.m.r.	nuclear magnetic resonance
ODC	ornithine decarboxylase
Ph	phenyl
Pht	phthaloyl
Ppoc	2-phenylisopropylloxycarbonyl
Pr ⁱ	iso-propyl
RCO-Im	acylimidazoles
RCO-ox	3-acyl-2-oxazolone
RCO-TT	3-acylthiazolidine-2-thione
Red.	reduction
Red-Al	sodium bis(2-methoxyethoxy)aluminiumhydride
r.t.	room temperature
spd	spermidine
TcBoc	2,2,2-trichloro- <u>tert</u> -butoxycarbonyl
TEA	triethylamine
TFA	trifluoroacetic acid
THF	tetrahydrofuran
t.l.c.	thin layer chromatography
TMG	N,N,N',N'-tetramethylguanidine

TMS	tetramethylsilane
Tos	tosyl
Troc	2,2,2-trichloroethoxycarbonyl
Z	benzyloxycarbonyl
Z(NO ₂)	4-nitrobenzyloxycarbonyl
ZOBt ²	benzyl benzotriazol-1-yl carbonate
Z(OMe)	4-methoxybenzyloxycarbonyl
Z-TT	3-benzyloxycarbonylthiazolidine-2-thione

Notes:

- a) The nomenclature of the protecting groups is that recommended by IUPAC-IUB as summarized in Pure Appl. Chem., 1984, **56**, 595.
- b) The bibliographic references are presented in order of their appearance. The abbreviations of the journals are those adopted by the Chemical Society of London.
- c) According to the ref. 35a, spermidine is numbered as follows:



1 - POLYAMINES

1.1 - Biological aspects

The aliphatic polyamines putrescine 1, spermidine 3, and spermine 4 (Fig. 1) constitute the principal members of a family of natural products present in a relatively high concentration in most living organisms.




Figure 1 Structural representation of the naturally occurring polyamines. Spermine is unique to eukaryotes whereas cadaverine is found mainly in prokaryotes.

During the last few years there has been an increasing interest in polyamines and great progress has been made in this field. Although their mechanisms of action are not yet understood in detail, many studies have shown that these substances play a key role in a variety of cellular processes. In this chapter some of the present developments will be briefly outlined whereas detailed aspects of polyamine biochemistry and physiology can be found in recent excellent

reviews¹⁻⁶.

The general metabolic reactions responsible for polyamine biosynthesis and interconversion are outlined in Fig. 2. The initial step is the formation of putrescine 1. In organisms lacking arginine decarboxylase, such as mammalian cells and lower eukaryotes, the only route to putrescine is that catalysed by the initial and rate-limiting enzyme ornithine decarboxylase (ODC). On the other hand, plants and some bacteria can initiate this synthetic pathway either directly from ornithine or indirectly from arginine through the activity of arginine decarboxylase via the intermediate agmatine. Putrescine is then converted to spermidine by the enzymatic coupling of an aminopropyl group, derived from decarboxylated S-adenosylmethionine, to one of the terminal amino groups by spermidine synthase. In a reaction catalysed by another aminopropyl transferase, spermine synthase, spermidine is converted to spermine. Prokaryotes lack this last enzyme and therefore spermine. Although the aminotransferase reactions are irreversible, these polyamines can be interconverted via consecutive enzymatic N-acetylation and oxidation reactions. This cyclic process has been explained in terms of the prevention of toxic levels of intracellular polyamines either by their interconversion and degradation or facilitation of their excretion from the cell due to a decrease in the net charge of the polyamine⁷.

Under physiological conditions the polyamines are largely protonated. Thus, these conformationally mobile polycations




Figure 2 Biosynthesis and interconversion of polyamines⁶.

can associate to various extents with anionic binding sites in nucleic acids and membrane phospholipids through electrostatic charge interactions^{4,6,8}. With such macromolecules these interactions have specific structural requirements. For instance, in double-helical regions of DNA the positively charged amino groups of spermine strongly bind to two phosphate groups on one DNA strand and the tetramethylene chain of the polyamine molecule bridges the minor groove to interact with two phosphates on the second DNA strand (Fig.3). Thus, spermine stabilizes the double helix by binding its two




Figure 3 Proposed model for spermine/double-helical DNA interaction^{8a}.

strands together. This effect is also correlated with the ability of the polyamines to induce the transition from the usually right-handed (B-DNA) to a left-handed (Z-DNA) double helix conformation of DNA. This property has been correlated with a large variety of biological effects such as the stabilization of nucleic acids and the stimulation of RNA, DNA, and protein biosynthesis⁶.

Many findings indicate that these natural polycations have a key role in cell growth and proliferation¹⁻⁶. These include the observation that: polyamines serve as growth factors for cultured cells; they are found in larger amounts in growing than in non-growing tissues; prokaryotic and eukaryotic mutants deficient in polyamine biosynthesis are auxotrophic for polyamines, and, a more recent and stronger evidence, depletion of intracellular polyamine levels by highly

specific inhibitors of their biosynthesis results in a cessation of cell growth. The latter feature has received great attention as it provides biological information regarding polyamine function and, moreover, it offers promising prospects for antineoplastic drug action¹⁻⁶. The potential of interference with polyamine biosynthesis and/or function as an anticancer chemotherapeutic strategy is due to the fact that polyamines are essential for cell proliferation, the rates of polyamine uptake and biosynthesis are increased in neoplastic tissues, and the ability of inhibitors to slow down neoplastic cell growth.

Ornithine decarboxylase (Fig. 2), has been the principal target for designing irreversible enzyme-activated inhibitors ("suicide inhibitors"). α -Difluoromethylornithine (DFMO) is a potent and specific ODC inhibitor¹⁻⁶. Its specificity is due to the fact that only ODC can decarboxylate and thus activate this substrate analogue. After the enzymatic decarboxylation, the highly reactive electrophilic intermediate alkylates a nucleophilic residue within the active site, thus inactivating the enzyme (Fig. 4)⁶. By inhibiting the ODC activity, DFMO is effective in depleting intracellular putrescine and spermidine. This process can be restored by adding exogenous polyamines to the cells treated. This inhibitor has proved important in delineating the effects of polyamine depletion in animals⁶. For instance, inhibition of polyamine synthesis by DFMO treatment in early embryogenesis suppresses protein synthesis and prevents development. Another important feature




Figure 4 Postulated mechanisms for the irreversible inhibition of ODC by DFMO⁶.

is that when cells in culture are depleted of polyamines with this inhibitor, their progression through the cell cycle is slowed down and sometimes arrested. This cytostatic effect of DFMO has been exploited in the treatment of tumour and cancer cells due to a difference in the response to the inhibitor between normal and transformed cells. Thus, either alone or in combination with other cytotoxic agents, DFMO has been tested clinically as an anticancer agent and encouraging results have been obtained in leukaemia and melanoma⁶. Besides being promising for an anticancer chemotherapy strategy, DFMO has also been exploited in the treatment of the major parasitic diseases, such as African trypanosomiasis (sleeping sickness), malaria, cryptosporidiosis, and more recently, *Pneumocystis carinii*, an opportunistic protozoan infection in patients with

acquired immune deficiency syndrome (AIDS)⁶.

Although DFMO is essentially nontoxic it has certain limitations. One of them is the cytostatic rather than cytotoxic response of most tumour systems where the arrest in tumour cell proliferation is rapidly reversed by the removal of the drug or uptake of polyamines. Other limitations are that DFMO is soon cleared from the body and enters into the cells by diffusion rather than by an active amino acid transport. Thus a high concentration is necessary.

Another promising alternative to specific inhibitors of polyamine biosynthetic enzymes is based on the use of polyamine analogues bearing a close structural resemblance to the natural polyamines, such as the ethylated spermidine and spermine derivatives^{4,5,9}. The basic idea behind the design of these analogues is that they may be taken up into the cell by the polyamine transport system which is more active in proliferating cells. Once inside the cell, polyamine analogues could exert antiproliferative effects by some of the following mechanisms: inhibition and/or regulation of polyamine biosynthetic enzymes; competition for polyamine binding sites and subsequent disruption of critical macromolecular structure and/or function, or as vector molecules for delivering to cancer cells biologically active moieties or small antineoplastic agents. Studies with N⁴- or N¹,N⁸-substituted spermidine analogues showed that the primary amino groups of spermidine are more critical than the secondary one as determinants of cellular uptake and in functions required for

cell proliferation. The central amino group is more important to regulatory activities of spermidine relevant to ODC activity. The alkylated spermidine analogues are more effective derivatives than the corresponding acylated ones because the presence of a positive charge at any of the amino functionalities also plays a very important role in uptake specificity^{9d}.

In accordance with these features, regulation of polyamine biosynthesis by polyamine analogues seems to represent another antiproliferative strategy⁹. Generally this approach might offer certain advantages relative to specific enzyme inhibitors (Table I): by utilizing the polyamine transport system, the derivatives should penetrate cells more effectively and at relatively low concentrations; the activity of more than one biosynthetic enzyme may be negatively regulated at the same time; compensatory increases in related enzymes may not occur as they do with enzyme inhibitors and might give substantial decreases in the pools of all polyamines including spermine.

Besides their existence in free form, the most common polyamines, spermidine and spermine, and their homologues are incorporated in many naturally occurring products such as sugars¹⁰, phospholipids¹¹, peptides¹², alkaloids¹³, and siderophores¹⁴, which also have a wide range of important biological activities (Table II, Fig. 4).

In conclusion, natural polyamines seem to have an important role in cell growth and differentiation and, like their

conjugates, have potential as drugs in medicine. These features have prompted the outline of synthetic routes to polyamines and analogues which will be reviewed in the next section.

Table I - Cellular effects by treatment with modulators of polyamine biosynthesis⁴.

Parameter	DFMO	DEspd ^a
Effective dose	1.5 mM	10-100 μ M
Growth effect	Cytostatic	Cytostatic
ODC activity	Depleted	Depleted
S-Adenosylmethionine DC activity	Increased	Decreased
Putrescine	Depleted	Depleted
Spermidine	Depleted	Depleted
Spermine	Increased	Decreased
S-Adenosylmethionine	Decreased	Unchanged
Decarboxy-S-adenosylmethionine	Increased	Unchanged
Spermidine uptake	Increased	Unchanged

^aN¹,N⁸-Diethylspermidine.

Table II - Examples of natural polyamine-containing compounds.

N ^o	Compound	Biological properties	ref.
5a-c	Glycocinnamoyl-spermidines	Broad-spectrum antibiotics	10
6	Trypanothione	Trypanosomatid metabolite	12
7	Solapalmitine	Tumour inhibitory	13c
8	Agrobactin	Iron-chelating	14a
9a-c	Acarnidines	Antiviral and antimicrobial	15
10	Spergualin	Antitumour antibiotic	16









Figure 5 Structures of some of the natural polyamine-containing compounds.

1.2 - Synthesis of polyamines and their analogues

1.2.1 - Introduction

The wide application of synthetic derivatives of polyamines for biological purposes has created a demand for efficient synthetic procedures. Thus, an important field in polyamine chemistry is the development of syntheses of such compounds. The remaining part of this chapter provides a comprehensive review of the previous work in this field.

A major problem in the synthesis of naturally occurring polyamines and their analogues is the selective modification of the different amino groups. As simple approaches suffer from problems of regioselectivity¹⁷, in recent years different procedures have been developed for selective modification and functionalization of various polyamines^{18,19}.

The two general strategies, which are based on total synthesis or on selective protection of preexisting polyamines, will be described below.


1.2.2 - Total synthesis of polyamines

This methodology is of general applicability by which the complete backbone of polyamines is elaborated using adapted preparative methods for amines such as alkylation and reduction reactions²⁰.

1.2.2.1- Alkylating methods

A. Alkylation of tosylamides


A key step for the assembly of the carbon-nitrogen framework is the mono- or di-N-alkylation of N-protected-N'-Tos-alkanediamines and N,N'-Tos₂-alkanediamines, respectively, with haloalkanes bearing a latent primary amine group to afford the protected polyamines (Scheme 1)²¹.



Scheme 1 Alkylation of tosylamides. Reagents: i, PhtN(CH₂)₃ Br, K₂CO₃ (DMF)²¹.

This method has been used for the synthesis of natural polyamine-containing substrates^{22,23}. The N-protecting groups most preferred during alkylation are either the phthaloyl²¹ or some of acyl²² or urethane²³ type. The alkylating agents are the N-haloalkylphthalimides^{21,22b} and dihaloalkanes^{22a}.

The first reported threefold protected spermidine, N⁸-Z-N¹Pht-N⁴-Tos-spermidine, designed by Eugster et al.²³ in their synthetic work with polyamine-containing lactams, is an example of the applicability of this approach (Scheme 2).




Scheme 2 Synthesis of a fully protected spermidine. Reagents: i, NaN_3 ; ii, H_2 , Pd-C (EtOH); iii, ZCl (aq. NaHCO_3); iv, NH_2NH_2 (EtOH); v, TosCl (aq. NaOH); vi, $\text{PhtN(CH}_2)_3\text{Br}$, K_2CO_3 (DMF).

More recently, in their studies of water-soluble carbodiimides which mimic the role of ATP/DNA via an autocatalytic pathway, Dörwald et al.²⁴ have designed carbodiimides bearing a DNA-binding side chain which, by increasing the relative stability of the mixed complex carbodiimide template increase the rate of autocatalysis. Thus, they have synthesized polyamine-carbodiimides in which the phosphate-activating property of carbodiimide and the DNA-binding property of spermine are combined.

The preparation of the key precursor for the synthesis of the target carbodiimides, the tetramethylated spermine, was based on this methodology as shown in Scheme 3. A suitable terminally differentiated diaminobutane, **22**, was subsequently N-alkylated with three alkylating agents of

type $X(\text{CH}_2)_3\text{R}$ (where $X = \text{Br}, \text{I}$ and $\text{R} = \text{Cl}, \text{N}(\text{Me})\text{Tos}, \text{NPht}$). Two alkylating reagents carry a latent functionality for a primary amino group, the third one a latent functionality for a dimethylamino group. Two routes were available for the assembly of the carbon-nitrogen framework both starting with monoalkylation of tosylamide **22**. The synthetic pathway using the phthalimide group as a primary amine genitor, has proved less effective. This is due to its low stability under the reaction conditions, mainly in the deformylation step and in the cleavage of the tosyl group. Once again, in the final step, the use of the phthalimide derivative was troublesome where the methylated spermine was contaminated with minor amounts of the corresponding alcohol. Instead, replacement of the chlorine atom and the conversion of bromide into azide followed by reduction to afford the spermine derivative **29** have turned out to be a better alternative.


Although this procedure has allowed the preparation of selectively modified spermidine and spermine, it is important to point out several aspects: the large number of steps involved, the drastic reaction conditions required for the removal of the protecting groups²⁵ and, in some cases, the absence of totally site-specific alkylation²⁶⁻²⁸, which makes this method less attractive.



Scheme 3 Synthesis of $\text{N}^1, \text{N}^1, \text{N}^4, \text{N}^8$ -Me₄-spermine. Reagents: i, Na, MeOH; ii, $\text{I}(\text{CH}_2)_3\text{N}(\text{Me})\text{Tos}$ (DMF); iii, conc. HCl (dioxan, reflux), base; iv, TosCl , TEA (CH_2Cl_2); v, NaH (DMF, 60 °C); vi, $\text{Br}(\text{CH}_2)_3\text{NPht}$; vii, $\text{Br}(\text{CH}_2)_3\text{Cl}$; viii, 48 % HBr (reflux); ix, 99 % HCOOH , 35 % HCHO (80 °C); x, NaN_3 (aq. MeOH, 60 °C); xi, Na_2S , (NaOH, 60 °C); xii, PhtNK_3 (DMF); xiii, NaOH (reflux), HCl (reflux), NaOH; xiv, 0.48 M CH_3ONa (MeOH)²⁴.

B. Alkylation of amino groups

Employing neutral alkylation with KF-celite³⁰ of amino groups, Samejima et al.²⁹ have prepared ¹⁵N-enriched spermidine and spermine. The key compounds of this method were putrescine or benzylamine as amine reagents and, Br(CH₂)_nNPht (n= 3 or 4) as an aminoalkyl donor. The Scheme 4 depicts the two general procedures.



Scheme 4 Synthesis of spermidine. Reagents: i, ZCl (AcONa, EtOH); ii, PhtN(CH₂)₃Br (KF-Celite, CH₃CN, reflux); iii, HCl; iv, PhtN(CH₂)₄Br² (KF-Celite, CH₃CN, reflux); v, NH₂NH₂; vi, H₂/PtO₂.

Procedure (a) involved an alkylation of monobenzyloxycarbonylputrescine in the presence of KF-celite to afford the diprotected spermidine **36** and traces of the dialkylated product. After removal of protecting groups by acid hydrolysis, spermidine was purified on a cation exchange column to afford the trihydrochloride salt of spermidine in 33 % overall yield. In procedure (b), benzylamine was successively alkylated with the proper alkylating agents to afford the triprotected spermidine **40**. The spermidine could be obtained in 30 % yield by successive removal of the protecting groups.

This method was extended to the synthesis of spermine based on the previous procedure (b). The first step involved the synthesis of dibenzylputrescine via reduction of the Schiff base formed from putrescine and benzaldehyde. The secondary amino groups were further alkylated in a similar way to afford the tetraprotected spermine which could be deprotected to afford the tetrahydrochloride salt of spermine.


Recently, Kunesch et al.³¹, have described the synthesis of N¹- and N⁸-monoacylated spermidines by monoalkylation of diaminoalkanes with the intermediate $\text{Br}(\text{CH}_2)_n\text{NHCOR}$ ³².

The main disadvantages of these approaches are the monoacylation and alkylation steps which afford to some extent the disubstituted derivatives leading to low yield reactions and purification problems. To minimize these by-products it is necessary to use large amounts of amine compound relatively to the acylating or alkylating agent.


1.2.2.2 - Reductive methods

A. Reduction of nitriles

The basic idea of this method is the reduction of nitriles, themselves obtained by cyanoalkylation, to afford part of the C-N polyamine backbone^{20,33}.



Thus, polyamines can be prepared by mono and dicyanoalkylation of the appropriate diamines followed by catalytic reduction of the resulting nitriles (Scheme 5)³⁴. The mono-cyanoethylated derivatives were prepared by dropwise addition of 1 equivalent of acrylonitrile into the diamine to avoid the possibility of dicyanoethylation which decreases with increase in the methylene chain of the diamine. In a similar way the dinitriles were prepared by 2 equivalents of acrylonitrile. The mononitriles could be purified by vacuum distillation below 160-170 °C. Above this temperature they were either reconverted to starting material by elimination of acrylonitrile or extensively decomposed. On the other hand, although the lower dinitriles could, to some extent, be isolated by distillation, the higher ones underwent extensive decomposition. The final step could then be carried out by catalytic hydrogenation (4 atm) with sponge Raney nickel in NH₃-saturated ethanol to suppress the formation of secondary amines.



(n = 2, 3, 4, 5, 6, 9, 10, 12)

Scheme 5 Synthesis of spermidine, spermine and their homologues. Reagents: i, $\text{CH}_2=\text{CHCN}$; ii, H_2/Ni (NH_3 , 3-4 atm)³⁴.

Besides, for the polyamines themselves, this methodology has proved useful for preparation of the precursors for synthesis of several polyamine-containing substances³⁵⁻³⁹.


Tabor et al.³⁵ have reported the synthesis of the two monoacetylspermidines starting from 1,3-diaminopropane or putrescine which were monoacetylated in glacial acetic acid with acetic anhydride (30-50 %). The $\text{N}^1\text{-Ac-}$ or $\text{N}^8\text{-Ac-}$ -spermidine was then prepared by reaction of 4-bromo-butyronitrile or acrylonitrile, respectively, followed by catalytic reduction (H_2/PtO_2) of the mononitrile.

In their synthetic work with alkaloids Quick et al.³⁶ have designed a synthesis of $\text{N}^4, \text{N}^8\text{-Boc}_2\text{-spermidine}$. The mononitrile adduct of putrescine, prepared as usual, was bis-tert-butoxycarbonylated with Boc-ON and, the resulting protected nitrile, was selectively reduced with LiAlH_4 in 70 % yield.

Thus, the N¹ nitrogen of spermidine remained free for further functionalization. In principle, this procedure can be extended to other mononitriles and dinitriles to afford the corresponding derivatives of spermine and its homologues. In the mononitrile series, only one primary nitrogen becomes free for selective modification and in the dinitrile one, it is only possible to differentiate the primary amino groups from the secondary ones. Therefore, this approach is of limited value for a general preparation of selectively protected polyamines. Moreover the reduction step may also be a limiting factor in the choice of the type of protective groups.

More recently, based on the above procedure, Bergeron et al.^{19,40} have developed a comprehensive methodology for selective modification of polyamines via their N-benzylated derivatives. The preparation of these compounds involved the same starting material, benzylamine, and the three basic reactions (Scheme 6). The first step consisted in consecutive monocyanoalkylations (spermidine derivatives) or dicyanoalkylation (symmetric spermidine homologues) of benzylamine. By selective reduction of the nitriles with LiAlH₄/AlCl₃ the monobenzylated polyamines **49**, **53**, and **55** were obtained. The corresponding free polyamines could then be obtained by debenzylation of these derivatives or directly from the benzylated nitriles by hydrogenolysis in acetic acid over a palladium catalyst.

Based on protection / deprotection of the amino groups of



Scheme 6 Synthesis of spermidine, homospermidine, and norspermidine via derivatives N-benzylated on their secondary amino groups. Reagents: i, $\text{CH}_2=\text{CHCN}$; ii, $\text{Cl(CH}_2\text{)}_3\text{CN}$ (butanol, Ca_2CO_3); iii, $\text{LiAlH}_4/\text{AlCl}_3$; iv, H_2/Pd ; v, H_2/PtO_2 .

these benzylated derivatives, this research group⁴¹ has synthesized the N^1, N^8 -Boc₂-spermidine (and its homologues) by direct tert-butoxycarbonylation with Boc-ON followed by hydrogenolysis of the benzyl group. Thus, these two types of polyamine precursors were further used with success as starting material for preparation of N^1, N^8 - and N^4 -substituted polyamines such as siderophores⁴².


Although this approach is convenient and relatively simple, it is of limited applicability and, therefore, as a

complement to the above precursors, this group designed a triprotected spermidine and its homologues⁴³, which could then be extended to the spermine series⁴⁴.

Scheme 7 summarizes the different steps for the preparation of the fully protected polyamines. The nitriles of the type 51 (see Scheme 6) were reduced to diamine 57 by the usual procedure. Surprisingly, the authors could selectively tert-butoxycarbonylate the primary amino group using one equivalent of Boc-ON at 0 °C in high yield⁴³. These protected diamines were then cyanoalkylated to give the protected mononitriles which by selective reduction gave the diprotected polyamine 60. The final step involved the protection of the free amino group with the orthogonal protective trifluoroacetyl group²⁵.

Alternatively, the diprotected triamine 60 was cyanoethylated followed by selective reduction of the nitrile group. The resulting diprotected spermine 63 could be selectively trifluoroacetylated⁴⁴ at its primary amino function using the active ester N-(trifluoroacetoxy)succinimide⁴⁵. The final tetraprotected polyamine was obtained by acylation with TcBocCl⁴⁶.

These fully protected polyamines have been used with success as precursors of the polyamine backbone of siderophores⁴⁷ and alkaloids⁴⁸ and the orthogonality of these protective groups^{43,44} should allow the access to a large number of selectively functionalized polyamines provided they are stable to the reaction conditions.




Scheme 7 Synthesis of selectively, fully protected spermidine and spermine. Reagents: i, H₂/Raney nickel (aq. EtOH, NaOH, 4 atm); ii, Boc-ON (THF, 0 °C); iii, Cl(CH₂)₃CN (BuOH, NaCO₃, KI); iv, (CF₃CO)₂O (TEA, CH₂Cl₂); v, CH₂=CHCN (MeOH); vi, benzene solution of 0.34 M N-CF₃CO-succinimide (CH₂Cl₂, 0 °C); vii, TcBocCl (ether, 0.2 M NaOH)^{43,44}.

The main drawback of this procedure is the reduction of the nitriles. Although the authors have previously accomplished the reduction with metal hydrides, the isolated yields were moderate (60 %). Further experiments, carried out by the same group⁴⁹, have shown that the best procedure for reduction of the nitriles was hydrogenolysis (2.5-3 atm) using Raney nickel in ethanol in the presence of sodium hydroxide. Substituting ammonium hydroxide for sodium hydroxide or pretreating the catalyst with sodium hydroxide gave only 17 % and 33 %, respectively, of the desired amine. Due to these results the authors concluded that the sodium hydroxide must play an active role in this process and not simply preactivate the catalyst. Although this reduction method worked nicely for the different nitriles of N⁴-benzylated polyamines giving high yields of these protected analogues, there are a few limitations. One of these is the low yield if the reaction is carried out on a small scale due to the adsorption of the reactant or product on the catalyst. Another limitation is the strong alkaline conditions of the reduction which restrict the choice of the protecting groups.

B. Reduction of amides


This method is based on the reduction of an amide group^{20,33} as the key step for assembly of the polyamine framework. The general procedure involves two main steps: the formation of an amide type bond and the reduction of the resulting polyamide as illustrated in Scheme 8⁵⁰.



Scheme 8 Synthesis of N -alkyl spermidines by reduction of amides. Reagents: i, RR^1NH (pyridine); ii, K_2CO_3 (MeOH, $50^\circ\text{H}_2\text{O}$, reflux); iii, $\text{CF}_3\text{CONH}(\text{CH}_2)_2\text{COCl}$ (pyridine); iv, $\text{BH}_3\cdot\text{Me}_2\text{S}$.

Using this approach, Das et al.⁵¹ have synthesized the N^4, N^8 - Boc_2 - N^1 -Pht-spermidine and N^1, N^8 - Boc_2 -spermidine as the key intermediates for synthesis of several acylated conjugates of spermidine (Scheme 9). The initial step was the condensation of N -Boc-4-aminobutyric acid with 3-amino-1-chloropropane using the mixed anhydride as coupling method. The resulting amide could then be selectively reduced to amine by $\text{Na}(\text{CF}_3\text{COO})\text{BH}_3$ ^{51b} or first deprotected followed by reduction of the amide group with borane^{51a}.

The main difficulty of this method is the reduction of the amide function. Although amides can be selectively reduced in the presence of urethane groups^{51b, 52}, it is of limited application. In some cases the desired amine has been obtained in low yield (30%)^{51a} or only the methylated amine was formed^{51a, 53, 54}.




Scheme 9 Synthesis of two protected spermidines. Reagents: i, ClCO_2Et (TEA, THF, 0-5 °C); ii, $\text{Cl}(\text{CH}_2)_3\text{NH}_2$ (TEA, $\text{CH}_2\text{CN-CH}_2\text{Cl}$);² iii, TFA (ether); iv, $\text{BH}_3\cdot\text{Me}_2\text{S}$ (THF, 80 °C, N_2);³ v, Boc_2O_2 (Na_2CO_3 , dioxan- H_2O); vi, PhtNNa (DMF, 60 °C, N_2);⁴ vii, N_2H_4 (EtOH, 80 °C); viii, $\text{Na}(\text{CF}_3\text{COO})\text{BH}_3$ (THF)⁵.


Although it is a general method for polyamines and their analogues, it is relatively limited in scope when the protected precursors are desired.

C. Reduction of imines

The key step in this method is the formation of a secondary amine by reductive alkylation of the corresponding primary amine^{20,33}.



Several naturally occurring polyamine derivatives^{55,56} have been synthesized using this general procedure which can be illustrated with the preparation of the N⁸-Ac-spermidine **85** as shown in Scheme 10⁵⁵. Condensation of monoacetyl putrescine with the protected aldehyde **82** gave the Schiff base **83** which was reduced to the protected spermidine **84**. The final product was then obtained by a simple deprotection step. Virtually, starting from different N-monoprotected diaminoalkanes and N-protected amino aldehydes it is possible to prepare selectively protected polyamines which can be modified at any nitrogen atom.




Scheme 10 Synthesis of N⁸-Ac-spermidine. Reagents: i, ZCl; ii, 0.06 M HCl (aq. dioxan 1:1); iii, AcNH(CH₂)₄NH₂ (CH₂Cl₂, Na₂SO₄); iv, NaBH₄ (MeOH); v, H₂/10 % Pd-C (EtOH)⁵⁵.


The stability of the imine formed is of crucial importance to the success of this approach. The condensation-reduction process can be carried out in two ways: reduction of the isolated Schiff base or reduction of the formed imine in situ. Although in the literature⁵⁷ high yields have been found in a "one-pot" synthesis, some authors^{56,58} have reported low yields in the range of 20-50 %. This is probably due to further reaction of the formed secondary imine with aldehyde^{20,33} giving rise to tertiary amine^{58a,59}. The procedure involving intermediate isolation seems to be more efficient but it is limited to stable imines⁶⁰.

D. Reduction of azides

The azide group can be reduced to primary amines by catalytic reduction^{20,33,61} or to secondary amines by their reductive alkylation with borane derivatives⁶².



Recently, based on this feature, Carrié et al.⁶³, have developed a method for building the polyamine backbone as outlined in Scheme 11. Thus, the amino azides⁶⁴ could be reductively alkylated with the dichloroboranes of the type



$n, m = 2, 3; l = 3$ $R^1 = H, \text{ alkyl, acyl}; R^2 = H; R^3 = H, \text{ alkyl}$

Scheme 11 Synthesis of polyamines by reduction of azides. Reagents: i, $Br(CH_2)_mBCl_2$ (CH_2Cl_2 or C_6H_6); ii, $MeOH$; iii, NaN_3 (H_2O_2 , reflux); iv, $NaOH$; v, $H_2/Pd-C$; vi, R^3BCl_2 ; vii, $Br(CH_2)_lBCl_2$.

$Br(CH_2)_mBCl_2$ ⁶⁵ to the derivatives **86** after methanolysis. They were then transformed to the diamino azides **87** by nucleophilic substitution of bromide with NaN_3 . These compounds could be either hydrogenated or again alkylated by a suitable dichloroborane to give, respectively, the polyamines **88**, **89** or **90**. If desired, the triamine bearing a bromide atom, could again be treated in the same way to afford higher polyamines.

Although only few alkylated and acylated polyamine derivatives are reported⁶³, this approach seems to allow a flexible synthesis not only of polyamines themselves but also of their selectively modified analogues by using appropriately protected starting materials or protecting the intermediate amino azides or bromides.

1.2.3 - Direct selective protection and modification of polyamines

In this section will be described the second strategy for synthesis of polyamine analogues which is based on regioselective acylation and alkoxycarbonylation of primary amino groups or transiently protected polyamine systems masked either as cyclic ureas or as hexahydropyrimidines.

1.2.3.1 - Regioselective acylation and alkoxycarbonylation of primary amines

It is known that secondary amines are more basic and therefore generally more nucleophilic than primary ones towards electrophilic reagents. Nevertheless, the primary amino groups of polyamines such as spermidine show higher reactivity which is explained in terms of an intramolecular hydrogen bonding between the secondary nitrogen atom and the hydrogen of the aminopropyl moiety or steric factors when bulkier agents are employed^{66,68}. This important feature has

been exploited and recently several procedures have been reported where the primary amino groups are selectively functionalized without affecting the secondary ones using different selective reagents^{66,68-76}.


A. 2-Oxazolones

Kunieda et al.⁶⁶ have developed series of 3-acyl and 3-alkoxycarbonyl-2-oxazolones (RCO-ox) and the corresponding polymers as carbonyl transfer agents to different nucleophiles.



a, R = -CH₃; **b**, R = -OCH₂Ph; **c**, R = -OBu^t

The approach used either the ready-to-use type reagents RCOox^{66a-66d} or the corresponding carboxylic acids via diphenyl-2-oxo-3-oxazolinyolphosphonate (DPP-ox) as carboxyl-activating reagent^{66e,66f} as depicted in the following scheme.



(See ref. 124 for preparation of Hox)

Scheme 12 Aminolysis of DPP-ox^{66f}.

These reagents have shown several interesting features in the acylation of different amines and aminoalcohols^{66d,66e}. The selective N-acylation or N,O-diacylation of aminoalcohols could be accomplished in the absence or presence of a catalyst. Another characteristic of these reagents and more relevant to this study, is their reactivity in relation to amines and polyamines which leads to highly preferential acylation of less hindered amino functions. The steric effects were more pronounced with the bulkier polymeric reagent (Table III).

Table III - Acetylation by polymeric reagent **93a**^{66d}.


Nucleophile	yield %
PhCH ₂ NH ₂	93
PhNH ₂	10 (91) ^b
1-Adamantanemethylamine	79 (85) ^b
1-Adamantylamine	13 (41) ^b
Cyclohexylamine	61
Dicyclohexylamine	0
CH ₃ CH ₂ NHCH ₂ CH ₂ NH ₂	80 ^c
NH ₂ (CH ₂) ₃ NH(CH ₂) ₂ NH ₂	93 ^c

^a Reaction conditions: THF, 6 hr, room temperature.

^b Acetylation by monomeric **92a**.

^c Only the primary amino groups were acylated.


Thus, the authors^{66f} have reported the synthesis of the bioactive maytenine **94** by direct simple acylation of the primary amino groups of spermidine with 3-trans-cinnamoyl-2-oxazolone.




Scheme 13 Synthesis of maytenine. Reagents: i, PhCH=CHCO-ox (CH₃CN)^{66f}.

B. Thiazolidine-2-thiones

The 3-acyl- and 3-alkoxycarbonylthiazolidine-2-thione derivatives (RCO-TT)⁶⁷ **95** have been employed as coupling agents in peptide chemistry and as acylating reagents for amines and alcohols⁶⁸.




The general sequence of the aminolysis reaction is shown in Scheme 14 which has a particular characteristic of being monitored by the disappearance of the yellow colour of the starting material (RCO-TT).



Scheme 14 Aminolysis of 3-acyl-thiazolidine-2-thiones^{68a}.


As in the case of the 2-oxazolones described above, this reagent also exhibited selectivity in relation to several amines. This is, however, affected by the electron density at the nitrogen atom and the sterical environment⁶⁸. Thus, this research group has also reported the syntheses of the spermidine alkaloids maytenine^{68a,68i} and codonocarpine^{68c,68e} by direct acylation of the primary amino functions, and the spermidine siderophore parabactin^{68j} **100** (Scheme 15), by previous direct selective protection of the primary amino groups with Z.



Scheme 15 Synthesis of parabactin. Reagents: i, Z-TT (CH_2Cl_2); ii, phenyl-bis(2-thione-1,3-thiazolidinyl)phosphine oxide, $(\text{Pr}^i)_2\text{NET}$ (CH_3CN , N_2 , reflux); iii, 25 % HBr-AcOH; iv, 2,3-diacetoxybenzoyl Chloride, Et_3N (THF); v, K_2CO_3 (MeOH)^{68j}.

C. Imidazoles


In 1984 Scott et al.⁶⁹ reported the synthesis of several N^1, N^8 -diacylspermidine derivatives by direct selective acylation at the primary nitrogen atoms with acylimidazoles (RCO-Im).



R= aryl

(RCO-Im)
101


The general approach involves a two-step reaction: formation of the (RCO-Im) by treating the corresponding carboxylic acids with N, N' -carbonyldiimidazole (CDI) followed by the acylation step as illustrated in Scheme 16 for the synthesis of the natural siderophore **105**⁶⁹.



Scheme 16 Synthesis of N^1, N^8 -bis(2,3-hydroxybenzoyl)spermidine. Reagents: i, CDI (CH_2Cl_2); ii, spermidine; iii, $\text{H}_2/\text{Pd-C}$ (MeOH, 5% AcOH)⁶⁹.

Although the procedure is simple and efficient, it has certain limitations. This depends upon the high steric hindrance at the carbonyl group to the attack by the nucleophile. Thus, as the authors have reported⁶⁹, the selectivity was lost and mixtures of products were obtained in the case of cinnamic acid.

More recently, based on this method, Sharma et al.⁷⁰ have synthesized spermidine siderophores via primarily protected polyamine precursors which was accomplished with benzyloxycarbonylimidazole in the presence of catalytic amounts of DMAP. The protected spermidine could then be acylated at the secondary nitrogen atom using succinimide esters of various acids. Subsequent removal of the Z groups allowed further symmetrical modification at the primary amino groups to afford for example the siderophore **106**.




106
Spermaxatol


D. Benzo- and Naphthosultones

Acher et al.⁷¹ have synthesized different strained five-membered benzosultones **107** and naphthosultones **108** and studied their reactivity as coupling agents in peptide

chemistry and as acylating reagents for amines.



107




108

a: X=Y=H; b: X=NO₂ Y=H; c: X=Y=NO₂

Generally, the aminolysis of these aryl esters takes place according to a two-step process involving a tetrahedral intermediate as represented in Scheme 17. The first step, the nucleophilic attack by the carboxylate anion, leads to a mixed anhydride which is then transformed to an activated ester by an intramolecular rearrangement, the seven-membered oxygen \rightarrow oxygen acyl transfer. The second step is the aminolysis of the activated ester to give the acylated amine. This reaction is fast in aprotic solvents which is explained in terms of an anchimeric assistance by the neighbouring SO_3^- group via an intramolecular general base catalysis.

The benzosultones do not give quantitative yields of the activated esters and, consequently, of amides, due to an incomplete acyl transfer reaction. This equilibrium can favourably be shifted with the naphthosultones because these form more rigid mixed anhydrides. Although the nitro derivatives are more reactive with respect to the nucleophilic attack by carboxylate ions, the aminolysis of the dinitro esters is slower due to the steric hindrance of the *o*-nitro substituent.



Scheme 17 Aminolysis of benzosultones⁷¹.

The reaction of primary and secondary amines with these reagents has shown that the acylation of the latter was a much slower process than with the former ones and the activated esters of the benzosultones were more selective than those of the naphthosultones. Thus, these authors⁷¹ have also reported the synthesis of maytenine in 71 % yield by direct selective acylation of N¹ and N⁸ of spermidine with 5-nitrobenzosultone **107b** and trans-cinnamic acid.


E. 1-Hydroxypiperidine esters

The peptide-coupling agents, the active 1-hydroxy-piperidine esters **109**, have also been used as acylating reagents for amines⁷².



109

The acylation of amines is relatively fast and Young et al.^{72b} have suggested that the transition state for the formation of the adduct (I), and the adduct itself, may be stabilized by hydrogen bonding and the subsequent proton transfer would greatly accelerate the final step (Scheme 18).



Scheme 18 Acylation of amines with 1-acyloxypiperidine^{72b}.


These reagents have also shown selectivity. For instance, it was reported that the reaction rates of benzoylation of amines with 1-benzoyloxypiperidine decreased in the following general order: n-butylamine > PhCH₂NH₂ >> (CH₃)₂CHNH₂ >>> (CH₃)₃CNH₂^{72b}. The authors have interpreted these data in such a way that increasing steric effects might prevent the amines from approaching sufficiently close for hydrogen bonding to be effective.

Exploiting these features, Husson et al.⁷³ have proposed these active esters for direct selective modification of terminal amino groups of polyamines. Thus, this research group

has reported the synthesis of several primary diacyl-spermidine derivatives such as maytenine and other dihydroxybenzoyl derivatives by reaction of spermidine with the corresponding active esters^{73d}.

F. Nefkens's reagent


The Nefkens's reagent, N-ethoxycarbonylphthalimide **110**, has been used in peptide chemistry for the introduction of phthaloyl as an N-protective group²⁵.



110

Recently, Sosnovsky et al.⁷⁴ have reported selective protection of primary amino groups by using this reagent. Thus, the authors have described the syntheses of several N,N'-bisphthaloylated polyamines in yields varying between 53-86 %. As illustrated in Scheme 19 for the preparation of N⁴-Bz-spermidine, these protected precursors were then used for selective acylation at the secondary nitrogen atom with acyl chlorides followed by selective deblocking of primary amino groups by hydrazinolysis.

Although this approach for selective protection is a simple one-step procedure based on readily available starting material, the main drawback is the strong conditions required for




Scheme 19 Synthesis of N⁴-benzoylspermidine. Reagents: i, N-(EtOCO)-phthalimide; ii, BzCl (TEA, CH₂Cl₂); iii, N₂H₄ (EtOH, reflux)⁷⁴.


the removal of the phthaloyl group. This may be a limiting factor with respect to other functional groups present in the target molecule.

G. Nitriles

Two methods have been devised for acylation of amino groups with nitriles^{75,76}. One exploits the ruthenium-catalysed condensation of nitriles with amines in the presence of two equivalents of water at high temperature⁷⁵.



The second and more recent procedure⁷⁶ uses direct acylation of amines with acyl cyanides⁷⁷ or cyanofornates⁷⁸.




As in the methods previously described, a selective differentiation between primary and secondary amino groups

has been achieved by acylation with nitriles^{75,76}. For example, N^1, N^8 - Ac_2 -spermidine and maytenine were synthesized by direct condensation of spermidine with the corresponding nitriles, acetonitrile and trans-cinnamitrile, in the presence of a Ru catalyst in good yield⁷⁵. It is worth emphasizing that the authors could also introduce selectively the Z group with benzyl cyanofornate (ZCN) to give the known diprotected precursor N^1, N^8 - Z_2 -spermidine⁷⁶.

The clean acyl cyanide reaction is preferable to the ruthenium-catalysed one because it proceeds under mild conditions, an important requirement for the synthesis of thermally unstable polyamines.

1.2.3.2 - "Transiently protected" polyamines

Ganem et al.^{18,79} have developed a method for selective modification of a polyamine which takes advantage of a temporarily protected polyamine either as a cyclic urea or as a hexahydropyrimidine. The methylene bridge between N^1 and N^4 of polyamines such as spermidine leads to a strain-free six-center cyclic derivative **114** in preference to a seven-membered structure corresponding to an N^4, N^8 -cyclization.




114

a, X= carbonyl
b, X= methylene

A. Cyclic ureas

The synthesis of cyclic urea **114a** is depicted in Scheme 20¹⁸. The first step involves exhaustive methoxycarbonylation of spermidine with methyl chloroformate. In refluxing alkaline media the resulting triprotected spermidine undergoes a preferential hydrolysis and decarboxylation of the terminal urethane groups followed by in situ cyclization via an intramolecular aminolysis between the free amino group of the propyl moiety and the urethane group at the secondary nitrogen atom.




Scheme 20 Formation of spermidine cyclic urea. Reagents: i, ClCO_2CH_3 ; ii, aq. $\text{Ba}(\text{OH})_2$ (reflux)¹⁸.

The removal of the carbonyl group can be carried out either by urea exchange or by reduction with LiAlH_4 . The first method consists in a transamination by heating the cyclic urea in a large excess of low boiling diamine such as 1,3-propanediamine, which forms the corresponding water-soluble urea

derivative. The second procedure is less practical because it leads to a mixture of spermidine and the N¹-methylated derivative difficult to separate.


These cyclic ureas can be extended to higher polyamines containing a 1,3-diaminopropyl moiety and in the spermine case give the corresponding bis urea **115**.



115

For spermine, however, this approach seems to be of less practical value since all nitrogen atoms are blocked to the most common electrophilic agents. On the other hand, in the cyclic spermidine derivative, the N⁸ or all three nitrogen atoms can be differentiated by exploiting selective reactions¹⁸.


Thus, the synthesis of native spermine **4** was reported where the cyclic urea was monocyanoethylated at N⁸ followed by selective reduction (borane) of the resulting nitrile. The last step consisted in the removal of the carbonyl group by urea exchange (Scheme 21)¹⁸. Although not reported in the literature, the intermediates **116** and **117** may be exploited as potential precursors for selective protection and modification of the different nitrogen atoms.



Scheme 21 Synthesis of spermine. Reagents: i, $\text{CH}_2=\text{CHCN}$ (C_6H_6); ii, $\text{BH}_3\text{-THF}$ (room temperature); iii, $\text{NH}_2(\text{CH}_2)_3\text{NH}_2$ (140°C).

B. Hexahydropyrimidines


As a complement and alternative to cyclic ureas, Ganem et al.^{18,79} have also prepared the corresponding hexahydropyrimidine **114b** by simple condensation reaction of the polyamine with aqueous formaldehyde.




The cleavage of this cyclic gem-diamine can be achieved by acid hydrolysis⁸⁰ or by ethyl hydrogen malonate and piperidine in refluxing ethanol (a Knoevenagel type reaction)⁷⁹. The latter method is more advantageous for acid-sensitive polyamine analogues.

As in the case of cyclic ureas, spermine also reacts with formaldehyde to give the corresponding bis(hexahydro-

pyrimidine) derivative **118**^{18,81}.



In both cases the originally secondary nitrogen atoms are protected against electrophilic reagents and therefore differentiated from the primary ones. The synthesis of terminally diacylated spermidine and spermine derivatives (maytenine and kukoamine A **119**) has been reported by direct acylation of **114b** and **118** with the corresponding acyl chlorides^{79,81}.




Kukoamine A

The same authors⁸² have also described the synthesis of N⁴-acylspermidine derivatives after performing a bis tert-butoxycarbonylation of the hexahydropyrimidine. Selective removal of the methylene bridge affords the diprotected N¹,N⁸-Boc₂-spermidine which can be selectively acylated on the secondary amino group.


Another important feature of the hexahydropyrimidine reported by Ganem^{80,83} is the possibility of differentiating between the secondary nitrogen atom and the primary one. For

selective modification at the secondary nitrogen, the described cases are based on the protection of the primary amino group in **114b** by generation of the Schiff base **120** (Scheme 22)⁸⁰ or by complexation with crown-ether as in the synthesis of N¹-Ac-spermidine⁸³. For the preparation of the N⁸-acetyl isomer, monoacylation was reported at the primary amino group by using a selective acetylating reagent⁸³.



Scheme 22 Synthesis of thermospermine. Reagents: i, PhCHO (C₆H₆, reflux); ii, CH₂=CH₂CN₀ (EtOH); iii, 2M HCl-MeOH (reflux); iv, CoCl₂·6H₂O, NaBH₄⁸⁰.

The synthesis of the cytotoxic spermidine metabolites **128** isolated from the coral *Sinularia Brougersma*⁸⁵ is an example of the usefulness of this approach for selective differentiation of the three nitrogen atoms. After starting from the cyclic urea it proceeds via the hexahydropyrimidine intermediate **125**, obtained by selective reduction of the carbonyl group (Scheme 23)⁸⁴.



Scheme 23 Synthesis of a cytotoxic spermidine derivative. Reagents: i, HCHO / 88 % HCO₂H; ii, LiAlH₄; iii, aq. Na₂CO₃, CH₂Cl₂; iv, HCO₂H, (heating); v, H₂/Pd-C (ethyl acetate)⁸².

In comparison with the cyclic ureas, the hexahydro-pyrimidines are preferable because they are more easily available, can be deprotected under milder conditions and be useful for diprimary modification of higher polyamines. The main limitation of this approach is that it is only applicable to polyamines containing the aminopropyl moiety.

1.2.4 - Conclusions

The different synthetic methods discussed in this chapter, available for total synthesis and selective protection and modification of polyamines, are summarized in Tables IVa and IVb for the spermidine case.

The methods involving total syntheses are general approaches affording the free polyamines as well as fully or partially protected derivatives. Among these procedures the nitrile approach developed by Bergeron appears to be the most versatile one with which it is possible to obtain protected polyamines in high yield. Moreover, mild conditions are normally employed for the removal afterwards of the remaining protecting groups. The main disadvantage of the total synthesis strategy is the often large number of steps required to afford the target compounds. Nevertheless, for synthetic targets with a new or unusual C,N backbone, total synthesis is the only alternative.

On the other hand, in the spermidine and spermine series, the use of selective protection and modification reagents or Ganem's spermidine-formaldehyde adduct (the hexahydro-pyrimidine **114b**) is generally advantageous. Symmetrical modification at the primary amino groups or selective functionalization of the secondary one can normally easily be accomplished.

Table IVa - Methods for synthesis of spermidine derivatives by a total synthesis approach with selected examples.

Method	Protected or modified spermidine (N°)	Overall yield % (steps)	Conditions for removal of protective groups	Ref.
Alkylation of tosylamides	$\text{PhtN}(\text{CH}_2)_3\text{NTos}(\text{CH}_2)_4\text{NHZ}$ 21	18 ^a (6)	NH_2NH_2 Na/NH_3 $\text{H}_2/\text{Pd}^3\text{-C}$	23
Alkylation of amines	$\text{PhtN}(\text{CH}_2)_3\text{NH}(\text{CH}_2)_4\text{NHZ}$ 36	30 ^b (2)	NH_2NH_2 $\text{H}_2/\text{Pd}^2\text{-C}$	29
	$\text{PhtN}(\text{CH}_2)_3\text{NBzl}(\text{CH}_2)_4\text{NPht}$ 40	43 ^c (2)		
Reduction of nitriles	$\text{NH}_2(\text{CH}_2)_3\text{NBoc}(\text{CH}_2)_4\text{NHBoc}$	49 ^d (2)	CF_3COOH	36
	$\text{NH}_2(\text{CH}_2)_3\text{NBzl}(\text{CH}_2)_4\text{NH}_2$ 53	36 ^c (3)	$\text{H}_2/\text{Pd-C}$	40
	$\text{BocNH}(\text{CH}_2)_3\text{NBzl}(\text{CH}_2)_4\text{NHCOCF}_3$ 61	60 ^c (6)	CF_3COOH $\text{H}_2/\text{Pd-C}$ $\text{K}_2\text{CO}_3/\text{CH}_3\text{OH}$	43
Reduction of amides	$\text{PhtN}(\text{CH}_2)_3\text{NBoc}(\text{CH}_2)_4\text{NHBoc}$ 76	69 ^e (4)	NH_2NH_2 CF_3COOH CF_3COOH	51a
	$\text{BocNH}(\text{CH}_2)_3\text{NH}(\text{CH}_2)_4\text{NHBoc}$ 79	49 ^e (4)		51b
Reduction of imines	$\text{ZNH}(\text{CH}_2)_3\text{NH}(\text{CH}_2)_4\text{NHAC}$ 84	36 ^f (4)	$\text{H}_2/\text{Pd-C}$ HCl or NH_2NH_2	55

- ^a From $\text{PhtN}(\text{CH}_2)_4\text{Br}$
^b From $\text{NH}_2(\text{CH}_2)_4\text{NH}_2$
^c From PhCH_2NH_2
^d From $\text{NC}(\text{CH}_2)_2\text{NH}(\text{CH}_2)_4\text{NH}_2$
^e From $\text{BocNH}(\text{CH}_2)_3\text{CO}_2\text{H}$
^f From $\text{NH}_2(\text{CH}_2)_2\text{CH}(\text{OEt})_2$

Table IVb - Some examples illustrating the selective acylation methods described in 1.2.3.


Reagent	Protected or modified spermidine	Yield %	Conditions for removal of protective groups	Ref.
2-Oxazolones	Maytenine 94	76	-	66f
Thiazolidine-2-thiones	Maytenine 94	79	-	68a
	ZNH(CH ₂) ₃ NH(CH ₂) ₄ NH _Z 96	69	H ₂ /Pd-C	68j
Imidazoles	ZNH(CH ₂) ₃ NH(CH ₂) ₄ NH _Z 96	76	H ₂ /Pd-C	70
Benzo-sultones	Maytenine 94	71	-	71
1-Hydroxypiperidine esters	Maytenine 94	62	-	73c
Nefkens's reagent	PhtN(CH ₂) ₃ NH(CH ₂) ₄ NPht 111	75	NH ₂ NH ₂	74
Nitriles	Maytenine 94	70	-	75
	ZNH(CH ₂) ₃ NH(CH ₂) ₄ NH _Z 96	99	H ₂ /Pd-C	76
Ganem's adduct	Maytenine 94	85 ^a	-	79
	BocNH(CH ₂) ₃ NH(CH ₂) ₄ NHBoc	54 ^a	CF ₃ COOH	82

^aIn two steps.


2 - BACKGROUND OF THE PRESENT WORK

2.1 - Introduction

In general, acylations of amines and alcohols can be accomplished with the corresponding anhydride or chloride in pyridine⁸⁶. However, this method often fails in cases of electronically deactivated or sterically hindered substrates. As recently reviewed⁸⁷⁻⁸⁹, certain 4-dialkylaminopyridines, DMAP **129** and MPP **130**, are nowadays commonly used as catalysts to facilitate such difficult acylations.



129
DMAP



130
MPP

Usually only small amounts of the catalyst (typical molar ratio substrate/DMAP 20:1) are required for an efficient acylation. Moreover, in more difficult cases it is often necessary to use at least an equimolar amount of an auxiliary base to remove the acid formed in the reaction. Alternatively, it is possible to use a stoichiometric amount of dialkylaminopyridine to act as both base and catalyst (Table V)⁸⁸.

Although some DMAP-catalysed reactions such as the formation of urethanes from alcohols and phenylisocyanates can involve general base catalysis⁹⁰, most of them probably


Table V - Acetylation of 1-methylcyclohexanol (100 mmol) with acetic anhydride⁸⁸.

Catalyst (mmol)	Base (150 mmol)	Reaction time /h	Yield of product (%)
-	Pyridine	16	0
-	TEA	15	0
DMAP (4.0)	Pyridine	18	66
DMAP (4.0)	TEA	17	89
DMAP (1.0)	TEA	16	63
DMAP (4.0)	a	17	39
DMAP (100.0)	a	18	84
MPP	TEA	17	90

^aMethylene chloride was used as solvent.

occur by nucleophilic catalysis via an N-acylpyridinium ion as acylating agent (Scheme 24)⁹¹.

It is important pointing out that for DMAP the equilibrium is more shifted to the formation of the N-acyl pyridinium intermediate than for unsubstituted pyridine. This is due to the electron-donating mesomeric effects of the 4-dialkylamino group (see resonance structure II). The formed adducts, which exist as ion pairs in non-polar solvents, are also more stable and in some cases they have been isolated^{88,92-94}. Thus, their relatively high concentrations facilitate the second step which is subjected to a general base catalysis by the counter ion. Therefore the reactivity of these adducts depends upon the nature of the anion, and in general, acid anhydrides (e.g. X= acetate) are stronger acylating agents than the



Scheme 24 Mechanism of DMAP-catalysed acylation of amino groups.


corresponding acid chlorides ($\text{X} = \text{Cl}$). This second step is also promoted by the presence of strong bases.

Thus, the ability of these reagents to catalyse acylations depends not only on their strongly basic character. They might be considered as "catalytic supports"⁸⁸ for an electrophile promoting reaction with a substrate.

2.2 - tert-Butoxycarbonylation of amide type functional groups

In 1983 Grieco et al.⁹⁵ reported for the first time the DMAP-catalysed tert-butoxycarbonylation of secondary amides and lactams with Boc_2O ⁹⁶. Nevertheless, it was not until recently that Grehn, Gunnarsson and Ragnarsson⁹⁷ began to explore a general procedure for exhaustive tert-butoxycarbonylation of various type of amides using the Boc_2O /DMAP approach.

Although the first authors used an equimolar amount of DMAP and CH_2Cl_2 as solvent⁹⁵, the formation of the Boc_2O /DMAP adduct turned out to be faster in CH_3CN and only catalytic amounts of DMAP were required (0.05-0.1 equivalent)⁹⁷.



for R, R^1 see Table VI

As shown in Table VI, this acylation reaction has proved to be a very efficient procedure and useful for the conversion of different compounds to the corresponding Boc analogues. However, as pointed out by the authors⁹⁷ and already observed by other research groups^{87-89,95}, steric factors are important. Due to the bulkiness of the activated intermediate when using sterically hindered substrates, long reaction times and an excess of reagent are required (see Table VI, entry 17). In one case, for pivalanilide, the

Table VI - tert-Butoxycarbonylation of amides R¹NHR with Boc₂O/DMAP⁹⁷.

Entry	Compound		Reaction time /h	Yield (%)
	R	R ¹		
1	Ph	HCO	1	92
2	Ph	CH ₃ CO	8	99
3	Ph	PhCO	4	92
4	Ph	PhCH ₂ OCO	20	95
5	Ph	Boc	3	96
6	PhCH ₂	CH ₃ CO	20	86
7	Ph(CH ₂) ₂	CH ₃ CO	20	97
8	PhCH ₂	Boc	48 ^a	100
9	Ph(CH ₂) ₂	Boc	50	94
10	4-EtOCOC ₆ H ₄	CH ₃ CO	2	85
11	4-NO ₂ -2-CF ₃ -C ₆ H ₃	CH ₃ CO	1	83
12	Ph	4-Me-C ₆ H ₄ -SO ₂	1	94
13	Ph	2-NO ₂ -C ₆ H ₄ -S ₂	1	96
14	Ph	Ph ₂ P(=O) ₆ H ₄	1	98
15	4-Bu ^t -C ₆ H ₄	CH ₃ CO	6	98
16	3-Bu ^t -C ₆ H ₄	CH ₃ CO	20	98
17	2-Bu ^t -C ₆ H ₄	CH ₃ CO	50	97

^a Prepared in one step from benzylamine and excess of Boc₂O.

desired product was not obtained.

This acylation reaction is by no means restricted to carboxamides. The *p*-toluenesulfonanilide and *o*-nitrobenzenesulfenanilide as well as the diphenylphosphinanilide smoothly afford the corresponding Boc analogues (see Table VI, entries 12-14)⁹⁷.

Furthermore, with reference to the scope of this reaction, in the presence of Boc₂O and DMAP, urethane NH groups generally undergo the same substitution as described above for amides. The products are formed in high yields and are of

normal stability⁹⁷. Only few substances of this general type, systematically named imidodicarbonates, had earlier been prepared.

Various urethanes play an important role in synthetic organic chemistry for the protection of amino functions²⁵. Judging from this aspect, it is obvious that the chemistry described in the preceding paragraph can be exploited in different ways for double protection of primary amines.

In attempts to extend this approach to other dicarbonate reagents, the stability of the latter to DMAP appeared to be a limiting factor. While Boc_2O forms a rather stable adduct with DMAP, other dicarbonates such as dimethyl dicarbonate and Z_2O decompose rapidly in the presence of DMAP⁹⁷.


2.3 - Selective cleavage of amides

In general, the cleavage of amides requires strong reaction conditions²⁵ which practically excludes the use of amides as protecting groups.

Grieco et al.⁹⁵ have reported a milder method in which the N-Boc derivatives of secondary amides and lactams are selectively hydrolyzed by LiOH or methanolized in the presence of NaOMe at the less hindered carbonyl group.

Using a related approach, Grehn, Gunnarsson and Ragnarsson^{98,99} introduced a novel mild procedure in which the previous N-tert-butoxycarbonylated substrates undergo a selec-

tive aminolysis or similar base-catalysed methanolysis to give the corresponding acid-labile tert-butyl carbamates.



The scope of this cleavage reaction is indicated in Table VII. In general, the rate of DEAEA-mediated aminolysis is enhanced by electron-withdrawing substituents and decreased for sterically hindered substrates. On the other hand, in the case of compounds containing an aliphatic amide moiety the cleavage proceeds remarkably slowly (see Table VII, entries 7, 8). However, the rate of deacylation can be enhanced by carrying out the reaction in methanol and in the presence of a strong base such as TMG. As the selectivity is retained, this can be an efficient alternative to aminolysis for resistant substrates which do not contain base-labile functions.

As reported by the authors⁹⁹, this procedure is not applicable to substances carrying non-carboxamide groups of tosyl (Tos), 2-nitrophenylsulfenyl (Nps) and diphenylphosphinyl (Dpp) type. The first one is reconverted to Tos-anilide and the latter are almost unchanged even after prolonged reaction times.

The Boc-derivatives of suitable urethanes mentioned in the preceding section, "double-protected amines", can be brought to undergo cleavage to the corresponding Bu^t carbamates provided the first urethane-protecting group is orthogonal to Boc. In the early work on aminolytic cleavage of

Table VII - Selective deacetylation of $\text{CH}_3\text{CON}(\text{Boc})\text{R}^{99}$.

Entry	R	Reaction conditions	Time /h	Yield (%)
1	Ph	DEAEA	24	90
		TMG/MeOH	0.2	98
2	4-Bu ^t -C ₆ H ₄	NH ₂ NH ₂	1	100
3	4-EtO ₂ C	DEAEA	2	96
4	2-Et-C ₆ H ₄	DEAEA	10	95
5	4-Bu ^t -2-NO ₂ -C ₆ H ₃	DEAEA	2	97
6	2-Thienyl	DEAEA	2	98
7	PhCH ₂	DEAEA	70	96
		TMG/MeOH	0.5	98
8	Ph(CH ₂) ₂	DEAEA	150	91
		NH ₂ NH ₂	1	98

DEAEA= 2-diethylaminoethylamine

TMG= N,N,N',N'-tetramethylguanidine

Boc-substituted amides, there was evidence for partially selective aminolytic cleavage of Z-groups from the compounds of benzyl tert-butyl imidodicarbonate type⁹⁹.


In conclusion, the novel chemistry outlined in this chapter based on exhaustive tert-butoxycarbonylation of amides and urethanes and subsequent selective deacylation seemed a promising basis for the development of new strategies of synthesis and selective protection of amines such as polyamines which will be the topic of the following chapters.

3 - OBJECTIVES AND SYNTHETIC METHODOLOGY

As previously reviewed, several methods are now available for selective modification of polyamines but there seems to exist no ideal general approach particularly to accomplish full protection of such substrates. In general many steps are required and/or strong conditions for the removal of the protecting groups afterwards. Therefore, an alternative procedure to the reported ones seemed desirable.

The main aim of this project was to explore the possibility of accomplishing selective protection of spermidine by extending to this substrate the novel approach to the use of DMAP-catalysed tert-butoxycarbonylation, described by Ragnarsson et al.⁹⁷⁻⁹⁹, in connection with tactics of protection in simple monoamine compounds.

Thus, the first part of this work consisted of the syntheses of protected spermidine derivatives with application of exhaustive tert-butoxycarbonylation of the previously protected amino groups. As outlined before⁹⁷, several possibilities were available for the initial protection of these functions. In this context, however, it was necessary to choose a protecting group which was orthogonal in relation to the Boc one. As the well-known benzyloxycarbonyl group (Z) fulfils this requirement and is cleaved under very mild conditions, we decided to use this group for temporary protection of the amino functions. Thus, Schemes 25 and 26 depict the synthetic strategy for the analogues




Scheme 25 Protection of spermidine. Reagents: i, ZCl (aq. Na_2CO_3); ii, Boc_2O , DMAP (CH_3CN); iii, $\text{H}_2/\text{Pd-C}$ (MeOH); iv, TMG (MeOH); v, Z_2O (CH_2Cl_2).


N^1, N^8 -bis(tert-butoxycarbonyl)spermidine **133** and N^1 -benzyl-oxycarbonyl- N^8 -tert-butoxycarbonylspermidine **138**. In the case of the protected derivative **133**, the novel methodology would be directly applied to spermidine itself. In the synthesis of **138**, the new approach would be accomplished starting with the spermidine-formaldehyde adduct **114b**¹⁸.

Once these protected precursors were obtained, the analogue **133** would be suitable for selective modification at the secondary nitrogen atom. The derivative **138**, by appropriate protection at the secondary amino group, would be advantageous to selective modification on N^1 or N^8 . Thus, the second part of this work aimed at studying their scope for synthetic work leading to selectively modified analogues, mainly those of biological interest. Then we decided to synthesize the monoacetylated spermidine analogues which are of importance as metabolites and excretory products⁷. The synthetic routes leading to all monoacetylated derivatives, **144**, **148** and **152**, are outlined in Schemes 27 and 28.

The ethyl analogues of spermidine are interesting substrates from a biological point of view⁹. Surprisingly their prior preparation seems to have been overlooked in the chemical literature where only scarce details regarding their synthesis and properties are available. Therefore, our goal was to prepare them by applying suitable alkylating methods to the previously protected precursors. The results of the attempted syntheses will be discussed in the next chapter.




Scheme 27 Synthesis of N⁴-Ac-spermidine. Reagents: i, Ac₂O; ii, Boc₂O, DMAP (CH₃CN); iii, TMG (MeOH); iv, 2.29 M HCl₂ in dioxan.



Scheme 28 Synthesis of N¹-Ac- and N⁸-Ac-spermidine derivatives. Reagents: i, Boc₂O (CH₂Cl₂); ii, H₂/Pd-C (MeOH); iii, Ac₂O (TEA, CH₂Cl₂); iv, 2.29 M HCl in dioxan; v, Z₂O (CH₂Cl₂).

Suitable protected NH_3 derivatives, such as Boc_2NH and BocNHZ , prepared in our laboratory^{100,101}, are potential reagents for a direct synthesis of protected amines¹⁰¹ via either the Gabriel¹⁰² or Mitsunobu¹⁰³ reactions using halides or alcohols, respectively. Being potentially useful also for synthesis of polyamines, a third part of the project was related to the preparation of several new similar reagents. The imidodicarbonates mentioned were prepared by using the Boc_2O /DMAP-mediated reaction with suitable substrates^{100,101}. As this approach cannot be used with anhydrides others than Boc_2O ⁹⁹, it was necessary to devise an alternative procedure. For the proposed alkyl benzyl imidodicarbonates **155** the chosen strategy, outlined in Scheme 29, was based on the well-known reaction between isocyanates and alcohols as reported recently by Kocovsky¹⁰⁴.



Scheme 29 Synthesis of alkyl benzyl imidodicarbonates. Reagents: i, $(\text{COCl})_2$ (CH_2Cl_2); ii, ROH (CH_2Cl_2).


4 - RESULTS AND CONCLUSIONS

4.1 - Selective protection of spermidine


4.1.1 - Synthesis of N¹,N⁸-bis(tert-butoxycarbonyl)spermidine

As shown in Scheme 25 (p. 61), the synthesis of N¹,N⁸-Boc₂-spermidine was based on a three-step sequence starting from spermidine.


The preparation of the N¹,N⁴,N⁸-Z₃-spermidine **131** was readily accomplished by using a slight excess of ZCl in aqueous Na₂CO₃-dioxan to give the crude product as an oil slightly contaminated with benzyl alcohol. The pure compound could then be obtained in high yield after a simple column chromatography.



The key reaction, the exhaustive tert-butoxycarbonylation of the terminal urethane groups in compound **131**, using the DMAP-catalysed reaction⁹⁷, also proceeded essentially quantitatively although, to complete the reaction, it was necessary to add a slight excess of the acylating reagent. Compound **132** was also obtained as an oil which could be purified by column chromatography.



The exhaustively protected derivative **132** could be converted either to product **133** by the removal of all Z groups or to derivative **134** by the selective cleavage of the terminal Z groups.



Thus, the preparation of the diprotected spermidine analogue **133** took advantage of the known orthogonality of the Z/Boc protecting groups²⁵. The reaction was easily accomplished by catalytic transfer hydrogenolysis¹⁰⁵ according to a procedure reported by Spatola et al.¹⁰⁶. The hydrogenolysis of compound **132** using ammonium formate as hydrogen donor in the presence of Pd-C in aqueous acetic acid gave the desired product **133**, in this case as a white solid in good yield after recrystallization.

A more important finding was that compound **132** could be debenzoyloxycarbonylated, essentially selectively, on the originally primary amino groups by the TMG-catalysed methanolysis⁹⁹ to afford compound **134** also as an oil in high yield after chromatography. It is worth mentioning

that ^1H n.m.r. of crude **134** indicated the presence of only trace amounts ($< 1\%$) of anomalous cleavage products. This high degree of selectivity was somehow unexpected. In earlier experiments was reported a rather low selectivity in the base-catalysed methanolysis of compound $\text{Z}(\text{Boc})\text{NPh}$ (product ratio $\text{BocNHPH} : \text{ZNHPH} \approx 6$) which was explained in terms of a relative similarity between the Z and Boc groups⁹⁹. This cleavage reaction is indeed the key step of the synthesis of the selectively protected spermidine derivative **138** described below.

Compound **134** was also obtained from the diprotected spermidine **133** by simple benzyloxycarbonylation on the secondary amino group.

The yields in the different synthetic pathways aiming at the "symmetrical", diprotected spermidine analogue **133** are summarized in Table VIII.

Table VIII - Yields and melting points of the N^1, N^8 - Boc_2 -spermidine **133**.

N°	Compound ^a	Yield ^b (%)	m.p. /°C	Thesis, page
131	$\text{N}^1, \text{N}^4, \text{N}^8\text{-Z}_3\text{-spd}$	84	38-40	113
132	$\text{N}^1, \text{N}^4, \text{N}^8\text{-Z}_3\text{-N}^1, \text{N}^8\text{-Boc}_2\text{-spd}$	92	Oil	114
133	$\text{N}^1, \text{N}^8\text{-Boc}_2\text{-spd}$	80	85.5-86.5 ^c	116
134	$\text{N}^4\text{-Z-N}^1, \text{N}^8\text{-Boc}_2\text{-spd}$	80 ^d 77 ^e	Oil	117 118


^aCharacterized by ^1H and ^{13}C n.m.r. spectra and elemental analysis (the latter only for **133**). ^bAfter purification.
^cLit. 82 79-80 °C. ^dFrom **132**. ^eFrom **133**.

4.1.2 - Synthesis of N¹-benzyloxycarbonyl-N⁸-tert-butoxy-carbonylspermidine


As shown in Scheme 26 (p. 62), the starting material of this synthesis was the cyclic spermidine derivative **114b**, easily prepared in quantitative yield from the triamine and a fresh formaldehyde solution according to the reported procedure^{80b}.

As in the previous case, the general strategy worked satisfactorily and only few minor modifications were made.


The attempted benzyloxycarbonylation of the cyclic spermidine **114b** with ZCl gave a rather intractable mixture which decomposed further on performing column chromatography on silica. The inspection of the ¹H n.m.r. spectrum of the crude mixture did not show any correlation with the structure expected. This outcome was probably due to the instability of the cyclic aminal derivative under these reaction conditions¹⁰⁷. This procedure was therefore abandoned and dibenzyloxycarbonylation was smoothly accomplished with Z₂O¹⁰⁸ under anhydrous conditions. The product **135** contaminated with benzyl alcohol was readily purified by column chromatography.



No problems were encountered in the DMAP-catalysed tert-butoxycarbonylation⁹⁷ step which gave product **136** in high yield after chromatography.



The synthesis of $\text{N}^1\text{-Z-N}^8\text{-Boc-spermidine}$ **138** could in principle be achieved by first selective removal of the methylene group followed by selective cleavage of the Z group on N^8 or by these procedures in the reversed order.



On pursuing the former synthetic pathway, compound **136** being treated with potassium monoethyl malonate and pyridine, no cleavage of the methylene group occurred. Instead, the ^1H n.m.r. spectrum of the isolated crude product showed that the Z group on N^8 had been selectively cleaved. The model compounds $\text{N}^1, \text{N}^8\text{-Boc}_2\text{-N}^1, \text{N}^4\text{-methylenspermidine}$ ⁸² and $\text{N}^1, \text{N}^2\text{-Z}_2\text{-N}^2\text{-Boc-N}^1\text{-Et-ethylenediamine}$ ^{109a,b} were therefore treated similarly. In the former case the starting material remained unchanged and in the latter the Z group on N^2 was cleaved off


again to give N^1 -Z- N^1 -Et- N^2 -Boc-ethylenediamine^{109a,b} (these experiments are not described in the experimental section). However, the reaction worked nicely when substituting by malonic acid⁸² the potassium salt mentioned above.

Thus, although the final product N^1 -Z- N^8 -Boc-spermidine **138** could be obtained from compound **136** via derivative **139** by carrying out the cleavage of the methylene group with malonic acid followed by TMG-catalysed methanolysis, the yields were relatively low. In the light of the previous experiments, this outcome was presumably due to a partial cleavage of the N^8 -Z group on performing the Knoevenagel type reaction and during the laborious chromatographic procedures.

Nevertheless, good results were obtained via the alternative synthetic pathway. Thus again the key reaction, the TMG-mediated methanolysis, could easily be performed on compound **136** and the selective cleavage of the Z group on the acyclic moiety gave the compound **137** in good yield after chromatography. It is worth mentioning that, although longer reaction times were needed, this reaction could also be carried out with a smaller amount of TMG (0.5 eq.) than that originally used (1.5 eq.). No problems were encountered in the removal of the methylene bridge on compound **137** by performing the reaction with malonic acid. The final product **138** was obtained as a solid in quite good yield after chromatography.

The structure of this compound was confirmed by an independent synthesis in which the corresponding diprotected

amide **140** was reduced with NaBH_4 -TFA^{51b} to afford **138** in low yield.



The yields in the different synthetic pathways leading to the selectively protected spermidine analogue **138** are summarized in Table IX.

Table IX - Yields and melting points of the compounds isolated during the synthesis of N^8 -Boc- N^1 -Z-spermidine **138**.


N°	Compound ^a	Yield ^b (%)	m.p. /°C	Thesis, page
135	N^1, N^8 -Z ₂ - N^1, N^4 -(CH ₂)-spd	76	Oil	119
136	N^1, N^8 -Z ₂ - N^8 -Boc- N^1, N^4 -(CH ₂)-spd	90	Oil	120
137	N^1 -Z- N^8 -Boc- N^1, N^4 -(CH ₂)-spd	88	Oil	121
139	N^1, N^8 -Z ₂ - N^8 -Boc-spd	48	Oil	123
138	N^1 -Z- N^8 -Boc-spd	78 ^c 65 ^d	63-64	121 123

^aCharacterized by ¹H and ¹³C n.m.r. spectra and elemental analysis (the latter only for **138**). ^bAfter purification.
^cFrom **137**. ^dFrom **139**.

4.1.3 - Attempted synthesis of N⁸-benzyloxycarbonyl-N¹-tert-butoxycarbonylspermidine

As reviewed in Section 1.2.3.1, many reports have recently appeared in the literature on regioselective acylations of primary amino groups by various reagents. This feature tempted us to undertake some work aiming at the synthesis of N⁸-Z-N¹-Boc-spermidine **161** as an alternative to the previous isomeric substance **138**.


Thus, by using the new reagent ZCN⁷⁶, readily available from ZCl and KCN⁷⁸, a preliminary experiment was performed on a model compound (N-ethylethylenediamine) in a "one-pot" procedure according to Scheme 30.



Scheme 30 Synthesis of N²-Z-N¹-Boc-N¹-Et-ethylenediamine. Reagents: i, ZCN (CH₂Cl₂); ii, Boc₂O, NMM (CH₂Cl₂).

This reaction worked excellently and compound **157** was obtained in good yield as white crystals after recrystallization. ¹H and ¹³C n.m.r. spectra and t.l.c. (ether) of this compound differed from those of the other isomer Et(Z)N(CH₂)₂NHBoc^{109a,b}.

Disappointing results were, however, obtained when performing this reaction on the monocyclic spermidine derivative **114b**.




T.l.c. and ^1H n.m.r. of the crude product showed a rather complex mixture containing $\approx 60\%$ of the desired product, $\text{N}^8\text{-Z-N}^1\text{-Boc-N}^1, \text{N}^4\text{-methylenespermidine } \mathbf{160}$, together with considerable amounts of the $\text{N}^1, \text{N}^8\text{-Z}_2\text{-N}^1, \text{N}^4\text{-methylene-spermidine } \mathbf{135}$ as well as other impurities. This crude mixture was difficult to separate by column chromatography on silica and afforded only a modest yield of $\mathbf{160}$. Discouraging results were also obtained in an attempt to isolate the intermediate $\text{N}^8\text{-Z-N}^1, \text{N}^4\text{-methylenespermidine } \mathbf{159}$ before tert-butoxycarbonylation of the secondary amino group. After a laborious work-up only 5% of nearly pure $\mathbf{159}$ was obtained. These poor results of the selective acylation of derivative $\mathbf{114b}$ were presumably due to its instability to cyanide ions. Although not a strict proof, a t.l.c. experiment showed that when compound $\mathbf{114b}$ was treated with Et_4NCN , a rather complex mixture was obtained.

4.2 - Synthesis of modified spermidines

4.2.1 - Synthesis of N⁴-acetylspermidine dioxalate


As shown in Scheme 27 (p. 64), the compound N⁴-Ac-spermidine **144** could be prepared by using the Boc₂O/DMAP approach⁹⁷ or by direct acylation of the previously prepared intermediate N¹,N⁸-Boc₂-spermidine **133**.

No problems were encountered in the acetylation of spermidine with Ac₂O either in an aqueous or anhydrous reaction system.




Under the latter conditions a direct column chromatography of the reaction mixture was preferred in which the triacetylated derivative **141** was obtained in better yield.

Although a moderate decrease in the rate of the Boc₂O/DMAP-catalysed reaction of aliphatic substrates⁹⁷ was reported earlier, the tert-butoxycarbonylation of compound **141** turned out to be remarkably slow (\approx one week reaction) and several additions of Boc₂O were required to complete the reaction.




After a usual column chromatography on silica, the fully protected intermediate **142** was obtained in a satisfactory yield.

The next step was easily performed where compound **142** underwent selective deacetylation by the TMG-mediated methanolysis and the desired product **143** was again obtained in good yield after chromatography. In the alternative procedure, this compound was easily prepared by acetylation of the precursor **133** with Ac_2O in the presence of TEA.



The final step required the removal of the Boc groups and could be accomplished by different reagents²⁵. The acidolysis by HCl was chosen in order to obtain directly the compound as the dihydrochloride salt **144**.



Although the procedure worked well, the resulting product turned out to be rather hygroscopic. Therefore, it was converted to its oxalate salt **144a** by passing through an anion exchange column to afford the product as a white solid,

homogeneous by t.l.c. and h.p.l.c. Elemental analysis indicated that the oxalate salt **144a** contained two molecules of oxalic acid and half a molecule of crystal water.

Table X summarizes the results for the synthesis of N^4 -Ac-spermidine.

Table X - Yields and melting points of the compounds isolated during the synthesis of N^4 -Ac-spermidine salts **144**.


N°	Compound ^a	Yield ^b (%)	m.p. /°C	Thesis, page
141	N^1, N^4, N^8 -Ac ₃ -spd	78	oil	130
142	N^1, N^4, N^8 -Ac ₃ - N^1, N^8 -Boc ₂ -spd	68	oil	131
143	N^4 -Ac- N^1, N^8 -Boc ₂ -spd	88 ^c 74 ^d	oil	133 134
144	N^4 -Ac-spd.2HCl	92 ^e	hygroscopic	134
144a	N^4 -Ac-spd.2H ₂ C ₂ O ₄ .1/2H ₂ O	80 ^e	187.5-188.5	134

^aCharacterized by ¹H and ¹³C n.m.r. spectra and elemental analysis (the latter only for the oxalate salt **144a**). ^bAfter purification. ^cFrom **142**. ^dFrom **133**. ^eFrom the corresponding HCl salt.


4.2.2 - Synthesis of N¹-acetylspermidine dihydrochloride

The synthetic methodology (Scheme 28, p. 65) leading to this compound took advantage of the selectively protected precursor N¹-Z-N⁸-Boc-spermidine **138**.


This four-step protection-deprotection strategy was easily carried out and no problems arose. The first step was accomplished by tert-butoxycarbonylation at N⁴ with Boc₂O to give intermediate **145** in good yield after purification.




The removal of the Z group by hydrogenolysis following the standard procedure afforded compound **146** in high yield. The crude product was essentially pure (t.l.c. and n.m.r.) and could be used for the next step without further purification.



The acetylation of compound **146** was easily accomplished by performing this reaction under anhydrous conditions and subsequent column chromatography on silica of the reaction mixture afforded compound **147** in high yield.



The final product was obtained as the dihydrochloride salt **148** by removing the Boc groups with HCl.



Although slightly hygroscopic, the analytical specimen could be obtained by recrystallization to afford the product as a white solid, homogeneous by t.l.c. and h.p.l.c. and its physical data agreed with those previously reported^{35a,110}.

The results for the synthesis of N¹-Ac-spermidine are summarized in Table XI.

Table XI - Yields and melting points of the compounds isolated during the synthesis of N¹-Ac-spermidine dihydrochloride **148**.


N°	Compound ^a	Yield ^b (%)	m.p. /°C	Thesis, page
145	N ¹ -Z-N ⁴ ,N ⁸ -Boc ₂ -spd	87	oil	135
146	N ⁴ ,N ⁸ -Boc ₂ -spd	98	oil	136
147	N ¹ -Ac-N ⁴ ,N ⁸ -Boc ₂ -spd	95	oil	137
148	N ¹ -Ac-spd.2HCl	97	191-193 ^c	137

^aCharacterized by ¹H and ¹³C n.m.r. spectra and by elemental analysis (the latter only for the salt **148**). ^bAfter purification. ^cLit. 173-178 °C^{35a}, 189-191°C¹¹⁰.

4.2.3 - Synthesis of N⁸-acetylspermidine dihydrochloride


This compound was easily obtained starting from the same protected spermidine derivative **138** by an analogous four-step strategy (Scheme 28, p. 65).

In this case the secondary amino group of intermediate **138** was protected by the Z group. Thus, compound **149** was obtained in good yield by using as reagents either Z₂O¹⁰⁸ or ZOBt¹¹¹.




Although the results were similar, the latter reagent required longer reaction times probably due to steric factors¹¹¹.

The next reaction afforded intermediate **150** by the removal of the Boc group with HCl.




After a simple extraction step, the product was isolated essentially pure (t.l.c. and n.m.r.) and directly used for acetylation of the free amino group.

Again the acetylated product **151** was obtained in good yield by the same procedure as described in 4.2.2.



The Z groups were cleaved by hydrogenolysis to give the final product as an oil which was also converted to its dihydrochloride salt **152** in good yield.



Recrystallization of the crude material afforded a white powder, homogeneous by t.l.c. and h.p.l.c. Its physical data were also in agreement with those reported in the literature^{35a,110}.

The results for the synthesis of N⁸-Ac-spermidine are summarized in Table XII.


Table XII - Yields and melting points of the compounds isolated during the synthesis of N⁸-Ac-spermidine dihydrochloride **152**.

N ^o	Compound ^a	Yield ^b (%)	m.p. /°C	Thesis, page
149	N ⁸ -Boc-N ¹ ,N ⁴ -Z ₂ -spd	83 ^c 88 ^d	oil	138
150	N ¹ ,N ⁴ -Z ₂ -spd	96	oil	140
151	N ⁸ -Ac-N ¹ ,N ⁴ -Z ₂ -spd	85	oil	140
152	N ⁸ -Ac-spd.2HCl	90	202-203 ^e	141

^aCharacterized by ¹H and ¹³C n.m.r. spectra and elemental analysis (the latter only for the salt of **152**). ^bAfter purification. ^cWith ZOBT. ^dWith Z₂O. ^eLit. 204-205 °C^{35a}, 203.5-205 °C¹¹⁰.

4.2.4 - Synthesis of N¹,N⁸-diethylspermidine

Based on a reported procedure for N-alkylation of urethane type groups with alkyl halides²⁸, the N¹,N⁸-diethylspermidine trioxalate **163** was prepared from the tribenzyloxy-carbonylated intermediate **131** according to Scheme 31.



Scheme 31 Synthesis of N¹,N⁸-Et₂-spermidine. Reagents: i, NaH, EtI (THF-DMF, reflux); ii, H₂/Pd-C iii, oxalic acid.

The first step, the diethylation of compound **131** required a long reaction time and even so t.l.c. still showed traces of starting material as well as an extra spot, presumably the monoethylated derivatives. Column chromatography on silica of the crude mixture afforded compound **162** in a satisfactory yield. When the reaction was scaled-up to about 4 mmol, the yield of the pure product decreased to 40-50 %.

The final step, the hydrogenolysis of compound **162**, was readily accomplished using the standard procedure to give the N¹,N⁸-diethylspermidine in good yield, slightly contaminated with traces of impurities.

As the previous experiments with the acetyl derivatives

showed that their oxalate salts were conveniently handled, the diethylated analogue was converted to its oxalate salt **163** by treating an ethanolic solution of the spermidine derivative with a solution of oxalic acid in ether. As oxalic acid can form a divalent anion it was originally expected that only 1.5 equivalents of oxalic acid were needed. Elemental analysis indicated, however, that the salt formed contained three molecules of oxalic acid. Thus, for further experiments the salt was always precipitated with a slight excess of three equivalents of oxalic acid. Recrystallization gave a very pure salt **163** as shiny white crystals.

The results of this synthesis are summarized in Table XIII.

Table XIII - Yields and melting points of the compounds isolated during the synthesis of N^1, N^8 -Et₂-spermidine trioxalate **163**.

N°	Compound ^a	Yield ^b (%)	m.p. /°C	Thesis, page
162	N^1, N^4, N^8 -Z ₃ - N^1, N^8 -Et ₂ -spd	65	oil	144
163	N^1, N^8 -Et ₂ -spd N^1, N^8 -Et ₂ -spd.3H ₂ C ₂ O ₄	92	oil 229.5-230.0	145


^aCharacterized by ¹H and ¹³C n.m.r. spectra and elemental analysis (the latter only for the oxalate salt **163**). ^bAfter purification.

4.2.5 - Attempted syntheses of N¹-ethyl- and N⁸-ethyl-spermidine

The synthesis of the title compounds turned out to be a more difficult task than the corresponding to N¹,N⁸-diethyl derivative **163**. In principle it should be possible to make them from proper intermediates previously obtained in this research work by selective alkylation or reduction.

A. Alkylation experiments

In order to synthesize N¹-Et-spermidine **164** by the procedure used for the diethyl derivative **163** (Scheme 31, p. 83), N¹,N⁴,N⁸,N⁸-Boc₄-spermidine **166** was considered a potential intermediate. It should be possible to make **166** in two steps by exhaustive tert-butoxycarbonylation of the key compound **138**, followed by selective removal of the Z group, according to the following scheme:



Before performing the synthesis of derivative **166**, it was worthwhile to study the alkylation conditions using simple model compounds. Thus, several attempts were carried out and the results can be summarized as follows:


a) alkylation with EtI/NaH

The model compound $\text{Ph}(\text{CH}_2)_2\text{N}(\text{Boc})_2$ ⁹⁷ was treated with NaH and EtI in THF-DMF. Although the compound was stable for about 2 hours at room temperature, it was readily converted (≈ 2 h) to the mono-Boc derivative under refluxing conditions. Thus, the idea of ethylating **166** under these conditions was abandoned.


b) alkylation with EtI/Ag₂O

In the literature it was reported the N-methylation of N-Z and N-Boc amino acids with methyl iodide in the presence of silver oxide¹¹². The stability of the model compound mentioned in a) under these conditions prompted us to adopt this procedure.

Preliminary experiments with the model compound N²-Z-N¹-Boc-N¹-Et-ethylenediamine **157** only led to the recovery of the starting material even after long reaction times or heating. It is worth mentioning that methyl iodide also failed to alkylate compound **157** under these conditions.




Repeating this experiment with one of the reported amino acids, N-Z-alanine¹¹², showed that the reaction worked nicely with methyl iodide as alkylating reagent but failed with ethyl iodide. The latter reagent afforded only the ester of the N-Z amino acid.



c) alkylation with $\text{CF}_3\text{SO}_3\text{CH}_2\text{CH}_3$

The properties of the alkyl perfluoralkanesulfonic esters as highly reactive alkylating agents¹¹³ prompted us to perform the alkylation reaction using ethyl trifluoromethanesulfonate at room temperature.

The model experiments revealed several interesting features. No alkylation occurred when compound **157** was treated only with $\text{CF}_3\text{SO}_3\text{CH}_2\text{CH}_3$ (triflate) in CH_2Cl_2 according to a described procedure¹¹⁴. The presence of a base (NaH) was necessary for the alkylation of a urethane group and the desired product **167** (Scheme 32) was readily obtained (≈ 4 h) in quantitative yield when using two equivalents of NaH and triflate. The product was an oil and it agreed by t.l.c. and ^1H n.m.r. with a product obtained by alkylation with EtI/NaH.



Scheme 32 Synthesis of $\text{N}^2\text{-Z-N}^1\text{-Boc-N}^1, \text{N}^2\text{-Et}_2\text{-ethylene-diamine}$. Reagents: i, $\text{CF}_3\text{SO}_3\text{CH}_2\text{CH}_3$, NaH (CH_2Cl_2) or EtI, NaH (THF:DMF, reflux).

This result seemed promising for the synthesis of N^1 -Et-spermidine as also the compound $\text{Ph}(\text{CH}_2)_2\text{N}(\text{Boc})_2$ referred to in a) was stable to NaH for several hours at room temperature.

However, h.p.l.c. experiments using the model compound **157** and its isomer N^1 -Z- N^2 -Boc- N^1 -Et-ethylenediamine **158** surprisingly indicated that the alkylation of derivative **158** was considerably slower than that of **157** (Fig. 6).





Figure 6 Alkylation of N^2 -Z- N^1 -Boc- N^1 -Et-ethylenediamine **157** (a) and N^1 -Z- N^2 -Boc- N^1 -Et-ethylenediamine **158** (b) with 1 equiv. of $\text{CF}_3\text{SO}_3\text{CH}_2\text{CH}_3$ and 1 equiv. of NaH. (The data were obtained by semiquantitative h.p.l.c.).

While the half-life of compound **157** was about 2 h, only 15 % of the desired product **167** was obtained after 24 h following the alkylation of **158** and a by-product was also detected.

d) selective alkylation

In view of the results described in c), it seemed worthwhile to selectively N-alkylate a benzyl carbamate group in the presence of a tert-butyl one. Thus, it was performed an explorative experiment with N¹-Z-N⁴-Boc-putrescine **168**.



However, rather disappointing results were obtained. It was afforded the product **169** in 35 % (by h.p.l.c.) together with three by-products. Moreover, compound **169** was contaminated with one of the by-products which could not be separated either by column chromatography or h.p.l.c.

In summary, the synthesis of N¹-Et-spermidine by N-ethylation of a carbamate group of the intermediate N¹,N⁴,N⁸,N⁸-Boc₄-spermidine **166** seemed to be less practical. First, a compound containing an N,N-Boc₂ moiety was unstable to NaH in refluxing THF-DMF. Second, the tert-butyl carbamate appeared to have a low reactivity towards triflate in the presence of NaH at room temperature. Third, the selective



N-alkylation of the benzyl carbamate group in N¹-Z-N⁴-Boc-putrescine **168** gave a low yield of the impure product.

B. Reduction experiments

The next series of experiments attempted had as basic idea either the reductive alkylation of amino groups or the reduction of amide groups. The results can be summarized as follows:

a) reductive alkylation

The reductive alkylation of the adducts between the key intermediates **146** or **150** (Scheme 28, p. 65) and acetaldehyde with reduction of the formed imine by NaBH₄, seemed a promising method.




Preliminary experiments were performed with the model compound N¹-Boc-N¹-Et-ethylenediamine **170** and again the results were discouraging. When compound **170** was treated with excess of acetaldehyde in the presence of molecular sieves in benzene followed by isolation of the product, a highly insoluble and reddish material was obtained (probably

due to polymerization of the imine). The alternative way, the reduction in situ of the imine according to a procedure reported by Olsen et al.^{58b}, gave a mixture of starting material, the desired product and the dialkyl derivative, difficult to separate. In view of this outcome this procedure was also abandoned.

b) reduction of the amide group



The reduction of the acetyl derivatives to afford the corresponding ethyl derivatives was one possible alternative (Scheme 33).



Scheme 33 Reduction of an N-acetyl to an N-ethyl group. Reagents: i, NaBH₄, TFA (THF); ii, 2.29 M HCl (dioxan); iii, Red-Al (dioxan, reflux).

Selective reduction of the acetamide group with NaBH₄/TFA of the model compound N²-Ac-N¹-Boc-N¹-Et-ethylenediamine **171** afforded a modest yield (35 %) of N¹-Boc-N¹,N²-Et₂-ethylenediamine **172** after column chromatography. Another possibility, the reduction of the amide group with Red-Al¹¹⁵ after removing the urethane group, gave 52 % of the reduced compound **174**.


Thus, the final syntheses of the title compounds were performed by the action of Red-Al on the deprotected acetyl derivatives **148** and **152** (Scheme 28, p. 65) in refluxing dioxan.




In both cases t.l.c. of the crude mixtures showed a by-product with the same R_f as spermidine. This was probably due to a partial hydrolysis of the acetyl group. The work-up was rather laborious as the products were continuously extracted from the aqueous solution. Column chromatography of the crude products gave the compounds as white powders which were homogeneous by t.l.c. but contaminated with inorganic material (elemental analysis gave $\approx 30\%$ of the product). Thus, it was necessary to carry out another extraction followed by precipitation as oxalate salts. After recrystallization, very pure products were obtained as white shiny crystals in a rather modest yield.

4.3 - Synthesis of alkyl benzyl imidodicarbonates

As shown in Scheme 29 (p. 66), the starting material for the preparation of the alkyl benzyl imidodicarbonates **155a-i** was benzyl carbamate **153** which was easily obtained according to a standard procedure reported by Carter et al.¹¹⁶.




The key intermediate, benzyloxycarbonyl isocyanate **154**, was obtained in satisfactory yield from benzyl carbamate and oxalyl chloride according to a known general method^{117,118}.



This intermediate, a colourless liquid, turned out to be very sensitive to moisture, and decomposition to the initial starting material occurred readily. Thus, it should be handled with special precautions and stored below -20 °C in a sealed vessel under nitrogen.

The next step involved the reaction of isocyanate **154** with different alcohols¹⁰⁴ and no particular problems were encountered.



For R see Table XIV

In general the conversion **154** → **155** proceeded smoothly to give the corresponding alkyl benzyl imidodicarbonates in excellent yield and high purity after a simple work up. It is worth pointing out that, in the case of the extremely acid-sensitive PpocNHZ **155d**, the crude product was contaminated with significant amounts of benzyl carbamate **153**. This was probably due to a partial decomposition of **155d** in the presence of acidic impurities in **154**. Therefore, the isocyanate must be redistilled before the preparation of **155d**.

As mentioned before, these reagents are potential intermediates in amine synthesis¹⁰¹ using the Gabriel and Mitsunobu reactions. Thus, they should also be useful for designing new pathways to polyamines. However, their full usefulness in general practice remains to be explored.

By selective removal of one of the blocking groups in such doubly protected amines, subsequent alkylation of the urethane function should be possible. To confirm their usefulness, a preliminary study was performed employing these newly synthesized compounds using conventional deprotection conditions^{25,120}. All deblocking reactions displayed an excellent selectivity and in no case even traces of an anomalous deprotection product could be detected in the crude reaction mixtures as judged by t.l.c. and ¹H n.m.r. The yields observed were also satisfactory after extractive work-up.

The experimental results obtained in the syntheses and in the deprotection study of these imidodicarbonates are summarized in tables XIV and XV.

Table XIV - Yields and melting points of the alkyl benzyl imidodicarbonates ZNHR **155a-i**.

Compound ^a	R	Yield ^b (%)	m.p. ^c /°C (Lit.)
155a	Z	99	109-109.5 (105.5-106.5 ¹¹⁹)
155b	Z(MeO)	≈100	92.5-93
155c	Z(NO ₂)	97	113.5-114
155d	Ppoc	91	83.5-84
155e	Boc	84	oil ¹⁰¹
155f	Adoc	98	112-112.5
155g	Aloc	94	79.5-80
155h	Troc	92	90-90.5
155i	Fmoc	92	112-113

^aCharacterized by ¹H and ¹³C n.m.r. spectra and elemental analysis (the latter for **155b, c, d, f-i**).

Table XV - Selective deprotection of alkyl benzyl imidodicarbonates **155**.

N°	Compound	Product	Reaction conditions	Ref.
155b	Z(MeO)NHZ	ZNH ₂	TFA/anisole (9:1), 1 h, 0 °C	25
155c	Z(NO ₂)NHZ	Z(NO ₂)NH ₂	HF/anisole (9:1), 1 h, 0 °C or TFA, reflux, 15 min.	25
155d	PpocNHZ	ZNH ₂	1.5 % TFA/CH ₂ Cl ₂ , 1 h, r.t.	25
155f	AdocNHZ	AdocNH ₂	H ₂ /Pd-C, MeOH, 1 h, r.t.	25
155g	AlocNHZ	ZNH ₂	(Ph ₃ P) ₃ RhCl, 90 % aq. EtOH, 1 h, 70 °C	120b
155h	TrocNHZ	TrocNH ₂ ZNH ₂	H ₂ /Pd-C, MeOH, 1 h, r.t. Zn, AcOH, 4 h, r.t.	25, 120a
155i	FmocNHZ	ZNH ₂	20% Piperidine, DMF, 1 h, r.t.	25, 120c

4.4 - Concluding comments

The most important spermidine derivatives obtained during this research are listed in Table XVI. The following aspects of the syntheses should be emphasized:

1- The novel protection methodology for amino functions, which differentiates between primary and secondary amino groups, was successful when applied directly to spermidine and gave N^1, N^8 -Boc₂-spermidine 133 as a solid by a simple three-step procedure in quite a satisfactory yield. Therefore, this is a good alternative to the previously reported syntheses. Bergeron et al.⁴¹ prepared it in five steps from benzylamine. Das et al.^{51b} have also reported a five-step sequence from 4-Boc-aminobutyric acid and 3-amino-1-chloropropane. More recently, Ganem et al.⁸² took advantage of the cyclic spermidine derivative and also obtained a crystalline product in three steps from spermidine. However, for an extension to other polyamines, our approach seems to be of general applicability whereas the Ganem procedure is limited to those containing the aminopropyl moiety. Developments of selective acylation reagents, such as acyl cyanides recently reported⁷⁶, might lead to an efficient one-step alternative. Nevertheless, the access to those reagents is still a restricting factor.

2- A selective protection of spermidine could also be achieved in a four-step route by performing this novel approach on a cyclic spermidine analogue in good overall

Table XVI - Summary of major protected and modified spermidine derivatives.

N ^o	Compounds ^a	Overall yield ^b , % (steps)	m.p. /°C	Thesis, page
133	N ¹ ,N ⁸ -Boc ₂ -spd	62 (3 from spd)	85.5-86.5 ^c	113
138	N ¹ -Z-N ⁸ -Boc-spd	45 (5 from spd)	63-64	119
144a	N ⁴ -Ac-spd.2H ₂ C ₂ O ₄ . 1/2H ₂ O	43 (4 from spd)	187.5-188.5	130
148	N ¹ -Ac-spd.2HCl	78 (4 from 138)	191-193 ^d	135
152	N ⁸ -Ac-spd.2HCl	65 (4 from 138)	202-203 ^e	138
163	N ¹ ,N ⁸ -Et ₂ -spd. 3H ₂ C ₂ O ₄	50 (3 from spd)	229.5-230	144
164	N ¹ -Et-spd.3H ₂ C ₂ O ₄	20 (2 from N ¹ -Ac- -N ⁴ ,N ⁸ -Boc ₂ -spd)	218.5-219	146
165	N ⁸ -Et-spd.3H ₂ C ₂ O ₄	22 (2 from N ⁸ -Ac- -N ¹ ,N ⁴ ,-Z ₂ -spd)	212.5-213	148

^aCharacterized by ¹H and ¹³C n.m.r. spectra and elemental analysis. ^bFrom pure compounds. ^cLit.⁸² 79-80 °C. ^dLit.^{35a,110} 173-174 or 189-191 °C. ^eLit.^{35a,110} 204-205 °C.

yield. Although previously referred to by Borowsky et al.¹²¹ as an in situ intermediate, this is the first complete synthesis reported for compound N¹-Z-N⁸-Boc-spermidine **138**. Thus, it is an alternative and complement to other derivatives such as N⁸-Z-N¹-Pht-N⁴-Tos-spermidine²³ and N⁴-Bzl-N¹-Boc-N⁸-(CF₃CO)spermidine⁴³, which have been prepared for the selective modification of spermidine. The first one requires at least six steps for its preparation and, moreover, rather

drastic conditions for the removal of its protecting groups. The second one, although containing more versatile protecting groups, also requires five-six steps. N^1 -Z- N^8 -Boc-spermidine can be easily obtained with the novel procedure. It takes advantage of the properties of the well established N-protecting groups Z and Boc which require rather mild orthogonal deprotection conditions. This approach, however, is limited to polyamines containing the aminopropyl moiety.

3- Although selective acylation of the secondary amino groups can be readily accomplished from the N^1, N^8 -Boc₂-spermidine **133** and similar disubstituted precursors, the Boc₂O/DMAP approach described in this thesis offers an efficient four-step alternative of wider application.

4- Compound N^8 -Boc- N^1 -Z-spermidine **138** turned out to be a good and rather convenient substrate for selective acylation on N^1 and N^8 via the intermediates N^4, N^8 -Boc₂-spermidine **146** and N^1, N^4 -Z₂-spermidine **150** (Scheme 28, p. 65), respectively. Thus, the monoacetylated spermidine derivatives were obtained in higher yield and purity compared to the reported ones^{35,55,83}. Tabor et al.³⁵ have first used a simple direct acetylation of spermidine which gave product mixtures and a low yield of the desired product. An improved later synthesis by the same authors³⁵ was based on the nitrile reduction methodology starting from monoacetylated diamines, putrescine and 1,3-propylenediamine, and proper nitriles, acrylonitrile and 4-bromobutyronitrile respectively. Slaich et al.⁵⁵ also

reported the syntheses of N^1 - and N^8 -acetylspermidine derivatives by the imine reduction approach in a relatively low overall yield. Ganem et al.⁸³ took advantage of the selective acetylation of the cyclic spermidine **114b** but rather impure products in low yields were obtained.

5- Prior protection with urethane groups (Z or Boc) provides an efficient route for alkylation of amines and in this particular context it worked rather satisfactorily for N^1, N^4, N^8 -Z₃-spermidine **131**. This compound seems to be more convenient in comparison with N-tosyl-protected polyamines^{9k} which require drastic conditions for the removal of the tosyl groups and a more laborious work-up.

6- Although the monoethyl spermidine derivatives were formed in rather modest yields by the reduction of the corresponding acetylated intermediates, they were obtained in a very high purity.

In conclusion, the protected spermidine derivatives N^1, N^8 -Boc₂-spermidine **133** and N^1 -Z- N^8 -Boc-spermidine **138**, readily obtained by simple three- and five-step routes respectively, are good intermediates for the selective acylation of spermidine. Therefore they are potentially useful for the synthesis of naturally occurring acylated spermidine analogues such as alkaloids and siderophores. On the other hand, the first attempts for selective alkylation of some related spermidine intermediates were not entirely successful.

In this context, a strategy involving a total synthesis may be a better alternative. The Gabriel type reagents prepared, the alkyl benzyl imidodicarbonates $ZNHCOOR$ **155a-i**, might also be useful as starting material in such a synthetic approach, leading to new biologically important spermidine analogues.

5 - EXPERIMENTAL

The syntheses carried out during the course of this work are fully described here in the following order:

- 1) experiments mainly with N¹-ethylethylenediamine;
- 2) experiments with spermidine;
- 3) experiments related to the syntheses of the imidodicarbonates.

A list of the compounds isolated is presented at the end in order to facilitate the consultation of this experimental section.

Materials. General methods

All solvents applied as reaction media were of analytical grade and dried for several days over a molecular sieve (4A). The spermidine used in this work was obtained from Fluka AG (purum). The content of ZCl in the commercial samples was measured by ¹H n.m.r. and the quantities applied were corrected accordingly.

Unless otherwise stated, all organic extracts were repeatedly washed in turn with half their volumes of aqueous 1M KHSO₄, aqueous 1M NaHCO₃, and saturated aqueous NaCl, and then dried over anhydrous MgSO₄ (for amines Na₂SO₄).

T.l.c. analyses were performed on 0.25 mm² thick precoated silica plates (Merck DC-Fertigplatten Kieselgel 60 F₂₅₄) using the following solvent systems: (A) toluene-acetonitrile (2:1); (B), (C) light petroleum-ether (2:1), (1:3); (D), (E) CH₂Cl₂-ether (12:1), (20:1); (F), (G), (H), (K) CH₂Cl₂-acetone (2:1), (4:1), (9:1), (20:1); (L) CH₂Cl₂-acetone-HOAc² (5:5:1); (M) EtOAc-acetone-HOAc-water (5:3:1:1)²; (N), (O), (P) CH₂Cl₂-methanol (4:1), (9:1), (20:1); (Q) CH₂Cl₂-methanol²-HOAc (18:2:1); (R) CHCl₃-ethanol-water-aqueous 25% NH₃ (10:50:4:1) and (S) CHCl₃-methanol-aqueous 25% NH₃ (2:2:1)³. Spots were visualized by³ inspection under u.v. light at 254 nm or, after brief heating, by exposure to Cl₂ followed by dicarboxidine spray¹²² (violet-blue spots).

The analytical h.p.l.c. equipment consisted of two LDC Constametric pumps, an LDC gradient master, a Rheodyne 7125 injector, an LDC Spectromonitor III variable wavelength u.v.-detector, an Altex 400 mixer, and a Shimadzu C-R3A integrator.

Column chromatography was performed on silica gel (Merck,

Kieselgel 60, 70-230 mesh ASTM). The dimensions of the columns were 20 x 2.7 cm, 15 x 5 cm and 20 x 6 cm for 50 g, 160 g and 260 g of silica, respectively.

N.m.r. spectra were routinely recorded for solutions in CDCl_3 on a JEOL FX90Q instrument at 90 MHz (^1H) or 22.5 MHz (^{13}C). The chemical shifts are generally reported relatively to TMS as internal standard but for spectra recorded in D_2O , they refer to sodium 2,2-dimethyl-2-silapentane-5-sulfonate (DSS). The signals, assigned by comparing chemical shifts and peak shapes, are tentative.


Melting points were determined in open capillary tubes on a Gallenkamp melting point apparatus and are uncorrected.

Elemental analyses of selected derivatives were carried out by Mikro Kemi AB, Uppsala, Sweden.

5.1 - Experiments with model compounds

5.1.1 - Synthesis of N²-benzyloxycarbonyl-N¹-tert-butoxy-carbonyl-N¹-ethylethylenediamine (157)

One-pot procedure:




To a solution of N¹-ethylethylenediamine (4.41 g, 50.1 mmol) in dry CH₂Cl₂ (100 ml), ZCN⁶¹ (8.22 g, 51.1 mmol), dissolved in dry CH₂Cl₂ (50 ml), was added dropwise at r.t. with vigorous stirring over a period of 4 h. The resulting colourless solution was stirred for 2 h with the exclusion of atmospheric moisture. A solution of Boc₂O (12.0 g, 55.1 mmol) in dry CH₂Cl₂ (50 ml) was then slowly introduced (≈ 1 h) followed by N-methylmorpholine (5.57 g, 55.1 mmol). The reaction mixture was then stirred overnight. The solvent was removed at reduced pressure below 30 °C. The remaining, almost colourless oil was partitioned between ether (1000 ml) and aqueous 0.2M citric acid (500 ml). The ethereal extract was washed successively with aqueous 0.2M citric acid, aqueous 1M NaHCO₃ and saturated aqueous NaCl (3 x 250 ml) and dried (MgSO₄). Evaporation to complete dryness left a pale yellow viscous residue which soon solidified upon trituration with cold heptane. The crude, essentially pure product 157 was obtained in quantitative yield. It was recrystallized from

light petroleum (20 ml/g, decolourizing carbon). After seeding and chilling to $-20\text{ }^{\circ}\text{C}$ overnight, the precipitated white crystals were collected, rinsed with small portions of cold light petroleum and dried in high vacuo to afford 13.1 g (81 %) of the pure compound **157**. T.l.c. (ether) gave one spot, different from the isomeric $\text{N}^2\text{-Boc-N}^1\text{-ethyl-N}^1\text{-Z-ethylenedi-amine}^{109}$; m.p. $47.5\text{-}48.0\text{ }^{\circ}\text{C}$; δ_{H} 7.33 (s, 5H, arom. H), 5.35 (broad, \approx 1H, NH), 5.09 (s, 2H, CH_2Ph), 3.10-3.33 (complex, 6H, CH_2N), 1.44 [s, 9H, $\text{C}(\text{CH}_3)_3$], and 1.08 (t, 3H, CH_2CH_3); δ_{C} 156.5 and 155.9 (CO), 136.6, 128.4, and 128.0 (arom. C), 79.7 [$\text{C}(\text{CH}_3)_3$], 66.5 (CH_2Ph), 46.0 ($\text{CH}_3\text{CH}_2\text{N}$), 42.5 (CH_2NEt), 40.3 (CH_2NHZ), 28.4 [$\text{C}(\text{CH}_3)_3$], and 13.6 (CH_2CH_3). (Found: C, 63.4; H, 8.1; N, 8.7. $\text{C}_{17}\text{H}_{26}\text{N}_2\text{O}_4$ requires C, 63.3; H, 8.1; N, 8.7%).

5.1.2 - Synthesis of $\text{N}^1\text{-benzyloxycarbonyl-N}^2\text{-tert-butoxy-carbonyl-N}^1, \text{N}^2\text{-diethylethylenediamine}$ (**167**)


A - Alkylation with ethyl iodide



To a solution of compound **157** (500 mg, 1.55 mmol) and EtI (2.20 g, 14.1 mmol) in anhydrous THF:DMF (10:1) (25 ml), NaH (80 % dispersion in oil, 150 mg, 5.00 mmol) was cautiously added at $0\text{ }^{\circ}\text{C}$ with gentle stirring under nitrogen. The resulting suspension was refluxed under nitrogen for 24 h.

The reaction mixture was cooled to r.t. (a white precipitate was formed) and water (2 ml) was added to decompose excess of NaH. The clear yellow solution was evaporated under reduced pressure and the residue partitioned between ether (200 ml) and aqueous 1M KHSO₄ (100 ml). The organic layer was washed and dried as usual. Evaporation to dryness gave a yellow oily residue (498 mg, 92 %). This crude material was chromatographed on silica (light petroleum-ether, 2:1). The appropriate fractions were pooled and evaporated under reduced pressure to afford 364 mg (67 %) of compound **167** as a pale yellow oil, homogeneous by t.l.c. (B or G); δ_H 7.35 (s, 5H, arom. H), 5.13 (s, 2H, CH₂Ph), 3.21-3.44 (complex, 8H, CH₂N), 1.45 [s, 9H, C(CH₃)₃], and 1.05-1.25 (m, 6H, CH₂CH₃); δ_C 155.8 and 155.2 (CO), 136.7, 128.4, 127.9, and 127.7 (arom. C), 79.3 [C(CH₃)₃], 66.9 (CH₂Ph), 45.6, 45.4, 44.8, and 42.8 (CH₂N), 28.4 [C(CH₃)₃], 13.9 and 13.5 (CH₂CH₃).


B - Alkylation with ethyl trifluoromethanesulfonate



A stirred solution of compound **157** (100 mg, 0.320 mmol) in dry CH₂Cl₂ (1 ml) was treated with NaH (20 mg, 0.640 mmol) followed by CF₃SO₃Et (114 mg, 820 μ l, 0.640 mmol) at r.t. After 3 h stirring, t.l.c. (G) indicated complete reaction.

The solvent was evaporated under reduced pressure and the yellowish residue chromatographed as under procedure A to afford the pure product **167** (110 mg, 98 %). T.l.c. and ^1H and ^{13}C n.m.r. spectra were identical with those given above.

5.1.3 - Synthesis of N^1 -tert-butoxycarbonyl- N^1 -ethylethylene-diamine (**170**)




Compound **157** (6.00 g, 18.6 mmol) was dissolved in dry methanol (350 ml) and hydrogenolyzed (1 atm, r.t.) in the presence of Pd-C (5 %, 1.00 g).

When the starting material had been consumed (t.l.c. (Q)), the catalyst was filtered off and rinsed with methanol. The colourless filtrate was taken to dryness to leave a light yellow oily residue which was partitioned between aqueous 0.2M citric acid (100 ml) and ether (100 ml). The aqueous phase was again extracted with ether (2 x 100 ml). After making alkaline with solid NaOH (pH \approx 13), the aqueous phase was extracted with ether (5 x 200 ml). The combined organic phases were washed with saturated aqueous NaCl (2 x 100 ml) and dried (Na_2SO_4). The extract was carefully evaporated to dryness under reduced pressure (the product was somewhat volatile and excessive drying caused losses) to afford 3.18 g (91 %) of compound **170** as a light yellow oil. T.l.c. (M or Q)

of this crude material showed mainly one spot and was suitable for further synthetic work; δ_{H} 3.13-3.36 (complex, 4H, CH_2N), 2.82 (t, 2H, CH_2NH_2), 1.98 (s, 2H, amine NH), 1.46 [s, 9H, $\text{C}(\text{CH}_3)_3$], and 1.10 (t, 3H, CH_2CH_3); δ_{C} 155.7 (CO), 79.3 [$\text{C}(\text{CH}_3)_3$], 49.7, 42.3, 40.7 (CH_2N), 28.5 [$\text{C}(\text{CH}_3)_3$], and 13.7 (CH_2CH_3).

5.1.4 - Synthesis of N^2 -acetyl- N^1 -tert-butoxycarbonyl- N^1 -ethylethylenediamine (171)




A solution of Ac_2O (1.63 g, 16.0 mmol) was added dropwise to a stirred ice-cold solution of compound **170** (2.15 g, 13.3 mmol) and TEA (2.02 g, 20.0 mmol) in dry CH_2Cl_2 (20 ml). The clear, pale yellow mixture was stirred for 1 h in ice-bath and overnight at r.t. The solvent was evaporated under reduced pressure and the yellow oily residue chromatographed on silica (CH_2Cl_2 -acetone 4:1). The appropriate fractions were pooled and evaporated to afford 2.90 g (94 %) of compound **171** as a light yellow oil. This product solidified at -20°C to a low melting solid, homogeneous by t.l.c. (A, G, or O); $\delta_{\text{H}} \approx 6.62$ (broad, $\approx 1\text{H}$, amide NH), 3.12-3.38 (complex, 6H, CH_2N), 1.96 (s, 3H, CH_3CON), 1.47 [s, 9H, $\text{C}(\text{CH}_3)_3$], and 1.10 (t, 3H,

CH₂CH₃); δ_C 170.5 (CO), 79.6 [C(CH₃)₃], 45.6, 42.5 and 39.3 (CH₂N), 28.4 [C(CH₃)₃], 23.1 (CH₃CON), and 13.7 (CH₃CH₂N).

5.1.5 - Synthesis of N¹-tert-butoxycarbonyl-N¹,N²-diethyl-ethylenediamine (172)


A - Reduction of compound 171



The compound **171** (653 mg, 2.84 mmol) in dry THF (15 ml) was treated with NaBH₄-TFA, essentially as outlined in ref. 51b. After stirring for 30 h, water (5 ml) was added to the reaction mixture and the solvent evaporated under reduced pressure. The residue was treated with aqueous 0.2M citric acid (20 ml) and extracted with ether (3 x 20 ml). The aqueous phase was made alkaline with solid NaOH (pH ≈ 13) and extracted with ether (4x50 ml). The combined organic layers were washed with saturated NaCl (2 x 20 ml) and dried (Na₂SO₄). Evaporation of the extract afforded 377 mg (61 %) of a yellowish oil. The crude product was chromatographed on silica (CHCl₃-EtOH-water-25 % aq. NH₃, 100:50:4:1) to give a semisolid residue. This material was dissolved in CHCl₃ and some white material filtered off. The solvent was evaporated to dryness to afford 227 mg (37 %) of compound **172** as a pale yellow oil, essentially pure by t.l.c. (R); δ_H 3.13-3.37 (m,


4H, CH_2NBoc), 2.55-2.83 (complex, 4H, CH_2NH), 1.46 [s, 9H, $\text{C}(\text{CH}_3)_3$], 1.40 (broad, \approx 1H, amine NH), and 1.10 (t, 6H, CH_2CH_3); δ_{C} 155.6 (CO), 79.3 [$\text{C}(\text{CH}_3)_3$], 48.2 ($\text{CH}_3\text{CH}_2\text{NH}$), 46.7 ($\text{CH}_3\text{CH}_2\text{NBoc}$), 44.0 (CH_2NHET), 42.4 [$\text{CH}_2\text{N}(\text{Et})\text{Boc}$], 28.5 [$\text{C}(\text{CH}_3)_3$], 15.4 ($\text{CH}_3\text{CH}_2\text{NH}$), and 13.6 [$\text{CH}_3\text{CH}_2\text{N}(\text{Et})\text{Boc}$].

B - Hydrogenolysis of compound 167



Compound **167** (217 mg, 0.620 mmol) was hydrogenolyzed and worked up as described for compound **170** and the product chromatographed as under procedure A to give 107 mg (80 %) of product **172** as a light yellow oil. T.l.c. and ^1H and ^{13}C n.m.r. spectra were identical with those given above.

5.1.6 - Synthesis of N^2 -acetyl- N^1 -ethylethylenediamine (173)




TFA (20 ml) was added to a flask containing compound **171** (505 mg, 2.19 mmol) and the resulting solution was stirred for 20 min. at r.t. The solvent was then quickly evaporated under reduced pressure. The residue was taken up in methanol (10 ml) and concentrated (twice). The remaining oily residue was treated with aqueous 30 % K_2CO_3 (15 ml) and extracted with

CHCl₃ (4 x 50 ml). The combined organic layers were dried (Na₂SO₄) and evaporated under reduced pressure to afford 242 mg (85 %) of product **173** as a brownish oil. This crude material was essentially pure by t.l.c. (M or Q) and was used without further purification; $\delta_{\text{H}} \approx 6.94$ (broad, $\approx 1\text{H}$, amide NH), 3.35 (q, 2H, CH_2NHAc), 2.56-2.83 (complex, 4H, CH_2NHET), 1.98 (s, 3H, CH_3CON), 1.26 (broad, $\approx 1\text{H}$, amine NH), and 1.12 (t, 3H, $\text{CH}_3\text{CH}_2\text{N}$); δ_{C} 171.1 (CO), 48.4, 43.6 and 38.9 (CH_2N), 23.0 (CH_3CON), and 14.7 ($\text{CH}_3\text{CH}_2\text{N}$).

5.1.7 - Synthesis of N¹,N²-diethylethylenediamine dihydrochloride (174)


A - Reduction of compound 173



To an ice-cold solution of compound **173** (109 mg, 0.837 mmol) in dry toluene (10 ml) was cautiously introduced a solution of 3.5M Red-Al in toluene (1.40 ml, 5.02 mmol) with gentle stirring under nitrogen (strong evolution of H₂). The resulting solution was refluxed under nitrogen for 2 h. The coloured reaction mixture was cooled to room temperature and then in an ice-bath. Water (2 ml) was gently added to


decompose excess of Red-Al (strong evolution of H₂) and the solvent evaporated under reduced pressure. The remaining residue was treated with aqueous 15 % NaOH (1 ml) and water (5 ml) (pH ≈ 13) and stirred for 1 h. The yellowish turbid aqueous solution was saturated with NaCl and extracted with CHCl₃ (4 x 20 ml). The combined extracts were dried (Na₂SO₄) and carefully evaporated under reduced pressure. The remaining oil was precipitated as dihydrochloride salt with excess of a solution of 2.29M HCl in dioxan to afford 82 mg (52 %) of product **174**. An analytical sample was obtained by recrystallization from ethanol (100 ml/g); m.p. 262-264 °C; δ_H 3.34 (s, 4H, NCH₂CH₂N), 3.15 (q, 4H, CH₃CH₂N), and 1.31 (t, 6H, CH₃CH₂N); δ_C 46.3 and 45.2 (CH₂N), and 13.2 (CH₃CH₂N).

B - Acidolysis of compound 172



Compound **172** (100 mg, 0.462 mmol) was treated with 2.29M HCl in dioxan (5 ml) with stirring at r.t. for 4 h. The solvent was evaporated under reduced pressure and the white residue was suspended in dry ether (5 ml) and concentrated (twice). The white solid residue was triturated with cold dry ether (2 ml), filtered off and rinsed with small portions of cold dry ether (3 x 1 ml) and dried in vacuo. The yield of crude **174** was 58 mg (66 %) and the recrystallized sample was identical with the product prepared according to procedure A.

5.1.8 - Synthesis of N¹-benzyloxycarbonyl-N⁴-tert-butoxy-carbonylputrescine (168)




To an ice-cold solution of N¹-Z-putrescine^{17b} (2.10 g, 9.04 mmol) in dry CH₂Cl₂ (10 ml) was added dropwise with stirring a solution of Boc₂O (2.17 g, 9.94 mmol) in dry CH₂Cl₂ (10 ml). Evolution of gas started and at the end of the addition of Boc₂O a white precipitate formed. The resulting mixture was then stirred at r.t. and after few minutes a clear, yellow solution was obtained. T.l.c. (O) indicated complete reaction after 2 h. The solvent was removed and the residue partitioned between ether (500 ml) and aqueous 1M KHSO₄ (200 ml). The yellowish ethereal extract was washed and dried as described before. The extract was evaporated to dryness, leaving 2.89 g (99 %) of crude **168** as a white solid, essentially pure by t.l.c. [ether or (O)]. Recrystallization from ethyl acetate-light petroleum (1:2) (40 ml/g) gave the pure product **168** as a white solid; m.p. 101.5-102.0 °C (lit.¹²³ m.p. 124-126 °C); δ_H 7.34 (s, 5H, arom. H), 5.09 (s, 2H, CH₂Ph), ≈ 4.95 and ≈ 4.63 (broad, ≈ 2H, amide NH), 2.98-3.98 (m, 4H, CH₂N), and 1.49-1.56 (m) and 1.43 (s) [together 13H, CCH₂C and C(CH₃)₃]; δ_C 156.4 and 160.0 (CO), 136.6, 128.5, and 128.0 (arom. C), 79.1 [C(CH₃)₃], 66.6 (CH₂Ph), 40.7, 40.2, (CH₂N), 28.4 [C(CH₃)₃], and 27.3 (CCH₂C).

5.2- Experiments with spermidine

5.2.1- Synthesis of N¹,N⁸-bis(tert-butoxycarbonyl)spermidine


N¹,N⁴,N⁸-Tribenzyloxycarbonylspermidine (131)



An ice-cold solution of spermidine (1.50 g, 10.0 mmol) in aqueous 10 % Na₂CO₃-dioxan (4:3, 70 ml) was treated dropwise under vigorous stirring with ZCl (7.40 g, 35.0 mmol) dissolved in dioxan (30 ml). When the addition was complete (≈ 1 h), the resulting mixture was stirred for 1 h in ice-bath (a precipitation of NaCl occurred) and 16 h at r.t. Most of the dioxan was stripped off under reduced pressure below 30 °C and the remaining suspension partitioned between aqueous 1 M KHSO₄ (100 ml) and ether (200 ml). The colourless, ethereal extract was washed and dried according to the general procedure described earlier. The extract was filtered and evaporated in vacuo to complete dryness to afford a pale yellow oil (4.80 g, 90 %) which, although contaminated with benzyl alcohol and minor impurities, was suitable for further synthetic work. The pure product was readily obtained by column chromatography on silica gel. Benzyl alcohol was eluted with neat CH₂Cl₂ and further elution with CH₂Cl₂-acetone (9:1)

gave 4.60 g (84 %) of compound **131** as a pale yellow oil, homogeneous by t.l.c. (A or B). The product solidified after several weeks at -20 °C, m.p. 38-40 °C; δ_{H} 7.31 (s, 15H, arom. H), \approx 5.6 (broad, \approx 2H, NH), 5.08 and 5.06 (2s, 6H, CH_2Ph), 2.98-3.38 (m, 8H, CH_2N), and 1.48-1.71 (m, 6H, CCH_2C); δ_{C} 156.4 (CO), 136.0, 128.5, 128.0, 127.9, 127.4, and 126.9 (arom. C), 67.1 and 66.6 (CH_2Ph), 46.4 and 44.1 (CH_2NCH_2), 40.5 and 37.7 (CH_2NHZ), and 28.9, 28.2, and 25.5 (CCH_2C).


$\text{N}^1, \text{N}^4, \text{N}^8$ -Tribenzyloxycarbonyl- N^1, N^8 -bis(tert-butoxycarbonyl)spermidine (**132**)



A solution of **131** (6.00 g, 11.0 mmol) and DMAP (134 mg, 1.10 mmol) in dry acetonitrile (50 ml) was treated with Boc_2O (5.00 g, 23.1 mmol) in one portion with agitation at r.t. when evolution of carbon dioxide occurred. After stirring for 2 h, t.l.c. (H) showed that starting material as well as another compound (the mono Boc-derivatives!) still remained in the mixture. More Boc_2O (2.50 g, 11.0 mmol) was introduced and the brownish mixture was left overnight. A new t.l.c. showed that the starting material was consumed but still indicated the presence of the mono Boc-products. Thus, more Boc_2O

(2.50 g, 11.0 mmol) was added and after 3 h the reaction was complete (t.l.c. showed one main spot). The solvent was stripped off at r.t. and the brownish syrupy residue was partitioned between ether (300 ml) and aqueous 1M KHSO₄ (150 ml). The brownish aqueous phase was discarded and the yellow ethereal extract worked up as described. After treatment with decolourizing carbon, the extract was taken to dryness under reduced pressure, leaving a dark yellow oil (7.90 g, 96 %). The crude product was chromatographed on silica with CH₂Cl₂-ether (20:1) as eluant. The appropriate fractions were pooled and evaporated to afford 7.60 g (92 %) of compound **132** as a pale yellow oil, homogeneous by t.l.c. (A or H); δ_{H} 7.34 and 7.31 (two s, 15H, arom. H), 5.20 (s, 4H, BocNCO₂CH₂Ph), 5.10 (s, 2H, third CH₂Ph), 3.56-3.62 [m, 4H, CH₂N(Z)Boc], 3.16-3.28 (m, 4H, CH₂NCH₂), and 1.71-1.94 (m) and 1.45 (s) [together 24H, CCH₂C + C(CH₃)]; δ_{C} 155.9, 153.8, 153.6, 152.0, and 151.8 (CO), 136.8, 135.5, 128.5, 128.3, 128.2, 127.8, and 127.7 (arom. C), 82.8 and 82.7 [OC(CH₃)₃], 68.2 (BocNCO₂CH₂Ph), 66.9 (third CH₂Ph), 46.4, 46.1, and 44.4 (CH₂N), 27.9 [C(CH₃)₃], and 26.2 and 25.4 (CCH₂C).

N¹,N⁸-Bis(tert-butoxycarbonyl)spermidine (133)




To a solution of compound **132** (8.70 g, 11.6 mmol) in aqueous 80 % acetic acid (200 ml) was added ammonium formate (13.0 g, 200 mmol). When all had dissolved, Pd-C (5 %, 4.00 g) was added in small portions with stirring under nitrogen at r.t. After stirring for 1 h, the catalyst was filtered off, rinsed with aqueous 80 % HOAc and the colourless filtrate was concentrated under reduced pressure. After the addition of aqueous 30 % K₂CO₃ (250 ml) the product was extracted with ether (3 x 300 ml). The combined colourless organic layers were washed with saturated aqueous NaCl (2 x 50 ml) and dried (Na₂SO₄). The colourless extract was filtered and evaporated under reduced pressure to give a yellow oil which slowly solidified (3.42 g, 85 %). Recrystallization of the crude material from light petroleum (100 ml/g) afforded **133** as a white solid (2.9 g, 72 %), homogeneous by t.l.c. (M), m.p. 85.5-86.5 °C (Lit.,⁸² 79-80 °C); δ_H ≈ 5.2 and 4.8 (broad, ≈ 2H, amide NH), 3.20 (q, 4H, CH₂NHBoc), 2.67 (m, CH₂NCH₂), and 1.52-1.79 (m) and 1.44 (s, together 25H, CCH₂C, C(CH₃)₃, and amine NH); δ_C 156.1 and 156.0 (CO), 79.0 [OC(CH₃)₃], 49.4 and 47.7 (CH₂N), 40.5 and 39.2 (CH₂NHBoc),

29.8, 27.9, and 27.3 ($\text{C}\underline{\text{C}}\text{H}_2\text{C}$), and 28,5 [$\text{C}(\underline{\text{C}}\text{H}_3)_3$]. (Found: C, 59.0; H, 10.1; N, 12.2. $\text{C}_{17}\text{H}_{35}\text{N}_3\text{O}_4$ requires C, 59.1; H, 10.2; N, 12.2 %).

5.2.2 - Synthesis of N^4 -benzyloxycarbonyl- N^1, N^8 -bis(tert-butoxycarbonyl)spermidine (134)


Method A - Methanolysis of compound 132 with excess of TMG



Compound 132 (1.41 g, 1.90 mmol) was dissolved in dry methanol (15 ml) and treated with TMG (656 mg, 5.70 mmol) with rapid stirring at room temperature. After 20 h stirring, the solvent was evaporated and the yellowish residue partitioned between ether (100 ml) and aqueous 1M KHSO_4 (50 ml). The yellowish extract was washed as usual and taken to dryness affording a pale yellow oil which was dried under high vacuum to remove volatiles. ^1H n.m.r. and t.l.c. (D) showed that the crude product (820 mg, 90 %) was still contaminated with benzyl alcohol as well as traces of 131. This material was chromatographed on silica with CH_2Cl_2 -acetone (20:1) as eluant to give compound 134 (725 g, 80 %) as a chromatographically pure oil (ether or E); δ_{H} 7.34 (s, 5H, arom. H), 5.12 (s, 2H, CH_2Ph), 3.08-3.30 (m, 8H, CH_2NHBoc and CH_2NZCH_2), and 1.47-1.74 (m) and 1.43 (s) [together 24H, $\text{C}\underline{\text{C}}\text{H}_2\text{C}$ and $\text{C}(\underline{\text{C}}\text{H}_3)_3$]; δ_{C} 155.9 (CO), 136.6, 128.5, 128.2, 128.0, and 127.8 (arom. H),

79.2 and 79.1 [$\text{OC}(\underline{\text{C}}\text{H}_3)_3$], 67.1 ($\underline{\text{C}}\text{H}_2\text{Ph}$), 46.4 and 44.3 [$\underline{\text{C}}\text{H}_2\text{N}(\text{Z})\underline{\text{C}}\text{H}_2$], 40.2 and 37.5 ($\underline{\text{C}}\text{H}_2\text{NHBoc}$), 28.4 [$\text{C}(\underline{\text{C}}\text{H}_3)_3$], and 28.4, 27.3, and 25.6 ($\text{C}\underline{\text{C}}\text{H}_2\text{C}$).


Method B - Benzyloxycarbonylation of compound 133



To an ice-cold solution of compound **133** (173 mg, 0.50 mmol) in dry CH_2Cl_2 (2 ml) was added dropwise with vigorous stirring a solution of Z_2O (157 mg, 0.55 mmol) in dry CH_2Cl_2 (1 ml). The clear, colourless solution was stirred for 1 h in ice-bath and overnight at r.t. The solvent was evaporated under reduced pressure and the crude yellowish residue was directly chromatographed on silica with ether as eluant to afford the desired product **134** (186 mg, 77 %) as a pale yellow oil. This sample was identical with the product obtained by procedure A (t.l.c. and ^1H and ^{13}C n.m.r. spectra).


5.2.3 - Synthesis of N¹-benzyloxycarbonyl-N⁸-tert-butoxy-carbonylspermidine

N¹,N⁴-Methylenespermidine (114b)



This compound was prepared from spermidine according to a previously described procedure⁸⁰, using a fresh formaldehyde solution. The yield of essentially pure **114b**, obtained as a colourless oil which soon solidified, was 95 %; δ_{H} 3.38 (s, 2H, NCH₂N), 2.51-2.88 (m, 6H, CH₂N), 2.17-2.33 (m, 2H, CH₂NH₂), and 1.40-1.73 (m, together 8H, CCH₂C and NH₂); δ_{C} 69.9 (NCH₂N), 55.4 and 53.1 (NCH₂CH₂CH₂N), 45.2 and 42.1 (CH₂N), and 31.8, 27.2, and 24.3 (CCH₂C).


N¹,N⁸-Dibenzyloxycarbonyl-N¹,N⁴-methylenespermidine (135)



A solution of Z₂O (12 g, 42 mmol) in dry CH₂Cl₂ (15 ml) was added dropwise to a cooled solution of compound **114b** (3.0 g, 19 mmol) in dry CH₂Cl₂ (15 ml). The resulting clear mixture was stirred for 1 h in ice-bath and 2 h at r.t. The solvent was removed under reduced pressure and the colourless residue was chromatographed on silica using CH₂Cl₂-acetone (4:1) as eluant to afford the chromatographically essentially pure (G)

product **135** as a pale yellow oil (6.1 g, 76 %); δ_{H} 7.34 (s, 10H, arom. H), 5.12 and 5.08 (2s, 4H, CH_2Ph), 4.14 (s, 2H, NCH_2N), 3.53 (t, 2H) and 3.13-3.23 (m, 2H, CH_2NZ), 2.71 (t, 2H), and 2.29-2.53 (m, 2H) (CH_2N), and 1.33-1.74 (m, 6H, CCH_2C); δ_{C} 156.4 and 155.0 (CO), 136.6, 128.4, and 128.0 (arom. C), 67.0 and 66.4 (CH_2Ph), 65.0 (NCH_2N), 52.4 and 52.2 (2 x CH_2N), 43.8 and 40.8 (2 x ZNCH_2C), and 27.6, 24.4, and 22.9 (CCH_2C).

N^1, N^8 -Dibenzoyloxycarbonyl- N^8 -tert-butoxycarbonyl- N^1, N^4 -methylenespermidine (**136**)




To a stirred solution of compound **135** (4.40 g, 10.3 mmol) in dry CH_3CN (20 ml) was added DMAP (128 mg, 1.03 mmol), followed by Boc_2O (2.47 g, 11.3 mmol). After 7 h stirring, t.l.c. (G) showed that starting material still remained in the mixture and more Boc_2O (1.24 g, 5.65 mmol) was introduced and left overnight. When the reaction was complete the solvent was evaporated under reduced pressure and the brownish residue was chromatographed (silica; ether) to give pure (A and G) compound **136** as a yellow oil (4.9 g, 90 %); δ_{H} 7.35 and 7.33 (two s, 10H, arom. H), 5.20 (s, 2H, $\text{BocNCO}_2\text{CH}_2\text{Ph}$), 5.12 (s, 2H, CH_2Ph), 4.12 (s, 2H, NCH_2N), 3.35-3.62 (m, together 4H, CH_2NZ and $\text{CH}_2\text{N}(\text{Z})\text{Boc}$), 2.67 (t, 2H) and 2.38 (t, 2H) (CH_2N), and 1.46-1.72 (m) and 1.46 (s) [together 15H, CCH_2C]

and $C(CH_3)_3$]; δ_C 154.8, 153.6, and 151.8 (CO), 136.5, 135.3, 128.3, 128.0, and 127.7 (arom. C), 82.5 [$OC(CH_3)_3$], 68.0 ($PhCH_2O_2CNBoc$), 66.9 (CH_2Ph), 65.2 (NCH_2N), 52.2 and 51.6 (CH_2N), 46.2 and 43.6 (CH_2NZ), 27.8 [$C(CH_3)_3$], and 26.6, 24.2, 22.9, and 22.5 (CCH_2C).

N^1 -Benzyloxycarbonyl- N^8 -tert-butoxycarbonylspermidine (138)

Method A: via the intermediate 137


- a) N^1 -Benzyloxycarbonyl- N^8 -tert-butoxycarbonyl- N^1, N^4 -methyl-
enespermidine (137)



A stirred solution of compound **136** (3.67 g, 6.98 mmol) in dry methanol (40 ml) was treated with TMG (402 mg, 3.50 mmol) at r.t. for 2 days. The coloured solution was evaporated under reduced pressure and the yellowish residue was chromatographed (silica; CH_2Cl_2 -acetone, 3:1) to afford the chromatographically pure (G) product **137** as a yellow oil (2.42 g, 88 %); δ_H 7.34 (s, 5H, arom. H), 5.13 (s, 2H, CH_2Ph), \approx 4.9 (broad signal, \approx 1H, NH), 4.13 (s, 2H, NCH_2N), 3.53 (t, 2H, CH_2NZ), 2.96-3.13 (m, 2H, CH_2NHBoc), 2.70 (t, 2H) and 2.17-2.40 (m, 2H) (CH_2N), and 1.34-1.75 (m) and 1.44 (s) [together 15H, CCH_2C and $C(CH_3)_3$]; δ_C 155.9 and 155.0 (CO), 136.6, 128.4, 128.0, and 127.9 (arom. C), 78.9 [$OC(CH_3)_3$], 67.1 (CH_2Ph), 65.1 (NCH_2N), 52.6 and 52.2 (CH_2N), 43.8

($\underline{\text{C}}\text{H}_2\text{NZ}$), 40.4 ($\underline{\text{C}}\text{H}_2\text{NHBoc}$), 28.4 [$\text{C}(\underline{\text{C}}\text{H}_3)_3$], and 27.7, 24.5, and 22.8 ($\text{C}\underline{\text{C}}\text{H}_2\text{C}$).

b) ring cleavage of compound 137




A solution of compound **137** (2.40 g, 6.10 mmol) in dry methanol (50 ml) was refluxed with pyridine (1.50 g, 19.0 mmol) and malonic acid (2.30 g, 22.4 mmol) with stirring for 2 h. The solvent was evaporated under reduced pressure and after the addition of aqueous 30 % K_2CO_3 (30 ml) the product was extracted with CHCl_3 (3 x 60 ml). The combined yellowish organic layers were washed with saturated aqueous NaCl (2 x 30 ml), dried (Na_2SO_4), and evaporated to afford a crude oil which was chromatographed on silica (CH_2Cl_2 - MeOH - HOAc , 18:2:1). The appropriate fractions were collected and again neutralized as for the crude product to give a yellow oil which was triturated with light petroleum to afford **138** as a white solid (1.81 g, 78 %); homogeneous by t.l.c. (M or P). An analytical specimen was obtained by recrystallization from heptane-ether (2:1; 100 ml/g); m.p. 63-64 °C; δ_{H} 7.34 (s, 5H, arom. H), \approx 5.65 (broad, \approx 1H, amide NH), 5.09 (s, 2H, $\underline{\text{C}}\text{H}_2\text{Ph}$), \approx 4.80 (broad, \approx 1H, amide NH), 3.06-3.30 (m, together 4H, $\underline{\text{C}}\text{H}_2\text{NHZ}$ and $\underline{\text{C}}\text{H}_2\text{NHBoc}$), 2.51-2.74 (m, 4H, $\underline{\text{C}}\text{H}_2\text{N}$), and 1.49-1.79 (m) and 1.43 (s) (together \approx 16H, $\text{C}\underline{\text{C}}\text{H}_2\text{C}$, $\text{C}(\underline{\text{C}}\text{H}_3)_3$, and

amine NH); δ_C 156.5 and 156.0 (CO), 136.7, 128.4, and 128.0 (arom. C), 79.0 [$\text{OC}(\text{CH}_3)_3$], 66.4 (CH_2Ph), 49.3 and 47.7 (CH_2N), 40.9 and 39.9 (CH_2NHBoc and CH_2NHZ), 29.5 (CCH_2C), 28.4 [$\text{C}(\text{CH}_3)_3$], and 27.8 and 27.2 (CCH_2C). (Found: C, 63.3; H, 8.6; N, 11.3. $\text{C}_{20}\text{H}_{33}\text{N}_3\text{O}_4$ requires C, 63.3; H, 8.8; N, 11.1%).

Method B: via compound 139

a) N^1, N^8 -Dibenzoyloxycarbonyl- N^8 -tert-butoxycarbonyl-spermidine (139)



To a solution of the N^1, N^4 -methylenespermidine derivative **136** (2.00 g, 3.80 mmol) in dry methanol (35 ml) were added pyridine (94.1 mg, 11.9 mmol) and malonic acid (1.45 g, 13.9 mmol). The mixture was refluxed with stirring for 2 h, then worked up as described in method A, b) (p. 122) except that the crude product was chromatographed with CH_2Cl_2 -acetone-HOAc (5:5:1) to furnish the essentially pure (L or M) compound **139** as a yellow oil (900 mg, 48 %); δ_H 7.35 and 7.33 (two s, 10H, arom. H), 5.65 (broad signal, \approx 1H, amide NH), 5.20 (s, 2H, $\text{PhCH}_2\text{O}_2\text{CNBoc}$), 5.08 (s, 2H, CH_2Ph), 3.65 (t, 2H, $\text{CH}_2\text{N}(\text{Z})\text{Boc}$), 3.24 (m, 2H, CH_2NHZ), 2.60 (q, 4H, CH_2N), and 1.46-1.77 (m) and 1.46 (s) (together \approx 16H, CCH_2C , $\text{C}(\text{CH}_3)_3$, and amine NH); δ_C 156.5, 153.9, and 152.1 (CO), 136.7, 135.5, 128.5, 128.2, and 128.0 (arom. C), 82.7 [$\text{OC}(\text{CH}_3)_3$], 68.2


(PhCH₂O₂CNBoc), 66.4 (CH₂Ph), 49.4 and 47.6 (CH₂NCH₂), 46.3 [CH₂N(Z)Boc], 39.9 (CH₂NHZ), 29.6 (CCH₂C), 28.0 [C(CH₃)₃], and 27.0 and 26.6 (CCH₂C).

b) methanolysis of compound 139 catalysed by TMG

A stirred solution of compound **139** (281 mg, 0.550 mmol) in dry methanol (2.50 ml) was treated with TMG (32.0 mg, 0.280 mmol) at r.t. for about 2 days. The solvent was removed under reduced pressure and the residue chromatographed and worked up as described in method A, b) (p. 122) to afford compound **138** (135 mg, 65 %); m.p. and ¹H and ¹³C n.m.r. spectra were in agreement with the foregoing data.

Method C: independent synthesis of compound 138


a) N¹-(Benzyloxycarbonyl-β-alanyl)-N⁴-tert-butoxycarbonyl-tetramethylenediamine (140)



To a solution of N¹-Boc-tetramethylenediamine^{17a,121} (1.13 g, 6.00 mmol) in dry acetonitrile (20 ml) Z-βAla-ONp (1.72 g, 5.00 mmol) dissolved in dry acetonitrile (20 ml) was added dropwise with vigorous stirring for 15 min. The resulting mixture immediately became bright yellow, and after 1 h agitation at r.t. a precipitate appeared. More

acetonitrile (10 ml) was added to facilitate stirring, which was continued overnight (20 h). The thick sludge was filtered by suction and the collected white solid was thoroughly triturated repeatedly with cold acetonitrile (3 x 5 ml) and sucked dry. The crude yield of the chromatographically pure product **140** was 1.82 g (92 %). Recrystallization from acetonitrile (30 ml/g) gave, after cooling for a few days, a white fluffy crystalline solid (90 % crystallization yield); t.l.c. (M or Q) gave one spot; m.p. 133-134 °C; δ_H 7.33 (s, 5H, arom. H), 6.10 (broad, \approx 1H, CONH), 5.62 (broad, \approx 1H, ZNH), 5.09 (s, 2H, \underline{CH}_2 Ph), 4.68 (broad, \approx 1H, BocNH), 3.46 (perturbed t, 2H), 3.18 (perturbed m, 2H), and 3.13 (perturbed s, 2H) (3 x \underline{NCH}_2), 2.39 (t, 2H, \underline{COCH}_2), \approx 1.49 (m, 4H, $\underline{CCH}_2\underline{CH}_2\underline{C}$), and 1.43 [s, 9H, $\underline{C(CH}_3)_3$]; δ_C 171.3 ($\underline{CH}_2\underline{CONH}$), 156.5 and 156.1 (2 x \underline{OCONH}), 136.5, 128.4, 128.0, and 127.9 (arom. C), 79.2 [$\underline{C(CH}_3)_3$], 66.6 (\underline{CH}_2 Ph), 40.1 and 39.1 ($\underline{CH}_2\underline{NHCO}_2$), 37.2 and 35.9 ($\underline{CH}_2\underline{CONHCH}_2$), 28.4 [$\underline{C(CH}_3)_3$], 27.5 and 26.5 ($\underline{CCH}_2\underline{CH}_2\underline{C}$).

b) reduction of the amide **140**




To a suspension of finely ground NaBH_4 (190 mg, 5.0 mmol) in dry THF (8 ml), TFA (385 μ l, 5.0 mmol) was added dropwise under vigorous stirring at r.t. for \approx 10 min (evolution of

gas). The resulting mixture was allowed to stay for 30 min. To this was added the amide **140** (393 mg, 1.00 mmol) in dry THF (10 ml). The suspension was stirred 4 h at r.t. and overnight at 40 °C. Then, 20 % aq. HOAc (3 ml) was carefully added to the reaction mixture under stirring at r.t. After 2 h, the solvent was evaporated and the residue partitioned between CH₂Cl₂ (60 ml) 30 % K₂CO₃-saturated NaCl (1:1) (30 ml). The aqueous phase was again extracted with CH₂Cl₂ (2 x 20 ml). The combined organic layers were washed with saturated NaCl (2 x 20 ml) and dried (Na₂SO₄). Evaporation to dryness afforded 380 mg of a crude mixture which was chromatographed on silica with CH₂Cl₂-MeOH-HOAc (18:2:1). First eluted the amide and some other impurities. The product was then eluted with CH₂Cl₂-MeOH-HOAc (16:4:1). The fractions contained pure compound were pooled and evaporated to dryness. The semisolid residue was worked up as for the crude mixture to give 88 mg (23 %) of a waxy solid. A recrystallized sample was identical with compound **138** as obtained earlier.

5.2.4 - Attempted synthesis of N⁸-benzyloxycarbonyl-N¹-tert-butoxycarbonylspermidine

N⁸-Benzyloxycarbonyl-N¹-tert-butoxycarbonyl-N¹,N⁴-methylenespermidine (160)


A - One pot procedure



A solution of dried cyclic spermidine **114b** (1.19 g, 7.56 mmol) in dry CH₂Cl₂ (30 ml) was treated dropwise over a period of 1 h with ZCN (1.34 g, 8.31 mmol) dissolved in dry CH₂Cl₂ with rapid stirring at r.t. The resulting, slightly turbid reaction mixture was stirred for a further 3 h with exclusion of moisture. A solution of Boc₂O (1.81 g, 8.32 mmol) in dry CH₂Cl₂ (20 ml) was slowly introduced during 20 min at r.t. followed by the dropwise addition of NMM (840 mg, 8.31 mmol) in dry CH₂Cl₂ (25 ml). After stirring overnight, most of the solvent was stripped off and the residue dissolved in ethyl acetate (300 ml). The extract was washed with 1M NaHCO₃ and saturated NaCl (3 x 100 ml each) and dried (Na₂SO₄). Evaporation to dryness afforded 2.95 g of a mixture containing ≈ 60 % of the desired product **160** as well as the N¹,N⁸-Z₂-derivative **135** (as judged from t.l.c. (P) and ¹H n.m.r.). Column chromatography on silica (CH₂Cl₂-MeOH, 30:1) gave small amounts of essentially pure product as a pale yellow oil; δ_H

7.33 (s, 5H, arom. H), \approx 5.40 (broad, \approx 1H, NH), 5.09 (s, 2H, CH_2Ph), 4.06 (s, 2H, NCH_2N), 3.44 (t, 2H, CH_2NBoc), 3.10-3.30 (m, 2H, CH_2NHZ), 2.69 (t, 2H), and 2.33-2.49 (m, 2H) (CH_2N), 1.49-1.62 (m), and 1.45 [s, together 15H, CCH_2C and $\text{C}(\text{CH}_3)_3$].


B - With isolation of $\text{N}^8\text{-Z-N}^1, \text{N}^4$ -methylenespermidine 159



To a solution of hexahydropyrimidine **114b** (4.30 g, 27.0 mmol) in dry CH_2Cl_2 (30 ml) ZCN (4.41 g, 27.0 mmol) in dry CH_2Cl_2 (30 ml) was added dropwise over a period of 3 h with stirring at r.t. After a further 2 h stirring, t.l.c. (CH_2Cl_2 -MeOH, 1:1) showed a rather complex mixture. The solvent was evaporated under reduced pressure and the oily residue chromatographed on silica with CH_2Cl_2 -MeOH (1:1) as eluant. The fractions containing the almost pure compound were pooled and evaporated to afford 448 mg (5 %) of $\text{N}^8\text{-Z-N}^1, \text{N}^4$ -methylenespermidine **159** as a yellow oil. This product (448 mg, 1.54 mmol) in dry CH_2Cl_2 (5 ml) was treated dropwise with Boc_2O in dry CH_2Cl_2 (5 ml) at room temperature. After 2 h stirring, the solvent was evaporated under reduced pressure and the yellowish residue worked up as under procedure A to afford crude **160**. This material was chromatographed on silica (CH_2Cl_2 -acetone, 3:1) to afford 283 mg (47 %) of a homogeneous yellow oil, identical with

compound **160** as obtained above.

N⁸-Benzyloxycarbonyl-N¹-tert-butoxycarbonylspermidine (161)




Compound **160** was treated and the product worked up as described for compound **138** under method A, b) (p. 122) to afford 167 mg (74 %) of **161** as a waxy solid, essentially pure by t.l.c. (M or P); δ_{H} 7.33 (s, 5H, arom. H), \approx 5.61 and \approx 5.31 (broad, \approx 2H, amide NH), 5.08 (s, 2H, CH_2Ph), 3.05-3.27 (m, 4H, CH_2NHBoc , CH_2NHZ), 2.55-2.75 (m, 4H, CH_2NHCH_2), 1.47-1.74 (m) and 1.43 (s, together 16H, CCH_2C , $\text{C}(\text{CH}_3)_3$ and amine NH).

5.2.5 - Synthesis of N⁴-acetylspermidine dioxalate


N¹,N⁴,N⁸-Triacetylspermidine (141)

A - Acetylation in aqueous conditions




Spermidine (1.12 g, 7.7 mmol) was dissolved in 1 M NaOH (20 ml) and, after cooling in ice water, simultaneously treated under stirring dropwise with Ac₂O (3.17 g, 31.0 mmol) and 1 M NaOH (80 ml) and then left for several hours. The solution was saturated with NaCl and extracted with CHCl₃ (4 x 50 ml). The extract was dried (MgSO₄) and evaporated to afford 1.63 g (78 %) of a colourless oil. This crude material was chromatographed on silica with CH₂Cl₂-MeOH (4:1) to yield 1.36 g (65 %) of compound **141** as a pale yellow oil, homogeneous by t.l.c. (N, L); δ_H ≈ 6.98 and 6.40 (broad, ≈ 2H, amide NH), 3.07-3.46 (m, 8H, CH₂N), 2.10 and 2.07 (two s, 3H, -N(CH₃CO)-), 1.98 (s, 6H, CH₃CONH), 1.49-1.87 (m, 6H, CCH₂C); δ_C 171.0 and 170.5 (CO), 48.4, 46.7, 45.2, 42.5, 38.7, 36.9 and 36.1 (CH₂N), 29.0, 27.7, 27.4, 27.0, 26.5, 25.9, 24.8, 23.3, 23.1, 23.0, 21.4 (other C).

B - Acetylation in anhydrous conditions



An ice-cooled solution of spermidine (1.00 g, 6.88 mmol) and TEA (2.16 g, 21.3 mmol) in dry CH_2Cl_2 (10 ml) was treated dropwise with Ac_2O (2.18 g, 21.3 mmol) and then stirred overnight at r.t. The solvent was evaporated and the colourless residue was chromatographed as described above to afford 1.37 g (73 %) of compound **141** as a pale yellow oil, essentially pure by t.l.c. (L, N) which was identical with the product obtained by procedure A.

$\text{N}^1, \text{N}^4, \text{N}^8$ -Triacetyl- N^1, N^8 -bis(tert-butoxycarbonyl)-spermidine (142)




A solution of compound **141** (0.787 g, 2.90 mmol) and DMAP (71 mg, 0.58 mmol) in CH_3CN (10 ml) was treated with Boc_2O (1.40 g, 6.40 mmol) in one portion and left with stirring at r.t. After 4 h t.l.c. (N) showed that more than 50 % of the starting material remained. Additional Boc_2O was added in six

portions (1 eq. each) at intervals during 5 days. T.l.c. (N) still showed some starting material left and two other major spots. The reaction mixture was therefore evaporated to dryness and the residue was again dissolved in CH_3CN (10 ml) and a new batch of Boc_2O (1 eq.) and DMAP (0.1 eq.) was added. The reaction was left overnight. This procedure was repeated once more until t.l.c. (N) of the reaction mixture showed one major spot. The solvent was evaporated in vacuo and the dark, brown residue partitioned between 1 M KHSO_4 (50 ml) and ether (100 ml). The solution was again extracted with ether (2 x 25 ml) and the combined organic layers were washed in turn with 1 M KHSO_4 , 1 M NaHCO_3 and saturated NaCl (2 x 50 ml each). The yellowish extract was dried (MgSO_4) and evaporated. The brown residue was chromatographed on silica using CH_2Cl_2 -acetone (9:1) to afford 932 mg (68 %) of **142**, pure by t.l.c. (N, ether); δ_{H} 3.68 [t, 4H, $\text{CH}_2\text{N}(\text{Ac})\text{Boc}$], 3.14-3.40 (m, 4H, $\text{CH}_2\text{NacCH}_2$), 2.46 [s, 6H, $\text{CH}_3\text{CO}(\text{Boc})\text{N}$], 2.07 (s, 3H, $-\text{N}(\text{CH}_3\text{CO})-$), 1.25-1.73 (m) and 1.54 [s, together \approx 24H, $\text{CCH}_2\text{C} + \text{C}(\text{CH}_3)_3$]; δ_{C} 172.9 (BocNCOCH_3), 170.1 and 170.0 (CH_3CON), 153.0 and 152.8 ($\text{Bu}^{\text{t}}\text{O-CO}$), 83.5, 83.3, 83.1 and 83.0 [$\text{OC}(\text{CH}_3)_3$], 48.2, 46.3, 45.2, 43.8, 43.6, 43.0, 42.1 and 41.7 (CH_2N), 28.1 [$\text{C}(\text{CH}_3)_3$], 26.9, 26.1, 25.9, 25.0 and 21.5 (other C).


N⁴-Acetyl-N¹,N⁸-bis(tert-butoxycarbonyl)spermidine (143)

Method A: Methanolysis of compound 142 in the presence of catalytic amounts of TMG




Compound **142** (547 mg, 1.2 mmol) was dissolved in dry methanol (10 ml) and treated with TMG (30 mg, 0.26 mmol) with stirring at r.t. for 4 h. The reaction mixture was evaporated in vacuo and the yellow residue partitioned between ether (60 ml) and 1M KHSO₄ (30 ml). After further extraction with ether (60 ml), the combined organic layers were washed in turn with 1M KHSO₄, 1M NaHCO₃ and saturated NaCl (2 x 30 ml) and dried (MgSO₄). The extract was evaporated to dryness to afford 418 mg (90 %) of a yellow oil. The crude product was chromatographed on silica (CH₂Cl₂-acetone, 2:1) to furnish 395 mg (88 %) of compound **143** as a pale yellow oil, homogeneous by t.l.c. (F, P); $\delta_{\text{H}} \approx 5.4$ and 4.7 (broad, $\approx 2\text{H}$, amide NH), 3.04-3.52 (m, 8H, CH₂N), 2.09 and 2.08 (two s, 3H, CH₃CON), 1.50-1.73 (m) and 1.44 [s, together 24H, CCH₂C + C(CH₃)₃]; δ_{C} 170.8 and 170.2 (CH₃CO), 156.1 and 156.0 (Bu^tO-CO), 79.4 and 78.9 [C(CH₃)₃], 48.3, 46.4, 45.3, 42.4, 40.0 and 37.4 (CH₂N), 28.4 [C(CH₃)₃], 29.7, 28.0, 27.6, 26.0, 25.0 and 21.4 (other C).

Method B: acetylation of compound 133



A solution of Ac₂O (123 mg, 1.20 mmol) in CH₂Cl₂ (5 ml) was added to a cooled solution of **133** (345 mg, 1.00 mmol) and TEA (152 mg, 1.50 mmol) in CH₂Cl₂ (10 ml) and the reaction was stirred for 4 h. The solvent was evaporated in vacuo and the remaining colourless residue worked up and chromatographed as described in Method A. It was afforded 288 mg (74 %) of compound **143** as a pale yellow oil. ¹H and ¹³C n.m.r. spectra were identical with those given under Method A.

N⁴-Acetylspermidine dioxalate (144a)




Compound **143** (324 mg, 0.84 mmol) was treated with 2.29 M HCl in dioxan (2 ml) with stirring at r.t. for 3 h. Most of the solvent was evaporated in vacuo and the sticky residue taken up in ether (20 ml) and evaporated twice. The product was then dissolved in distilled water (40 ml) and the resulting solution extracted with ether (3 x 20 ml). The aqueous layer was flushed with N₂ to remove ether and

lyophilized to afford 202 mg (92 %) of a sticky white residue, essentially pure by t.l.c. (S). This material was converted to its oxalate salt by loading a portion (100 mg), dissolved in water, onto a QAE-Sephadex A-25 column (oxalate form) and eluting with distilled water to afford 111 mg of a white residue after lyophilization. Recrystallization from water-ethanol (1:20, 25 ml) gave the oxalate salt **144a** as a white solid; pure by h.p.l.c. (see Figs. 7 and 8, p. 142 and 143); m.p. 187.5-188.5 °C; δ_{H} (D_2O) 3.27-3.55 (m, 4H, $\text{CH}_2\text{N}(\text{Ac})\text{CH}_2$), 2.82-3.13 (m, 4H, CH_2NH_2), 2.14 and 2.13 (two s, 3H, CH_3CON), 1.54-2.08 (m, 6H, CCH_2C); δ_{C} 177.2, 176.7 and 176.0 (CO), 51.0, 48.7, 47.9, 45.0, 41.8 and 39.5 (CH_2N), 28.5, 27.6, 27.5, 26.8, 26.4, 23.3 and 23.1 (other C). (Found: C, 46.1; H, 8.0; N, 14.7. $\text{C}_9\text{H}_{21}\text{N}_3\text{O}\cdot\text{C}_2\text{H}_2\text{O}_4\cdot 1/2\text{H}_2\text{O}$ requires C, 46.1; H, 8.45; N, 14.7 %).

5.2.6 - Synthesis of N^1 -acetylspermidine dihydrochloride


N^1 -Benzyloxycarbonyl- N^4, N^8 -bis(tert-butoxycarbonyl)-spermidine (145)



To an ice-cold solution of compound **138** (1.90 g, 5.01 mmol) in CH_2Cl_2 (10 ml) was added dropwise a solution of Boc_2O

(1.15 g, 5.26 mmol) in dry CH_2Cl_2 (10 ml). The colourless reaction mixture was stirred for 30 min. in ice and overnight at r.t. The solvent was evaporated and the residue partitioned between 1 M KHSO_4 (100 ml) and ether (500 ml). The extract was washed and dried as described before and evaporated to afford 3.0 g of a pale yellow oil. Column chromatography (silica, ether-light petroleum, 3:1) furnished 2.10 g (87 %) of the product **145**, homogeneous by t.l.c. (C,F); δ_{H} 7.34 (s, 5H, arom. H), 5.10 (s, 2H, CH_2Ph), \approx 5.70 and 4.60 (broad, \approx 2H, amide NH), 3.08-3.32 (m, 8H, CH_2NBoc , CH_2NHZ), 1.51-1.78 (m) and 1.44 [s, together 24H, $\text{CCH}_2\text{C} + \text{C}(\text{CH}_3)_3$]; δ_{C} 156.4 and 155.9 (CO), 136.6, 128.4 and 128.0 (arom. C), 79.7 and 79.2 [$\text{OC}(\text{CH}_3)_3$], 66.4 (OCH_2Ph), 46.6 and 43.7 [$\text{CH}_2\text{N}(\text{Boc})\text{CH}_2$], 40.2 and 37.8 (CH_2NHBoc , CH_2NHZ), 28.4 [$\text{C}(\text{CH}_3)_3$], 27.4 and 25.6 (CCH_2C).


N^4, N^8 -Bis(tert-butoxycarbonyl)spermidine (146)



Compound **145** (1.90 g, 3.96 mmol) was hydrogenolyzed as described for model compound **170** (p. 106). The catalyst was filtered off and rinsed with methanol. The colourless filtrate was evaporated under reduced pressure to give 1.35 g (98 %) of the product **146** as a colourless oil, essentially pure by


t.l.c. (M, Q); $\delta_{\text{H}} \approx 4.60$ (broad, $\approx 1\text{H}$, amide NH), 3.09–3.46 [m, 6H, CH_2NHBoc , $\text{CH}_2\text{N}(\text{Boc})\text{CH}_2$], 2.69 (t, 2H, CH_2NH_2), 1.53–1.71 (m), 1.45 and 1.44 [two s, together 26H, CCH_2C , $\text{C}(\text{CH}_3)_3 + \text{NH}_2$]; δ_{C} 156.0 and 155.7 (CO), 79.4 and 79.1 [$\text{OC}(\text{CH}_3)_3$], 46.5, 44.2, 40.3 and 39.3 (CH_2N), 32.3, 27.5 and 25.7 (CCH_2C), 28.5 [$\text{C}(\text{CH}_3)_3$].

N^1 -Acetyl- N^4, N^8 -bis(tert-butoxycarbonyl)spermidine (147)



A solution of compound **146** (1.22 g, 3.53 mmol) was treated and the product purified in a similar manner as described for the derivative **143** (Method B, p. 134): yield 1.30 g (95 %) of the product **147** obtained as an oil; $\delta_{\text{H}} \approx 6.75$ and 4.60 (broad, amide NH), 3.02–3.33 (m, 8H, CH_2N), 1.98 (s, 3H, CH_3CON), 1.53–1.72 (m) and 1.46 and 1.44 [two s, together $\approx 24\text{H}$, $\text{CCH}_2\text{C} + \text{C}(\text{CH}_3)_3$]; δ_{C} 170.2 (CH_3CO), 156.0 ($\text{Bu}^t\text{O-CO}$), 79.8 and 79.2 [$\text{OC}(\text{CH}_3)_3$], 46.6, 44.1, 40.1 and 35.9 (CH_2N), 28.4 ($\text{C}(\text{CH}_3)_3$), 27.7, 27.5 and 25.6 (CCH_2C), 23.4 (CH_3CON).

N^1 -Acetylspermidine dihydrochloride (148)




Compound **147** (539 mg, 1.39 mmol) was treated with 2.29 M HCl in dioxan (5 ml) at r.t. for 3 h. The solvent was evaporated in vacuo and the white residue suspended in ether (20 ml) and evaporated twice. It was afforded 350 mg (97 %) of the dihydrochloride salt **148**. It was recrystallized from ethanol and was pure by h.p.l.c. (see Figs. 7 and 8, p. 142 and 143); m.p. 191-193 °C (lit.^{35a,110} 173-178 °C or 189-191 °C; δ_{H} (D₂O) 3.28 (t, 2H, J=6.7 Hz, CH_2NHAc), 2.98-3.15 (m, 6H, CH_2N), 2.00 (s, 3H, CH_3CO), 1.74-1.88 (m, 6H, CCH_2C); δ_{C} 177.2 (CO), 49.6, 47.7, 41.4 and 38.7 (CH_2N), 28.2, 26.6 and 25.4 (CCH_2C), 24.5 (CH_3CON). (Found: C, 41.3; H, 8.8; N, 15.9. C₉H₂₁N₃O.2HCl requires C, 41.54; H, 8.91; N, 16.15%).

5.2.7 - Synthesis of N⁸-acetylspermidine dihydrochloride

N¹,N⁴-Dibenzoyloxycarbonyl-N⁸-tert-butoxycarbonyl-spermidine (149)


A - Benzyloxycarbonylation with Z₂O



A solution of compound **138** (380 mg, 1.00 mmol) in dry CH₂Cl₂ (2 ml) was treated with Z₂O¹⁰⁸ as described for compound **135** (p. 119). After completion of the reaction

[t.l.c. (L)] the solvent was evaporated under reduced pressure. The oily residue was partitioned between ether (50 ml) and aqueous 1 M KHSO₄ (25 ml). The organic layer was washed and dried as usual. Evaporation to dryness afforded the crude product (638 mg) as a pale yellow oil contaminated with benzyl alcohol. The crude material was chromatographed on silica (ether-light petroleum, 3:1) to give 452 mg (88 %) of product **149** as a light yellow oil, homogeneous by t.l.c. (ether, C); δ_{H} 7.33 and 7.32 (two s, 10H, arom. H), 5.11 and 5.08 (2s, 4H, CH₂Ph), 3.04-3.38 (m, 8H, CH₂N), 1.51-1.76 (m) and 1.43 [s, together 15H, CCH₂C + C(CH₃)₃]; δ_{C} 156.4 and 155.9 (CO), 136.6, 128.5, 128.4 and 128.0 (arom. C), 79.2 [OC(CH₃)₃], 67.1 and 66.5 (CH₂Ph), 46.5, 44.1, 40.1 and 37.6 (CH₂N), 28.4 [C(CH₃)₃], 28.2, 27.4 and 25.6 (CCH₂C).


B - Benzyloxycarbonylation with ZOBT



To a stirred suspension of ZOBT¹¹¹ (1.30 g, 4.82 mmol) in dry CH₃CN (40 ml) was added a solution of compound **138** (1.18 g, 4.77 mmol) in the same solvent (30 ml). The clear solution obtained was left overnight at r.t. The solvent was evaporated and the product worked up and chromatographed as described in procedure A to furnish 2.04 g (83 %) of compound **149** as a pale yellow oil. T.l.c. and ¹H and ¹³C spectra were in agreement with those reported for the product obtained


under procedure A.

N¹,N⁴-Dibenzyloxycarbonylspermidine (150)



Compound **149** (1.84 g, 3.58 mmol) was treated with 2.29 M HCl in dioxan (15 ml) and stirred at r.t. for 3 h. The solvent was evaporated and the white residue treated with 30 % K₂CO₃ (40 ml) and extracted with CHCl₃ (5 x 100 ml). The combined organic layers were dried (Na₂SO₄) and evaporated to afford 1.42 g (96 %) of product **150** as a pale yellow oil, nearly pure by t.l.c. (M, Q); δ_H 7.34 and 7.33 (two s, 10H, arom. H), ≈ 5.60 (broad, ≈1H, amide NH), 5.11 and 5.08 (2s, 4H, CH₂Ph), 3.05-3.39 (m, 6H, CH₂NHZ, CH₂N(Z)CH₂), 2.66 (t, 2H, CH₂NH₂), 1.25-1.84 (m, 8H, CCH₂C + NH₂). δ_C 156.4 (CO), 136.6, 128.5, 128.4, 128.0 and 127.9 (arom. C), 67.1 and 66.5 (CH₂Ph), 46.8, 44.1, 41.7 and 37.7 (CH₂N), 30.7, 28.1 and 25.8 (CCH₂C).


N⁸-Acetyl-N¹,N⁴-dibenzyloxycarbonylspermidine (151)



A solution of compound **150** (1.18 g, 2.85 mmol) was treated

and the product purified in a similar manner as described for derivative **143** (Method B, p. 134) to give 1.10 g (85 %) of product **151** as an oil; δ_{H} 7.33 (s, 10H, arom. H), \approx 6.00 and 5.60 (broad, \approx 2H, amide NH), 5.11 and 5.08 (two s, 4H, CH_2Ph), 3.05-3.37 (m, 8H, CH_2NZ , CH_2NAc), 1.92 (s, 3H, CH_3CON), 1.33-1.77 (m, 6H, CCH_2C); δ_{C} 170.2 (CH_3CO), 156.5 (Bz10- CO), 136.5, 128.6, 128.4, 128.0 and 127.9 (arom. C), 67.2 and 66.5 (CH_2Ph), 46.5, 44.3, 39.0 and 37.9 (CH_2N), 28.4, 26.7 and 25.7 (CCH_2C), 23.2 (CH_3CON).

N⁸-Acetylspermidine dihydrochloride (152)



A solution of compound **151** (497 mg, 1.09 mmol) was hydrogenolyzed in a similar manner as described for model compound **170** (p. 106) to give the free amine (200 mg, 98 %) as a colourless oil. This derivative was converted to its dihydrochloride salt with excess of 2.29 M HCl in dioxan to afford 250 mg (90 %) of the salt **152**. It was recrystallized from ethanol and was pure by h.p.l.c. (see Figs. 7 and 8, p. 142 and 143); m.p. 202-203 °C (lit.,^{35a,110} 204-205.5 °C or 203.5-205 °C); δ_{H} (D_2O) 3.01-3.24 (m, 8H, CH_2N), 1.91-2.26 (m, 2H, CCH_2C), 1.98 (s, 3H, CH_3CO), 1.53-1.77 (m, 4H, $\text{CCH}_2\text{CH}_2\text{C}$). δ_{C} 176.1 (CO), 49.8, 46.9, 41.1 and 39.1 (CH_2N), 28.0, 26.3 and 25.5 (CCH_2C), 24.4 (CH_3CON). (Found: C, 40.8; H, 8.8; N, 15.6. $\text{C}_9\text{H}_{21}\text{N}_3\text{O} \cdot 2\text{HCl}$ requires C, 41.54; H, 8.91; N, 16.15 %).




Figure 7 Separation of the three monoacetyl spermidine derivatives by reversed-phase ion pair chromatography using a u.v.-absorbing counter ion¹²⁶.

Mobile phase: 0.01 M phosphate buffer, pH= 2.0 / ethanol (94:6),
Flow rate 1.0 ml/min.

Counter ion: Naphthalene-2-sulfonate (0.0004 M).

Support: PLRP-S, 100 A, 5 μ m (150 x 4.6 mm I.D.).

Peaks: **144**= N⁴-Ac-spermidine; **148**= N¹-Ac-spermidine;
152= N⁸-Ac-spermidine and **S**= system peak.

Temperature: 25 °C \pm 1 °C.





Figure 8 Reversed-phase ion pair chromatography of: **144**= N^4 -Ac-spermidine; **148**= N^1 -Ac-spermidine and **152**= N^8 -Ac-spermidine. Conditions as in Fig. 7.

5.2.8 - Synthesis of N¹,N⁸-diethylspermidine trioxalate


N¹,N⁴,N⁸-Tribenzyloxycarbonyl-N¹,N⁸-diethylspermidine (162)



A solution of triprotected spermidine **131** (549 mg, 1.00 mmol) and EtI (2.50 g, 16.0 mmol) in anhydrous THF-DMF (10:1) (25 ml), was cautiously treated with NaH (80 % dispersion in oil, 180 mg, 6.00 mmol) at 0 °C under nitrogen. The resulting suspension was refluxed under nitrogen for 24 h. The yellowish, reaction mixture was cooled to r.t. (a white precipitate was formed) and water (2 ml) was added. The clear yellow solution was evaporated under reduced pressure and the residue partitioned between ether (200 ml) and 1 M KHSO₄ (100 ml). The aqueous solution was again extracted with ether (100 ml). The combined organic layers were washed and dried as usual. Evaporation to dryness gave a crude mixture which was chromatographed on silica with ether-light petroleum (2:1). The fractions containing pure compound were collected and evaporated to afford 392 mg (65 %) of product **162** as a light yellow oil, homogeneous by t.l.c. (G); δ_H, 7.32 (s, 15H, arom. H), 5.11 (s, 6H, CH₂Ph), 3.02-3.28 (m, 12H, CH₂N), 1.61-1.83 (m, 2H, CCH₂C), 1.31-1.58 (m, 4H, CCH₂CH₂C), and 1.09 (t, 6H, CH₃CH₂N); δ_C 160.0 (CO), 136.9, 136.7, 128.4, 127.8 and 127.7

(arom. C), 66.9 and 66.8 ($\underline{\text{C}}\text{H}_2\text{Ph}$), 46.5, 44.6 and 41.9 ($\underline{\text{C}}\text{H}_2\text{N}$), 27.4 and 25.6 ($\underline{\text{C}}\text{CH}_2\text{C}$), and 13.7 ($\underline{\text{C}}\text{H}_3\text{CH}_2\text{N}$).




N^1, N^8 -Diethylspermidine trioxalate (163)



Compound **162** (850 mg, 1.41 mmol) was hydrogenolyzed as described for the model compound **170** (p. 106). The crude product (265 mg, 94 %), essentially pure by t.l.c. (S), was dissolved in ethanol-ether (1:1) (5 ml) and precipitated as its oxalate salt with excess of a solution of 0.1 M oxalic acid in ether (40 ml). The white solid was centrifuged to afford 444 mg (71 %) of the salt. Recrystallization from water-ethanol (1:1) (100 ml/g) gave the pure oxalate salt **163** as light shiny white crystals, chromatographically homogeneous (S); m.p. 229.5-230.0 °C; δ_{H} (D_2O) 2.83-3.21 (m, 12H, $\underline{\text{C}}\text{H}_2\text{N}$), 1.83-2.81 (m, 2H, $\underline{\text{C}}\text{CH}_2\text{C}$), 1.56-1.80 (m, 4H, $\underline{\text{C}}\text{CH}_2\text{CH}_2\text{C}$), and 1.27 (t, 6H, $\underline{\text{C}}\text{H}_3\text{CH}_2\text{N}$); δ_{C} 168.6 (CO, oxalate), 49.7, 48.9, 47.2, 46.5, 45.7 and 45.6 ($\underline{\text{C}}\text{H}_2\text{N}$), 25.5 ($\underline{\text{C}}\text{CH}_2\text{C}$), and 13.2 ($\underline{\text{C}}\text{H}_3\text{CH}_2\text{N}$). (Found: C, 43.2; H, 7.0; N, 8.8. $\text{C}_{11}\text{H}_{27}\text{N}_3 \cdot 3\text{H}_2\text{C}_2\text{O}_4$ requires C, 43.31; H, 7.05; N, 8.91 %).

5.2.9 - Attempted syntheses of N¹-ethyl- and N⁸-ethyl-spermidines

N¹-Ethylspermidine trioxalate (164)




164

The crude N¹-Ac-spermidine dihydrochloride **148**, obtained from acidolysis of the corresponding Boc₂-derivative **147** (1.15 g, 2.97 mmol), was treated with aqueous 30 % K₂CO₃ (20 ml). After saturating with NaCl, the yellowish aqueous phase was extracted with CHCl₃ (5 x 50 ml). The combined organic layers were evaporated under reduced pressure to afford 393 mg (71 %) of the free amine as an oil, essentially pure by t.l.c. (S). This crude material was suspended in dry dioxan (40 ml) and cautiously treated with excess of Red-Al as described for model compound **174** under procedure A (p. 110). After refluxing for 4 h, t.l.c. (S) showed a major spot and a minor one with the same R_f as spermidine. The reaction mixture was cooled and worked up as described for compound **174** (p.110) but, in this case, the turbid basic aqueous solution (50 ml) was continuously extracted with CHCl₃ (150 ml) for about 6 h. The extract was dried (Na₂SO₄) and evaporated to afford 498 mg


of a yellow liquid. This crude material was chromatographed on silica (CHCl_3 -MeOH-aqueous 25 % NH_3 , 2:2:1). The fractions containing the pure product were pooled and evaporated under reduced pressure. Two phases were formed during the evaporation. To avoid bumping, it was necessary to keep the temperature below 25 °C until the chloroform evaporated (only one phase). The temperature was then increased to 40 °C to complete evaporation. A white solid was obtained which was triturated with CHCl_3 (10 ml) and concentrated (twice). The precipitate was again taken up in CHCl_3 (5 ml), filtered and rinsed with small portions of CHCl_3 . T.l.c. (S) showed that the yellowish filtrate did not contain the product and the solid (256 mg) was homogeneous. Elemental analysis indicated, however, that this material contained \approx 30 % of the desired product. This solid was dissolved in 15 % NaOH (20 ml) and after saturating with NaCl, the aqueous solution was extracted with CHCl_3 (5 x 40 ml). The combined organic layers were dried and evaporated to dryness to afford 147 mg of a yellowish liquid. The product was dissolved in ethanol-ether (1:1) (5 ml) and isolated as its oxalate salt (0.1 M oxalic acid in ether, 30 ml). The white precipitate was centrifuged to afford 218 mg (23 %) of the salt. Recrystallization from water-ethanol (1:2) (90 ml/g) afforded 197 mg (21%) of pure oxalate salt **164** as light shiny white crystals, homogeneous by t.l.c. (S); m.p. 218.5-219.0 °C; δ_{H} (D_2O) 3.05-3.18 (m, 10H, CH_2N), 2.05-2.14 (m, 2H, CCH_2C), 1.73-1.77 (m, 4H, $\text{CCH}_2\text{CH}_2\text{C}$), and 1.27 (t, 3H, $\text{CH}_3\text{CH}_2\text{N}$); δ_{C} 168.6 (CO, oxalate), 50.0, 47.4,

46.8, 46.0 and 41.7 ($\underline{\text{C}}\text{H}_2\text{N}$), 26.9, 25.7 and 25.6 ($\text{C}\underline{\text{C}}\text{H}_2\text{C}$), and 13.4 ($\underline{\text{C}}\text{H}_3\text{CH}_2\text{N}$). (Found: C, 40.7; H, 6.7; N, 9.4. $\text{C}_9\text{H}_{23}\text{N}_3 \cdot 3\text{H}_2\text{C}_2\text{O}_4$ requires C, 40.63; H, 6.59; N, 9.48 %).

N^8 -Ethylspermidine trioxalate (165)



- 1) Dioxan
- 2) aq. NaOH
- 3) $\text{C}_2\text{H}_2\text{O}_4$




165

The crude N^8 -Ac-spermidine, obtained from hydrogenolysis of the corresponding Z_2 -derivative **151** (1.17 g, 2.57 mmol), was treated with Red-Al and the product worked up and purified in a similar manner as described above for compound **164**. The yield of the pure oxalate salt **165**, as light shiny crystals, was 252 mg (22 %); m.p. 212.5–213.0 °C; δ_{H} (D_2O) 3.04–3.17 (m, 10H, $\underline{\text{C}}\text{H}_2\text{N}$), 2.05–2.10 (m, 2H, $\text{C}\underline{\text{C}}\text{H}_2\text{C}$), 1.74–1.77 (m, 4H, $\text{C}\underline{\text{C}}\text{H}_2\text{CH}_2\text{C}$), and 1.26 (t, 3H, $\underline{\text{C}}\text{H}_3\text{CH}_2\text{N}$); δ_{C} 168.4 (CO, oxalate), 49.7, 48.9, 47.2, 45.6 and 39.2 ($\underline{\text{C}}\text{H}_2\text{N}$), 26.4 and 25.5 ($\text{C}\underline{\text{C}}\text{H}_2\text{C}$), and 13.2 ($\underline{\text{C}}\text{H}_3\text{CH}_2\text{N}$). (Found: C, 40.0; H, 6.7; N, 9.2. $\text{C}_9\text{H}_{23}\text{N}_3 \cdot 3\text{H}_2\text{C}_2\text{O}_4$ requires C, 40.63; H, 6.59; N, 9.48 %).

5.3 - Synthesis of alkyl benzyl imidodicarbonates

5.3.1- Benzyloxycarbonyl isocyanate (154)




To a suspension of dry, finely ground benzyl carbamate¹¹⁶, **153**, (30.8 g, 0.20 mol) in dry CH₂Cl₂ (300 ml) was added dropwise with efficient stirring a solution of oxalyl chloride (38.1 g, 0.30 mol) in dry CH₂Cl₂ (150 ml) over a period of 1 h with ice-cooling under dry nitrogen. The initially clear mixture gradually became turbid after stirring for 1 h at 0 °C. The stirring was continued for 4 h at r.t. and overnight (15 h) under reflux. The mixture was then concentrated to about 2/3 of its original volume and the fine-grained precipitate was filtered off and washed with cold, dry CH₂Cl₂ (3 x 20 ml). The combined, pale yellow filtrate was evaporated to dryness at r.t. with a minimum exposure to atmospheric moisture. The residual yellowish turbid oil was distilled at reduced pressure under nitrogen. After a forerun, consisting largely of oxalyl chloride and benzyl chloride, product **154** was collected at 78-80 °C/ 0.15-0.20 mm Hg. The yield of **154** was 18.9 g (53 %). This material, a colourless liquid, containing <1% of benzyl chloride, was suitable for further work, except in the synthesis of **155d** which required the removal of remaining traces of acidic impurities by a second distillation. This compound being very sensitive to moisture

was stored below $-20\text{ }^{\circ}\text{C}$ in a sealed vessel; δ_{H} 7.38 (s, 5H, arom. H), 5.20 (s, 2H, CH_2Ph); δ_{C} 148.6 (PhCH_2OCON), 129.5 ($\text{N}=\text{C}=\text{O}$), 133.6, 128.5, 128.2 and 128.1 (arom. C), and 70.2 (CH_2Ph).

The white precipitate from above, weighing 4.9 g, consisted largely of N,N'-dibenzoyloxycarbonyloxamide (for related derivatives, see Lit.¹¹⁸); m.p. 211-212 $^{\circ}\text{C}$ (dec., from 1,2-dichloroethane ($\approx 1\text{ l/g}$)). The crystalline, analytical specimen tenaciously retains the solvent, even after drying in high vacuo for several days; δ_{H} (DMSO-d_6) 11.60 (broad, $\approx 2\text{H}$, NH), 7.38 (s, 10H, arom. H), and 5.17 (s, 4H, CH_2Ph); δ_{C} 165.0 (CO-CO), 152.9 (PhCH_2OCON), 135.3, 128.7 and 128.4 (arom. C), and 67.5 (CH_2Ph).

5.3.2- Alkyl benzyl imidodicarbonates (155a-i)

General procedure



To a vigorously stirred solution of the alcohol ROH (20 mmol) in dry CH_2Cl_2 (30 ml), under nitrogen and cooled in ice, a solution of benzyloxycarbonyl isocyanate, **154** (3.10 g, 19 mmol in the preparations of **155a**, **b**, **d**, **e**, **g**, **h** involving volatile alcohols; 3.43 g, 21 mmol for **155c**, **f**, **i**), in dry CH_2Cl_2 (30 ml) was added over a period of 30 min. In the preparation of **155d**, the crude **154** is redistilled before use.

After the addition, the resulting colourless solution was stirred under nitrogen in the ice-bath for 1 h and overnight at room temperature. The solvent was evaporated and, except for **155e**, the residue was thoroughly triturated with cold light petroleum (\approx 30 ml, for **155c** and **155i** cold ether). After several hours in the cold the white crystalline solid was collected by filtration, rinsed with small portions of cold solvent and dried over paraffin chips at reduced pressure.

The analytical samples were obtained by recrystallization (for **155e** by chromatography) and the details of the purification and properties of **155a-i** are compiled in Tables XVII and XVIII.

Table XVII - Alkyl benzyl imidodicarbonates **155a-i** prepared.

Com-pound	Yield ^a (%)	Solvent for recrystallization	m.p. /°C	Elemental analysis or Lit. m.p. /°C
155a	99	CH ₂ Cl ₂ -Et ₂ O (1:7, 50 ml/g)	109-109.5	105.5-106.5 ¹¹⁹
155b	100	See 155a	92.5-93	Found: C, 64.8; H, 5.3; N, 4.4 C ₁₇ H ₁₇ N ₂ O ₅ requires: C, 64.75; H, 5.43; N, 4.44 %
155c	97	CH ₂ Cl ₂ -Et ₂ O (1:4, 60 ml/g)	113.5-114	Found: C, 58.2; H, 4.2; N, 8.4 C ₁₆ H ₁₄ N ₂ O ₆ requires: C, 58.18; H, 4.27; N, 8.48 %

155d	91b	Et ₂ O-light petroleum (1:4, 30 ml/g) ^c	83.5-84	Found: C, 68.9; H, 6.2; N, 4.5 C ₁₈ H ₁₉ NO ₄ requires: C, 69.00; H, 6.11; N, 4.47 %
155e	97 ^d	purified by chromatography	oil	oil ¹⁰¹
155f	98	Et ₂ O (10 ml/g)	112-112.5	Found: C, 69.2; H, 7.0; N, 4.3 C ₁₉ H ₂₃ NO ₄ requires: C, 69.28; H, 7.04; N, 4.25 %
155g	94	Et ₂ O-hexane (1:1, 20 ml/g)	79.5-80	Found: C, 61.2; H, 5.5; N, 6.0 C ₁₂ H ₁₃ NO ₄ requires: C, 61.27; H, 5.57; N, 5.95 %
155h	92	See 155d	90-90.5	Found: C, 40.5; H, 2.9; N, 4.2 C ₁₁ H ₁₀ Cl ₃ NO ₄ requires: C, 40.46; H, 3.09; N, 4.29 %
155i	92	CH ₂ Cl ₂ -Et ₂ O (1:10, 70 ml/g)	112-113 ^e	Found: C, 74.1; H, 5.0; N, 3.8 C ₂₃ H ₁₉ NO ₄ requires: C, 73.98; H, 5.13; N, 3.75 %

^a Yield of essentially pure product before recrystallization (<1% of impurities as judged from ¹H n.m.r. and t.l.c. (A), (G), or ether).

^b Yield corrected for ≈ 4 % of **153** in the crude product.

^c Recrystallized product contained traces of **153**. Attempted chromatography on silica caused partial decomposition of the compound.

^d Crude product contained 1-2% of **153**. Yield after chromatography (silica light petroleum-ether, 3:1) 84 %.

^e Softens at ≈ 107 °C.

155e	as previously reported ¹⁰¹	as previously reported ¹⁰¹
155f	1.65 and 2.12 (2 br sign, 15H, H _{alif}); 5.17 (s, 2H, CH ₂); ≈7.10 (br s, ≈1H, NH); 7.34 (s, 5H _{arom}).	30.9, 36.0, 41.2 (C _{alif,Adoc}); 67.4 (CH ₂); 82.4 (C _{quart}); 128.4, 128.5, 128.6, 135.2 (C _{arom}); 148.9 (CO _{Adoc}); 150.9 (CO _Z).
155g	4.60 and 4.67 (2t, 2H, CH ₂ CH); 5.17 [s, 2H, CH ₂ (Z)]; 5.30 and 5.30 (2m, 2H, =CH ₂); 5.70-6.12 (m, 1H, =CH-); 7.34 (s, 5H _{arom}); ≈7.50 (br s, ≈1H, NH).	66.8 (CH ₂ -CH); 67.8 [CH ₂ (Z)]; 119.1 (=CH ₂); 128.5, 128.6, 134.9 (C _{arom}); 131.3 (=CH-); 150.5, 150.6 (CO).
155h	4.78 [s, 2H, CH ₂ (Troc)]; 5.22 [s, 2H, (Z)]; 7.37 (s, 5H _{arom}); ≈7.53 (br s, ≈1H, NH).	68.3 [CH ₂ , (Z)]; 74.7 [CH ₂ (Troc)]; 94.3 (CCl ₃); 128.6, 128.7, 128.8, 134.6 (C _{arom}); 149.0 (CO _{Troc}); 150.4 (CO _Z).
155i	4.14-4.28 (4 sign, 1H _{alif} , (Fmoc)); 4.45-4.53 [3 sign, 2H, CH ₂ (Fmoc)]; 5.19 [s,	46.6 [C _{alif} (Fmoc)]; 67.8 (CH ₂); 128.5, 128.6, 134.8 [C _{arom} Z]; 120.0,

2H, CH₂ (Z)]; 7.18-7.80 (m) 124.9, 127.1, 127.9,
+ 7.35 (s, together 14H, 141.2, 142.2 [C_{arom}
H_{arom} + NH) (Fmoc)]; 150.5, 150.7
(CO).

Z= Benzyloxycarbonyl
Z(OMe)= 4-Methoxybenzyloxycarbonyl
Z(NO₂)= 4-Nitrobenzyloxycarbonyl
Ppoc≡ 2-Phenylisopropylloxycarbonyl
Adoc= 1-Adamantylloxycarbonyl
Troc= 2,2,2-Trichloroethoxycarbonyl
Fmoc= 9-Fluorenylmethoxycarbonyl

Model compounds synthesized

	page	ref.
N ² -Benzyloxycarbonyl-N ¹ - <u>tert</u> -butoxycarbonyl-N ¹ -ethylethylenediamine 157	103	-
N ¹ -Benzyloxycarbonyl-N ² - <u>tert</u> -butoxycarbonyl-N ¹ ,N ² -diethylethylenediamine 167	104	-
N ¹ - <u>tert</u> -Butoxycarbonyl-N ¹ -ethylenediamine 170	106	-
N ² -Acetyl-N ¹ - <u>tert</u> -butoxycarbonyl-N ¹ -ethyl-ethylenediamine 171	107	-
N ¹ - <u>tert</u> -Butoxycarbonyl-N ¹ ,N ² -diethylethylenediamine 172	108	-
N ² -Acetyl-N ¹ -ethylethylenediamine 173	109	-
N ¹ ,N ² -Diethylethylenediamine dihydrochloride 174	110	-
N ¹ -Benzyloxycarbonyl-N ⁴ - <u>tert</u> -butoxycarbonyl-putrescine 168	112	123

Spermidine derivatives synthesized

	page	ref.
N^1, N^4, N^4 -Tribenzyloxycarbonylspermidine 131 .	113	109a, 109b
N^1, N^4, N^8 -Tribenzyloxycarbonyl- N^1, N^8 -bis(<u>tert</u> -butoxycarbonyl)spermidine 132	114	109a, 109b
N^1, N^8 -Bis(<u>tert</u> -butoxycarbonyl)spermidine 133	116	19, 41, 51b, 82, 109a, 109b
N^4 -Benzyloxycarbonyl- N^1, N^8 -bis(<u>tert</u> -butoxycarbonyl)spermidine 134	117	109a, 109b
N^1, N^4 -Methylenespermidine 114b	119	18, 79, 80 109b
N^1, N^8 -Dibenzyloxycarbonyl- N^1, N^4 -methylene-spermidine 135	119	109b
N^1, N^8 -Dibenzyloxycarbonyl- N^8 - <u>tert</u> -butoxycarbonyl- N^1, N^4 -methylenespermidine 136	120	109b
N^1 -Benzyloxycarbonyl- N^8 - <u>tert</u> -butoxycarbonyl- N^1, N^4 -methylenespermidine 137	121	109b
N^1 -Benzyloxycarbonyl- N^8 - <u>tert</u> -butoxycarbonyl-spermidine 138	121	121, 109b
N^1, N^8 -Dibenzyloxycarbonyl- N^8 - <u>tert</u> -butoxycarbonylspermidine 139	123	109b
N^1, N^4, N^8 -Triacetylspermidine 141	130	125, 109c
N^1, N^4, N^8 -Triacetyl- N^1, N^8 -bis(<u>tert</u> -butoxycarbonyl)spermidine 142	131	109c
N^4 -Acetyl- N^1, N^8 -bis(<u>tert</u> -butoxycarbonyl)-spermidine 143	133	109c
N^4 -Acetylspermidine dioxalate 144a	134	109c
N^1 -Benzyloxycarbonyl- N^4, N^8 -bis(<u>tert</u> -butoxycarbonyl)spermidine 145	135	121, 109c

N^4, N^8 -Bis(<u>tert</u> -butoxycarbonyl)spermidine 146	136	36, 37, 48, 51a, 121, 109c
N^1 -Acetyl- N^4, N^8 -bis(<u>tert</u> -butoxycarbonyl)- spermidine 147	137	109c
N^1 -Acetylspermidine dihydrochloride 148	137	35, 55, 83, 109c
N^1, N^4 -Dibenzyloxycarbonyl- N^8 - <u>tert</u> -butoxy- carbonylspermidine 149	138	109c
N^1, N^4 -Dibenzyloxycarbonylspermidine 150	140	109c
N^8 -Acetyl- N^1, N^4 -dibenzyloxycarbonyl- spermidine 151	140	109c
N^8 -Acetylspermidine dihydrochloride 152	141	35a, 55, 83, 109c
N^1, N^4, N^8 -Tribenzyloxycarbonyl- N^1, N^8 -diethyl- spermidine 162	144	-
N^1, N^8 -Diethylspermidine trioxalate 163	145	-
N^1 -Ethylspermidine trioxalate 164	146	-
N^8 -Ethylspermidine trioxalate 165	148	-

Alkyl benzyl imidodicarbonates synthesized

	page	ref.
Dibenzyl imidodicarbonate 155a	150	119, 109d
Benzyl <i>p</i> -methoxybenzyl imidodicarbonate 155b ..	150	109d
Benzyl <i>p</i> -nitrobenzyl imidodicarbonate 155c	150	109d
Benzyl 2-phenylisopropyl imidodicarbonate 155d	150	109d
Benzyl <i>tert</i> -butyl imidodicarbonate 155e	150	101, 109d
1-Adamantyl benzyl imidodicarbonate 155j	150	109d
Allyl benzyl imidodicarbonate 155g	150	109d
Benzyl 2,2,2-trichloroethyl imidodi- carbonate 155h	150	109d
Benzyl 9-fluorenylmethyl imidodicarbonate 155i	150	109d

REFERENCES

- 1 D. M. L. Morgan, Essays Biochem., 1987, **23**, 82.
- 2 A. E. Pegg, Biochem. J., 1986, **234**, 249.
- 3 C. W. Tabor and H. Tabor, Ann. Rev. Biochem., 1984, **53**, 749.
- 4 C. Porter and J. R. Sufrin, Anticancer Res., 1986, **6**, 525.
- 5 A. E. Pegg, Cancer Res., 1988, **48**, 759.
- 6 "Inhibition of Polyamine Metabolism: Biological Significance and Basis for New Therapies", eds. P. P. McCann, A. E. Pegg and A. Sjoerdsma, Academic Press, New York, 1987, 371 pp.
- 7 N. Seiler, Can. J. Physiol. Pharmacol., 1987, **65**, 2024.
- 8 a) M. Tsuboi, Bull. Chem. Soc. Jpn., 1964, **37**, 1514; b) B. G. Feuerstein, N. Pattabiraman and L. J. Marton, Proc. Natl. Acad. Sci. USA, 1986, **83**, 5948; c) J. E. Morgan, J. W. Blankenship and H. R. Matthews, Arch. Biochem. Biophys., 1986, **246**, 225; d) J. E. Morgan, J. W. Blankenship and H. R. Matthews, Biochemistry, 1987, **26**, 3643; e) P. M. Vertino, R. J. Bergeron, P. F. Cavanaugh, Jr. and C. W. Porter, Biopolymers, 1987, **26**, 691.
- 9 a) C. W. Porter, R. J. Bergeron and N. J. Stolowich, Cancer Res., 1982, **42**, 4072; b) R. A. Casero, Jr., R. J. Bergeron and C. W. Porter, J. Cell. Physiol., 1984, **121**, 476; c) C. W. Porter, J. Miller and R. J. Bergeron, Cancer Res., 1984, **44**, 126; d) C. W. Porter, P. F. Cavanaugh, Jr., N. Stolowich, B. Ganis, E. Kelly and R. J. Bergeron, Cancer Res., 1985, **45**, 2050; e) B. G. Erwin and A. E. Pegg, Biochem. J., 1986, **238**, 581; f) C. W. Porter, B. Ganis, T. Vinson, L. J. Marton, D. L. Kramer and R. J. Bergeron, Cancer Res., 1986, **46**, 6279; g) C. W. Porter, F. G. Berger, A. E. Pegg, B. Ganis and R. J. Bergeron, Biochem. J., 1987, **242**, 43; h) C. W. Porter, J. McManis, R. A. Casero and R. J. Bergeron, Cancer Res., 1987, **47**, 2821; i) R. A. Casero, Jr., B. Go, H. W. Theiss, J. Smith, S. B. Baylin and G. D. Luk, Cancer Res. 1987, **47**, 3964; j) S. C. Denstman, S. J. Ervin and R. A. Casero, Jr., Biochem. Biophys. Res. Commun., 1987, **149**, 194; k) A. E. Pegg, R. Madhubala, T. Kameji and R. J. Bergeron, J. Biol. Chem., 1988, **263**, 11008; l) R. J. Bergeron, A. H. Neims, J. S. McManis, T. R. Hawthorne, J. R. T. Vinson, R. Bortell and M. J. Ingenu, J. Med. Chem.,

- 1988, **31**, 1183; m) R. A. Casero, Jr., S. J. Ervin, P. Celano, S. B. Baylin and R. J. Bergeron, Cancer Res., 1989, **49**, 639.
- 10 G. A. Ellestad, D. B. Cosulich, R. W. Broschard, J.H. Martin, M. P. Kunstmann, G. O. Morton, J. E. Lancaster, W. Fulmor and F. M. Lovell, J. Am. Chem. Soc., 1978, **100**, 2515.
- 11 T. Kosaki, T. Ikoda, Y. Kotani, S. Nakagawa and T. Saka, Science, 1958, **127**, 1176.
- 12 G. B. Henderson, P. Ulrich, A. H. Fairlamb and A. Cerami, J. Chem. Soc., Chem. Commun., 1986, 593.
- 13 a) K. Wiesner, D. M. MacDonald and C. Bankiewicz, J. Am. Chem. Soc., 1953, **75**, 6348; b) G. Englert, K. Klinga, R. Hamet, E. Schlittler and W. Vetter, Helv. Chim. Acta, 1973, **56**, 474; c) S. M. Kupchan, A. P. Davies, S. J. Barboutis, H. K. Schoes and A. L. Burlingame, J. Am. Chem. Soc., 1967, **89**, 5718.
- 14 a) S. A. Ong, T. Peterson and J. B. Neilands, J. Biol. Chem., 1979, **254**, 1860; b) F. L. Weitzl and K. N. Raymond, J. Org. Chem., 1981, **46**, 5234; c) S. Funayama, K. Yoshida, C. Konno and H. Hikino, Tetrahedron Lett., 1980, **21**, 1355.
- 15 G. T. Carter and K. L. Rinehart, Jr., J. Am. Chem. Soc., 1978, **100**, 4302.
- 16 K. Umezawa and T. Takeuchi, Biomedicine & Pharmacotherapy 1987, **41**, 227.
- 17 a) J. B. Hansen, M. C. Nielsen, U. Ehrbar and O. Buchardt, Synthesis, 1982, 404; b) G. J. Atwell and W. A. Denny, Synthesis, 1984, 1032; c) A. R. Jacobson, A. N. Makris and L. M. Sayre, J. Org. Chem., 1987, **52**, 2592.
- 18 B. Ganem, Acc. Chem. Res., 1982, **15**, 290.
- 19 R. J. Bergeron, Acc. Chem. Res., 1986, **19**, 105.
- 20 J. R. Malpass, "Comprehensive Organic Chemistry: The Synthesis and Reactions of Organic Compounds", ed. I. O. Sutherland, Pergamon Press, Oxford, 1979, Vol. 2, p. 3.
- 21 J. R. Piper and T. P. Johnston, J. Org. Chem., 1968, **33**, 636.
- 22 a) A. Guggisberg, P. van den Broek, M. Hess, H. Schmid, F. Schneider and K. Bernauer, Helv. Chim. Acta, 1976, **59**, 3013; b) A. Guggisberg, R. W. Gray and M. Hesse, Helv.

- Chim. Acta, 1977, **60**, 112; c) B. M. Trost and J. Cossy, J. Am. Chem. Soc., 1982, **104**, 6881.
- 23 E. Walchli-Schaer and C. H. Eugster, Helv. Chim. Acta, 1978, **61**, 928.
- 24 G. von Kiedrowski and F. Z. Dörwald, Liebigs Ann. Chem., 1988, 787.
- 25 T. W. Greene, "Protective Groups in Organic Chemistry", J. Wiley and Sons, New York, 1981, p. 218.
- 26 a) N. L. Benoiton and J. R. Coggins, "Progress in Peptide Research", ed. S. Lande, Gordon and Breach, New York, 1972, Vol. 2, p.22-1; b) S. T. Cheung and N. L. Benoiton, Can. J. Chem., 1977, **55**, 906.
- 27 P. D. Croce, C. La Rosa and A. Ritieni, J. Chem. Res. (S), 1988, 346.
- 28 R. Sulsky and J. P. Demers, Tetrahedron Lett., 1989, **30**, 31.
- 29 a) K. Samejima, Y. Takeda, M. Kawase, M. Okada and Y. Kyogoku, Chem. Pharm. Bull., 1984, **32**, 3428; b) M. Niitsu and K. Samejima, Chem. Pharm. Bull., 1986, **34**, 1032.
- 30 T. Ando and J. Yamawaki, Chem. Lett., 1979, 45.
- 31 F. Ramiandrasoa, M.-L. Milat, G. Kunesch and S. Chuilon, Tetrahedron Lett., 1989, **30**, 1365.
- 32 G. Kunesch, Tetrahedron Lett., 1983, **24**, 5211.
- 33 M. Hudlicky, "Reductions in Organic Chemistry", J. Wiley and Sons, New York, 1983, 309 pp.
- 34 M. Israel, J. S. Rosenfield and E. J. Modest, J. Med. Chem., 1964, **7**, 710.
- 35 a) H. Tabor, C. W. Tabor and L. de Meis, Methods Enzymol., 1971, **17B**, 829; c) H. Tabor and C. W. Tabor, Methods Enzymol., 1983, **94**, 420.
- 36 a) M. Humora and J. Quick, J. Org. Chem., 1979, **44**, 1166; b) M. J. Humora, D. E. Seitz and J. Quick, Tetrahedron Lett., 1980, **21**, 3971.
- 37 H.- M. Shieh, D. Campbell and K. Folkers, Biochem. Biophys. Res. Commun., 1984, **122**, 21.
- 38 R. Walchli, A. Guggisberg and M. Hesse, Helv. Chem. Acta, 1984, **67**, 2178.

- 39 T. L. Shih, J. Ruiz-Sanchez and H. Mrozik, Tetrahedron Lett., 1987, **28**, 6015.
- 40 a) R. J. Bergeron, K. A. McGovern, M. A. Channing and P. S. Burton, J. Org. Chem., 1980, **45**, 1589; b) R. J. Bergeron, P. S. Burton, K. A. MacGovern and S. J. Kline, Synthesis, 1981, 732.
- 41 R. J. Bergeron and N. J. Stolowich, Synthesis, 1982, 689.
- 42 a) R. J. Bergeron, P. S. Burton, S. J. Kline and K. A. McGovern, J. Org. Chem., 1981, **46**, 3712; b) R. J. Bergeron, S. J. Kline, N. J. Stolowich, K. A. McGovern and P. S. Burton, J. Org. Chem., 1981, **46**, 4524; c) R. J. Bergeron and S. J. Kline, J. Am. Chem. Soc., 1982, **104**, 4489; d) R. J. Bergeron, N. J. Stolowich and S. J. Kline, J. Org. Chem., 1983, **48**, 3432; e) R. J. Bergeron, J. S. McManis, J. B. Dionis and J.R. Garlich, J. Org. Chem., 1985, **50**, 2780.
- 43 R. J. Bergeron, J. R. Garlich and N. J. Stolowich, J. Org. Chem., 1984, **49**, 2997.
- 44 R. J. Bergeron and J. S. McManis, J. Org. Chem., 1988, **53**, 3108.
- 45 S. Sakakibara and N. Inukai, Bull. Chem. Soc. Jpn., 1965, **38**, 1979.
- 46 H. Eckert, M. Listl and I. Ugi, Angew. Chem., Int. Ed. Engl., 1978, **17**, 361.
- 47 R. J. Bergeron, J. R. Garlich and J. S. McManis, Tetrahedron, 1985, **41**, 507.
- 48 R. J. Bergeron and J. S. McManis, J. Org. Chem., 1987, **52**, 1700.
- 49 R. J. Bergeron and J. R. Garlich, Synthesis, 1984, 782.
- 50 J. E. Nordlander, M. J. Payne, M. A. Balk, J. L. Gress, F. D. Harris, J. S. Lane, R. F. Lewe, S. E. Marshall, D. Nagy and D. J. Rachlin, J. Org. Chem., 1984, **49**, 133.
- 51 a) R. Sundaramoorthi, J.-L. Fourrey and B. C. Das, J. Chem. Soc., Perkin Trans. 1, 1984, 2759; b) R. Sundaramoorthi, C. Marazano, J.-L. Fourrey and B. C. Das, Tetrahedron Lett., 1984, **25**, 3191.
- 52 W. V. Curran and R. B. Angier, J. Org. Chem., 1966, **31**, 3867.

- 53 N. Umino, T. Iwakuma and N. Itoh, Tetrahedron Lett., 1976, 763.
- 54 B. T. Golding, M. C. O'Sullivan and L. L. Smith, Tetrahedron Lett., 1988, **29**, 6651.
- 55 a) J. Boukouvalas, B. T. Golding, R. W. McCabe and P. K. Slaich, Angew. Chem., Int. Ed. Engl., 1983, **22**, 618; b) J. Boukouvalas, B. T. Golding, R. W. McCabe and P. K. Slaich, Angew. Chem. Suppl., 1983, 860.
- 56 S. C. Yorke, J. W. Blunt, M. H. G. Munro, J. C. Cook and K. L. Rinehart Jr, Aust. J. Chem., 1986, **39**, 447.
- 57 a) R. F. Borch and A. I. Hassid, J. Org. Chem., 1972, **37**, 1673; b) Y. Ohfune, N. Kurokawa, N. Higuchi, M. Saito, M. Hashimoto and T. Tanaka, Chem. Lett., 1984, 441; c) M. Hayashi, M. Yhara and Y. Sawazaki, Banyu Pharmaceutical Co., Japanese Patent, 62, 26,255 (1987), Chem. Abstr., 1987, **107**, 97120c.
- 58 a) K. A. Schellenberg, J. Org. Chem., 1963, **28**, 3259; b) D. B. Olsen, T. W. Hepburn, M. Moos, P. S. Mariano and D. Dunaway-Mariano, Biochemistry, 1988, **27**, 2229.
- 59 P. C. Unangst, D. T. Connor and S. R. Stabler, J. Heterocycl. Chem., 1987, **24**, 817.
- 60 a) J. March, "Advanced Organic Chemistry", J. Wiley and Sons, New York, 3rd ed., 1985, p. 796-800; b) S. Dayagi and Y. Degani, "The Chemistry of C-N Double Bond", ed. S. Patai, J. Wiley and Sons, New York, 1970, p. 64-83; c) R. L. Reeves, "The Chemistry of the Carbonyl Group", ed. S. Patai, J. Wiley and Sons, 1966, p. 600-614; d) R. W. Layer, Chem. Rev., 1963, 489.
- 61 F. Rolla, J. Org. Chem., 1982, **47**, 4327.
- 62 a) H. C. Brown, M. M. Midland and A. B. Levy, J. Am. Chem. Soc., 1973, **95**, 2394; b) H. C. Brown, M. M. Midland, A. B. Levy, A. Suzuki, S. Sono and M. Itoh, Tetrahedron, 1987, **43**, 4079.
- 63 B. Carboni, M. Vaultier and R. Carrié, Tetrahedron Lett., 1988, **29**, 1279.
- 64 B. Carboni, M. Vaultier and R. Carrié, Tetrahedron, 1987, **43**, 1799.
- 65 a) H. C. Brown, N. Ravindran and S. U. Kulkarni, J. Org. Chem., 1980, **45**, 384; b) H. C. Brown and J. B. Campbell, Jr., J. Org. Chem., 1980, **45**, 389; c) H. C. Brown, P. K. Jadhav and M. C. Desai, J. Am. Chem. Soc., 1982, **104**,

- 66 a) T. Kunieda, T. Higuchi, Y. Abe and M. Hirobe, Tetrahedron Lett., 1980, **21**, 3065; b) Y. Abe and T. Kunieda, Tetrahedron Lett., 1979, **52**, 5007; c) T. Kunieda, Y. Abe, Y. Iitaka and M. Hirobe, J. Org. Chem., 1982, **47**, 4291; d) T. Kunieda, T. Higuchi, Y. Abe and M. Hirobe, Tetrahedron Lett., 1982, **23**, 1159; e) T. Kunieda, Y. Abe, T. Higuchi and M. Hirobe, Tetrahedron Lett., 1981, **22**, 1257; f) T. Kunieda, T. Higuchi, Y. Abe and M. Hirobe, Tetrahedron, 1983, **39**, 3253.
- 67 a) Y. Nagao, K. Kawabata, K. Seno and E. Fujita, J. Chem. Soc., Perkin Trans. 1, 1980, 2470; b) E. Fujita, Y. Nagao, K. Seno, S. Takao, T. Miyasaka, M. Kimura and W. H. Watson, J. Chem. Soc., Perkin Trans. 1, 1981, 914.
- 68 a) Y. Nagao, K. Seno, K. Kawabata, T. Miyasaka, S. Takao and E. Fujita, Tetrahedron Lett., 1980, **21**, 841; b) Y. Nagao, K. Seno, T. Miyasaka and E. Fujita, Chem. Lett., 1980, 159; c) Y. Nagao, K. Seno and E. Fujita, Tetrahedron Lett., 1980, **21**, 4931; d) Y. Nagao, T. Miyasaka, K. Seno, M. Yagi and E. Fujita, Chem. Lett., 1981, 463; e) E. Fujita, Pure Appl. Chem., 1981, **53**, 1141; f) Y. Nagao, T. Ikeda, M. Yagi, E. Fujita and M. Shiro, J. Am. Chem. Soc., 1982, **104**, 2079; g) Y. Nagao and E. Fujita, Heterocycles, 1982, **17**, 537; h) Y. Nagao, S. Takao, E. Fujita, C. Murayama, T. Mori, T. Asao and T. Suzue, Experientia, 1983, **39**, 1116; i) Y. Nagao, K. Seno, K. Kawabata, T. Miyasaka, S. Takao and E. Fujita, Chem. Pharm. Bull., 1984, **32**, 2687; j) Y. Nagao, T. Miyasaka, Y. Hagiwara and E. Fujita, J. Chem. Soc., Perkin Trans. 1, 1984, 183.
- 69 A. V. Joshua and J. R. Scott, Tetrahedron Lett., 1984, **25**, 5725.
- 70 S. K. Sharma, M. J. Miller and S. M. Payne, J. Med. Chem., 1989, **32**, 357.
- 71 F. Acher and M. Wakselman, J. Org. Chem., 1984, **49**, 4133.
- 72 a) S. M. Beaumont, B. O. Handford, J. H. Jones and G. T. Young, J. Chem. Soc., Chem. Commun., 1965, 53; b) B. O. Handford, J. H. Jones, G. T. Young and T. F. N. Johnson, J. Chem. Soc., 1965, 6814; c) J. H. Jones and G. T. Young, J. Chem. Soc. (C), 1968, 53.
- 73 a) H.-P. Husson, C. Poupat, B. Rodriguez and P. Potier, Tetrahedron, 1973, **29**, 1405; b) C. Poupat, Tetrahedron Lett., 1976, **20**, 1669; c) H.-P. Husson, C. Poupat and P. Potier, C. R. Acad. Sc., Paris, (C), 1973, **276**, 1039; d) A. Husson, R. Besselievre and H.-P. Husson, Tetrahedron

- Lett., 1983, **24**, 1031.
- 74 G. Sosnovsky and J. Lukszo, Z. Naturforsch., B, 1986, **41**, 122.
- 75 S.-I. Murahashi, T. Naota and E. Saito, J. Am. Chem. Soc., 1986, **108**, 7846.
- 76 S.-I. Murahashi, T. Naota and N. Nakajima, Chem. Lett., 1987, 879.
- 77 a) S.-I. Murahashi, T. Naota and N. Nakajima, Tetrahedron Lett., 1985, **26**, 925; b) S. Hunig and R. Schaller, Angew. Chem., Int. Ed. Engl., 1982, **21**, 36.
- 78 M. E. Childs and W. P. Weber, J. Org. Chem., 1976, **41**, 3486.
- 79 J. S. McManis and B. Ganem, J. Org. Chem., 1980, **45**, 2041.
- 80 a) K. Chantrapromma, J. McManis and B. Ganem, Tetrahedron Lett., 1980, **21**, 2475; b) B. Ganem and K. Chantrapromma, Methods Enzymol., 1983, **94**, 416.
- 81 K. Chantrapromma and B. Ganem, Tetrahedron Lett., 1981, **22**, 23.
- 82 S. Nagarajan and B. Ganem, J. Org. Chem., 1985, **50**, 5735.
- 83 C. M. Tice and B. Ganem, J. Org. Chem., 1983, **48**, 2106.
- 84 K. Chantrapromma, J. S. McManis and B. Ganem, Tetrahedron Lett., 1980, **21**, 2605.
- 85 F. J. Schmitz, K. H. Hollenbeak and R. S. Prasad, Tetrahedron Lett., 1979, **20**, 3387.
- 86 N. L. Allinger, M. P. Cava, D. C. Dejongh, C. R. Johnson, N. A. Lebel and C. L. Stevens, "Organic Chemistry", 2nd ed., Worth Publishers, Inc., New York, 1976, p. 539-540.
- 87 G. Höfle, W. Steglich and H. Vorbrüggen, Angew. Chem., Int. Ed. Engl., 1978, **17**, 569.
- 88 DMAP Update, Reilly Report, Reilley Tar & Chem. Corp. Indianapolis, IN, 1982.
- 89 E. F. V. Scriven, Chem. Soc. Rev., 1983, **12**, 129.
- 90 R. B. Moodie and P. S. Sansom, J. Chem. Soc., Perkin Trans.2, 1981, 664.

- 91 A. R. Fersht and W. P. Jencks, J. Am. Chem. Soc., 1970, **92**, 5432.
- 92 M. Wakselman and E. Guibé-Jampel, J. Chem. Soc., Chem. Commun., 1976, 21.
- 93 E. Guibé-Jampel and M. Wakselman, J. Chem. Soc., Chem. Commun., 1980, 993.
- 94 O. Hernandez, S. K. Chaudhary, R. H. Cox and J. Porter, Tetrahedron Lett., 1981, **22**, 1491.
- 95 D. L. Flynn, R. E. Zelle and P. A. Grieco, J. Org. Chem., 1983, **48**, 2424.
- 96 D. S. Tarbell, Y. Yamamoto and B. M. Pope, Proc. Natl. Acad. Sci. USA, 1972, **69**, 730.
- 97 L. Grehn, K. Gunnarsson and U. Ragnarsson, Acta Chem. Scand., Ser. B, 1986, **40**, 745.
- 98 L. Grehn, K. Gunnarsson and U. Ragnarsson, J. Chem. Soc., Chem. Commun., 1985, 1317.
- 99 L. Grehn, K. Gunnarsson and U. Ragnarsson, Acta Chem. Scand., Ser. B, 1987, **41**, 18.
- 100 L. Grehn and U. Ragnarsson, Synthesis, 1987, 275.
- 101 L. Grehn and U. Ragnarsson, Collect. Czech. Chem. Commun., 1988, **53**, 2778.
- 102 M. S. Gibson and R. W. Bradshaw, Angew. Chem., Int. Ed. Engl., 1968, **7**, 919.
- 103 O. Mitsunobu, Synthesis, 1981, 1.
- 104 P. Kočovský, Tetrahedron Lett., 1986, **27**, 5521.
- 105 G. Brieger and T. J. Nestrick, Chem. Rev., 1974, **74**, 567.
- 106 M. K. Anwer and A. F. Spatola, Synthesis, 1980, 929.
- 107 a) R. F. Evans, Aust. J. Chem., 1967, **20**, 1643; b) L. Duhamel, "The Chemistry of Functional Groups" Supplement F: The chemistry of amino, nitroso and nitro compounds and their derivatives, ed. S. Patai, J. Wiley and Sons, New York, 1982, p. 849.
- 108 E. Wunsch, Synthesis, 1986, 958.
- 109 a) M. L. S. Almeida, L. Grehn and U. Ragnarsson, J. Chem. Soc., Chem. Commun., 1987, 1250; b) M. L. S. Almeida, L.

- Grehn and U. Ragnarsson, J. Chem. Soc., Perkin Trans. 1, 1988, 1905; c) M. L. S. Almeida, L. Grehn and U. Ragnarsson, Acta Chem. Scand., Ser. B, 1989, 43, 990; d) L. Grehn, M. L. S. Almeida and U. Ragnarsson, Synthesis, 1988, 992.
- 110 P. K. Bondy and Z. N. Canallakis, J. Chromatogr., 1981, 224, 371.
- 111 S. Kim and H. Chang, Bull. Korean Chem. Soc., 1986, 7, 70.
- 112 R. K. Olsen, J. Org. Chem., 1970, 35, 1912.
- 113 P. J. Stang, M. Hanack and L. R. Subramanian, Synthesis, 1982, 85.
- 114 T. Kametany, T. Suzuki and K. Ogasawara, Chem. Pharm. Bull., 1972, 20, 1057.
- 115 a) M. Černý, J. Málek, M. Čapka and V. Chvalovský, Collect. Czech. Chem. Commun., 1969, 34, 1033; b) J. Vit, Technical Information Bulletin, Hexcel Corporation, Michigan, 1986.
- 116 H. E. Carter, R. L. Frank and H. W. Johnston, Org. Synth. Coll. Vol. III, 1955, 167.
- 117 a) A. J. Speziale and L. R. Smith, J. Org. Chem., 1962, 27, 3742; b) A. J. Speziale and R. L. Smith, J. Org. Chem., 1963, 28, 1805; c) A. J. Speziale, L. R. Smith and J. E. Fedder, J. Org. Chem., 1965, 30, 4306.
- 118 G. Zinner and R. Stoffel, Arch. Pharm. (Weinheim, Ger.), 1969, 302, 691.
- 119 R. Milcent, M. Guevrekian-Sohomoniantz and G. Barbier, J. Heterocycl. Chem., 1986, 23, 1845.
- 120 a) Y. Kiso, M. Inai, K. Kitagawa and T. Akita, Chem. Lett., 1983, 739; b) H. Waldmann and H. Kunz, Liebigs Ann. Chem., 1983, 1712; c) L. A. Carpino, Acc. Chem. Res., 1987, 20, 401.
- 121 R. Andruszkiewicz, H. Wojciechowska and E. Borowski, Pol. J. Chem., 1978, 52, 1167.
- 122 C. M. Svahn and J. Gyllander, J. Chromatogr., 1979, 170, 292.
- 123 S. Fuchs and W. Voelter, Z. Naturforsch., B, 1976, 31, 1410.
- 124 K.-H. Scholz, H.-G. Heine and W. Hartmann, Liebigs Ann.

Chem., 1976, 1319.

- 125 H. J. Veith, M. Hesse and H. Schmid, Helv. Chim. Acta, 1970, **53**, 1355.
- 126 G. Schill and J. Crommen, TrAc, Trends Anal. Chem. (Pers. Ed.), 1987, **6**, 111.

ABSTRACT

A new, simple and efficient preparative procedure of a potentially wide scope for the selective protection of mixed primary/secondary amines is presented. Its applicability is demonstrated on spermidine to make both protected and simple substituted derivatives.

The results obtained during this research are the main subject of this thesis. It also includes an introductory section on the biological aspects of polyamines and a review on the methods commonly used for their synthesis.

The new approach is based on tert-butoxycarbonylation of carbamate groups (exhaustive tert-butoxycarbonylation) derived from the primary amino functions only. In most cases, benzyl polycarbamates are used for this purpose. Subsequent removal of all benzyloxycarbonyl (Z) groups from the resulting intermediates by catalytic hydrogenolysis liberates the secondary amino functions, whereas tert-butoxycarbonyl (Boc) is retained on the primary ones. Alternatively, selective removal of Z only from amino functions, protected by both Z and Boc, which can be accomplished by base-catalysed methanolysis, results in protected polyamines with Boc and Z on their primary and secondary amino groups, respectively. The new reaction has been performed on spermidine to give N^1, N^8 -Boc₂-spermidine. By virtue of the non-equivalence of the two primary amino groups in this molecule, the synthesis of

N^8 -Boc- N^1 -Z-spermidine, starting instead with an easily available cyclic derivative (hexahydropyrimidine), is also presented. The yields of most intermediates as well as of the two products were high.

The synthesis of all three monoacetylated spermidines is also reported. The N^4 -acetyl derivative was obtained in four steps from spermidine via the triacetyl intermediate by selective deacetylation after exhaustive tert-butoxy-carbonylation as well as directly from N^1, N^8 -Boc₂-spermidine. The N^1 -acetyl and N^8 -acetyl derivatives were both obtained in four simple protection/deprotection steps from a common intermediate, N^8 -Boc- N^1 -Z-spermidine mentioned above, thus illustrating the versatility of this compound.

The synthesis of a few N-ethylspermidines is also described. The N^1, N^8 -diethylspermidine was obtained in three steps from spermidine via the Z₃-derivative by N^1, N^8 -diethylation followed by removal of the Z groups. The attempted syntheses of the N^1 -ethyl and N^8 -ethyl derivatives were performed by reduction of the amide groups of the corresponding acetyl spermidines.

New mixed alkyl benzyl imidodicarbonates were prepared by reaction of benzyloxycarbonyl isocyanate with appropriate alcohols. The compounds are of interest as potential Gabriel reagents. Completely selective removal of one of the alkoxycarbonyl groups from the N-atom of the imidodicarbonates was demonstrated in several instances, giving either benzyl or the alternative carbamate.

RESUMO

Poliaminas naturais (como por ex. espermidina e espermina) e seus derivados constituem um grupo de compostos com grande importância em muitos processos biológicos a nível celular.

Recentemente foi verificado que tais compostos são potenciais agentes antineoplásticos, sendo ainda úteis no tratamento de doenças parasitárias.


Em face da importância biológica referida, a síntese de poliaminas e seus derivados reveste-se de grande interesse. Na maioria dos casos, o êxito da síntese vai depender da modificação selectiva dos diferentes grupos amina. Em 1986, embora na literatura estivessem já descritos alguns métodos para a protecção selectiva dessas aminas, parecia não existir um método ideal, particularmente, no que se refere à protecção total desses compostos. A metodologia usada requeria um grande número de fases e/ou condições drásticas para a remoção dos grupos de protecção (por ex. grupos tosilo e ftaloilo), tornando-se por isso desejável um novo método alternativo mais simples.

Iniciou-se, então, na escola do Doutor Ulf Ragnarsson, Instituto de Bioquímica, Centro de Biomédicas, Universidade de Uppsala, Suécia, um projecto de investigação a fim de explorar um método simples e geral para a protecção selectiva de aminas primárias/secundárias. Deu-se particular atenção ao caso da espermidina, usando como reacção chave a N-tert-butoxi-

carbonilação de grupos uretanos catalisada por 4-dimetilamino-
piridina (DMAP). Para este fim, foi usado o grupo
benziloxicarbonilo (Z) o qual é ortogonalmente removido na
presença do grupo Boc e em condições suaves. É de referir que
antes de se usar a espermidina como substrato foram sempre
realizados estudos preliminares com um composto modelo mais
simples, a N-etil-1,2-etanodiamina.

No Esquema I está representado a nova metodologia para o
caso da espermidina. Depois de se introduzirem os três grupos
Z, efectuou-se a N-tert-butoxicarbonilação nos grupos NH
terminais. Em seguida, a protecção Z foi removida por dois
métodos alternativos:

- a) remoção total por hidrogenólise catalítica, dando origem
ao derivado N¹,N⁸-Boc₂-espermidina;
- b) remoção selectiva por metanólise catalisada por base,
para se obter o composto N⁴-Z-N¹,N⁸-Boc₂-espermidina.



Esquema I

Por outro lado, como a espermidina contém dois grupos -NH_2 não equivalentes, conseguiu-se a partir do derivado cíclico (hexa-hidropirimidina), resultante da reacção entre a triamina e formaldeído, sintetizar o composto chave, $\text{ZNH(CH}_2)_3\text{NH(CH}_2)_4\text{NHBoc}$.

Uma vez obtidos estes precursores protegidos, foram estudadas as suas potencialidades como reagentes chave na síntese de derivados da espermidina com interesse biológico. Assim, foram sintetizados os seguintes derivados monoacetilados os quais são importantes como metabólitos e produtos de excreção:

N^4 -acetilespermidina $[\text{NH}_2(\text{CH}_2)_3\text{NAC}(\text{CH}_2)_4\text{NH}_2]$;

N^1 -acetilespermidina $[\text{AcNH}(\text{CH}_2)_3\text{NH}(\text{CH}_2)_4\text{NH}_2]$;

N^8 -acetilespermidina $[\text{NH}_2(\text{CH}_2)_3\text{NH}(\text{CH}_2)_4\text{NHAc}]$.

A N^4 -Ac-espermidina foi obtida em quatro fases por conversão da espermidina no derivado triacetilado, seguindo-se a N-tert-butoxicarbonilação exaustiva e remoção selectiva dos grupos acetilo. Este derivado também foi obtido directamente a partir do precursor N^1, N^8 -Boc₂-espermidina.


Os derivados N^1 -Ac- e N^8 -Ac-espermidina foram obtidos em quatro fases simples de protecção/desprotecção a partir do composto chave N^1 -Z- N^8 -Boc-espermidina, ilustrando assim a versatilidade deste composto. No caso do derivado acetilado em N^1 , começou-se por proteger o grupo amina secundária com a protecção Boc. Em seguida removeu-se o grupo Z para se obter o intermediário $\text{NH}_2(\text{CH}_2)_3\text{NBoc}(\text{CH}_2)_4\text{NHBoc}$. Este derivado foi acetilado no grupo amina, tendo-se depois removido os grupos

Boc. No caso da N⁸-Ac-espermidina, o grupo amina secundária foi protegido com o grupo Z, seguindo-se a remoção selectiva da protecção Boc. De um modo semelhante, o derivado resultante, ZNH(CH₂)₃NZ(CH₂)₄NH₂, após acetilação seguida de remoção dos grupos Z, deu o composto acetilado em N⁸.

Foram também sintetizados os derivados etilados da espermidina, EtNH(CH₂)₃NH(CH₂)₄NH₂, EtNH(CH₂)₃NH(CH₂)₄NH₂ e NH₂(CH₂)₃NH(CH₂)₄NH₂. Estes compostos podem ser também importantes sob o ponto de vista biológico, nomeadamente em quimioterapia.

A N¹,N⁸-Et₂-espermidina foi obtida em dois passos por N¹,N⁸-dietilação do intermediário N¹,N⁴,N⁸-Z₃-espermidina seguida da remoção dos grupos Z. Os N¹-etil- e N⁸-etil-derivados da espermidina foram sintetizados por redução do grupo amida das correspondentes acetilespermidinas.

Foi possível, paralelamente ao trabalho mencionado, iniciar um estudo que permitisse a síntese total de poliaminas a partir de precursores adequadamente protegidos, tais como, Boc₂NH. Assim, começou-se por sintetizar novos reagentes de Gabriel, imidodicarbonatos de alquilo e benzilo, por reacção do isocianato de benziloxycarbonilo com alcoóis apropriados (Esquema II).



Esquema II

A completa remoção selectiva de um dos grupos alcóxicarbonilo do átomo de azoto destes imidodicarbonatos foi demonstrada com vários exemplos, obtendo-se o carbamato de benzilo ou o carbamato alternativo.

Em conclusão, os objectivos do projecto inicial foram amplamente alcançados uma vez que se conseguiu estabelecer um novo método, simples e eficiente, para a protecção selectiva de poliaminas o qual poderá ainda ser relevante no campo dos alcalóides. Por outro lado, os imidodicarbonatos preparados são reagentes promissores para uma estratégia envolvendo a síntese total de poliaminas, usando as reacções de Gabriel ou de Mitsunobu.

Os resultados obtidos durante este projecto de investigação constituem a parte essencial e original desta dissertação, que inclui uma breve introdução sobre os aspectos biológicos das poliaminas naturais e ainda uma revisão sobre os métodos mais usados na sua síntese.

Parte dos resultados obtidos nesta investigação deram já origem a quatro artigos publicados em colaboração ^{109a-c}.