

U. PORTO

FMUP FACULDADE DE MEDICINA
UNIVERSIDADE DO PORTO

MESTRADO INTEGRADO EM MEDICINA

2017/2018

João Vítor Barrau Mendes Ferreira Dinis

Qualidade do Sono e Depressão em Estudantes Universitários:

Uma Revisão Sistemática

Quality of Sleep and Depression in College Students:

A Systematic Review

março, 2018

FMUP



FACULDADE DE MEDICINA
UNIVERSIDADE DO PORTO

MESTRADO INTEGRADO EM MEDICINA

João Vitor Barrau Mendes Ferreira Dinis
**Qualidade do Sono e Depressão em Estudantes Universitários:
Uma Revisão Sistemática**
**Quality of Sleep and Depression in College Students:
A Systematic Review**

Mestrado Integrado em Medicina

Área: Ciências médicas e da saúde - Medicina clínica

Tipologia: Monografia

Trabalho efetuado sob a Orientação de:
Professor Doutor Miguel Ângelo Marques Bragança

Trabalho organizado de acordo com as normas da revista:
Sleep Science

março, 2018


FMUP

Eu, João Vitor Barrau Mendes Ferreira Dinis, abaixo assinado, nº mecanográfico 201203440, estudante do 6º ano do Ciclo de Estudos Integrado em Medicina, na Faculdade de Medicina da Universidade do Porto, declaro ter atuado com absoluta integridade na elaboração deste projeto de opção.

Neste sentido, confirmo que **NÃO** incorri em plágio (ato pelo qual um indivíduo, mesmo por omissão, assume a autoria de um determinado trabalho intelectual, ou partes dele). Mais declaro que todas as frases que retirei de trabalhos anteriores pertencentes a outros autores, foram referenciadas, ou redigidas com novas palavras, tendo colocado, neste caso, a citação da fonte bibliográfica.

Faculdade de Medicina da Universidade do Porto, 13/03/2018

Assinatura conforme cartão de identificação:

 _____

NOME

João Vitor Barrau Mendes Ferreira Dinis

NÚMERO DE ESTUDANTE

201203440

E-MAIL

joavitordinis@gmail.com

DESIGNAÇÃO DA ÁREA DO PROJECTO

Ciências médicas e da saúde > Medicina clínica

TÍTULO DISSERTAÇÃO/MONOGRAFIA (riscar o que não interessa)

Quality of Sleep and Depression in College Students: A Systematic Review

ORIENTADOR

Miguel Ângelo Marques Bragança

COORIENTADOR (se aplicável)

ASSINALE APENAS UMA DAS OPÇÕES:

É AUTORIZADA A REPRODUÇÃO INTEGRAL DESTA OBRA APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input type="checkbox"/>
É AUTORIZADA A REPRODUÇÃO PARCIAL DESTA OBRA (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input type="checkbox"/>
DE ACORDO COM A LEGISLAÇÃO EM VIGOR, (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) NÃO É PERMITIDA A REPRODUÇÃO DE QUALQUER PARTE DESTA OBRA.	<input checked="" type="checkbox"/>

Faculdade de Medicina da Universidade do Porto, 13/03/2018

Assinatura conforme cartão de identificação:

João Vitor Dinis

Dedico este trabalho a todos os que me ajudaram a chegar até aqui, porque o esforço de alguém sozinho não move a montanha.

Title: Quality of Sleep and Depression in College Students: A Systematic Review

Running Title: Quality of Sleep and Depression in College Students

Declaration of Funding

None of the authors had any direct or indirect funding in support of this study

Abstract

Nowadays, sleep-related problems are a prevalent occurrence among university students. Poor sleep quality is one of the most studied aspects of sleep complaints, affecting from 10% to 50% of this population. Poor sleep quality consequences are many and have a profound impact in the student's psychobiological health. University students live through a period of psychological challenge and adaptation, since the transition from high school to professional life. Abrupt autonomy challenges students to deal with many choices, from their academic and social life to their intimate habits. Frequently, sleep hygiene is neglected, or they are unable to use proper coping mechanisms, resulting in disturbing consequences that could impact their lives as adults. Research has found a significant association between sleep quality and depression or depressive symptoms, but this relationship is still somewhat difficult to interpret. The objective of this review is to appraise the current knowledge around the relationship of sleep with depression in this group of young adults. The methodology, the relationship between sleep and depression and possible influencing factors were examined. The current literature still supports a bidirectional relationship between sleep and depression, however, the importance of sleep quality is becoming the most relevant variable. Education and the application of policies regarding sleep hygiene may prevent, in some cases, the development of depression and improve the quality of sleep in other cases. Future research should clarify the relationship between sleep problems and depression in a way they could be prevented or, at least, minimized with effective and achievable interventions.

Key words: sleep quality; depression; college students;

1. Introduction

When young adults go through university, they experience a period of psychological challenge and adaptation, since the transition from high school to professional life. The sudden autonomy challenges students with varying choices, from their academic and social life to their intimate habits. Frequently, some important dimensions of their lives are sacrificed, being sleep hygiene often neglected, and in the presence of frustration or life changes, they are unable to use proper coping mechanisms, resulting in problematic consequences that could impact their lives through academic life and as adults.

The occurrence of poor mental health among university students varies, but nonetheless the rates of depressive symptoms can go as high as 50% in some countries.(1) Along with this situation, the deterioration of good sleep habits may be contributing to the poor mental health seen in this population. Sleep quality, a measurement that is related to sleep hygiene, has been shown to be poor in this population in different countries, affecting in some cases around 50% of the students.(1-3) An accumulation of research shows that the relationship between disturbed sleep and depression is complex and bidirectional. As this relation is better understood, a dedicated intervention both on sleep quality and depression could prevent the emerging problem in the affected student populations, as well as serving as prevention in the post-graduated young adults.

The aim of this systematic review is therefore to appraise the current available literature that specifically probes this relation, focusing mainly which of the variables of depression, depression symptoms or poor sleep quality may contribute more to the problem. Additionally, this review also aims to appreciate some of the factors that can potentially interfere in this relation.

2. Material and Methods

This review methodological approach has been based in PRISMA's guidelines, available at "prisma-statement.org/". Pubmed was queried for all references from 1970 through July 22, 2017 for studies related to depression associated to sleep in college students or young adults. The search terms were the following: ((Sleep[MeSH Terms]) AND depression[MeSH Terms]) AND adults, young[MeSH Terms], with 318 results; ((Sleep[MeSH Terms]) AND depression[MeSH Terms]) AND students[MeSH Terms] with 40 results; (("sleep"[MeSH Terms]) AND depression[MeSH Terms]) AND students) with 75 results; ((Sleep[Title/Abstract]) AND Depression[Title/Abstract]) AND Students[Title/Abstract]) with 272 results; In a first phase selection, repeated articles were sorted out, and the remaining articles were selected by title, having as inclusion criteria the combination of the terms "sleep", "sleep quality", "depression", "depressive symptoms", "college", "students", "young adults", "university", "population". In a second phase, the selected articles were individually read and submitted to the following criteria: 1) any type of article, excluding reviews, meta-analyses and editorial notes; 2) population attending college or young adults; 3) clear methodology and generally accepted data collection instruments, related to assess depression, depressive symptoms, sleep and sleep quality, coherent with the intervention; 4) the outcome would be associated with sleep, sleep quality, depression and depressive symptoms; 5) exclusion of any article in which the outcome would relate intentionally to insomnia or other sleep disturbances, other than affecting sleep quality.

The data retrieved from each article included the study type, the country, the population context, the methodology, the reported limitations, the outcomes regarding the instrument

used. Also, data was retrieved concerning other factors the authors consider having influence in the main outcomes in each study.

3.Results

From the initial search, 705 articles were selected. After the first phase selection, 66 were selected, which followed to be selected in the second phase selection, resulting in 32 articles which were included in the review (Figure 1).




Figure 1 – Flowchart of included articles

From all articles, 13 were from North America (11 – U.S.A; 1 Canada; 1 Mexico), 10 were from Asia (5- Japan; 3 China; 1 Thailand; 1 Nepal), 2 from Oceania (2- Australia), 2 from Europe (1- Austria; 1- Poland), 2 from Middle East (1- Turkey; 1- Lebanon), 2 from Africa (1- Nigeria; 1- Ethiopia) and 1 study regarding various geographic proveniences.

The study types included 22 cross-sectional studies, 7 longitudinal studies, 1 experimental study, 1 pilot study and 1 quasi-experimental study.

3.1 Population

A great portion of the studies included a general college student population (26), although some were specific college schools (Medicine – 2; Nursery – 1; Arts – 1) and two studies included both college students and young adults.

3.2 Methodology applied

Regarding sleep and sleep quality the most used instrument used in was the Pittsburgh Sleep Quality Score (PSQI),21 articles, followed by questionnaires made by the authors ,6 articles, and other questionnaires, 6 articles: Sleep Habits Questionnaire, Tokyo Metropolitan Institute for Neuroscience Life Habits Inventory, Auckland Sleep Questionnaire and sleep diaries. One article used the PSQI associated with actinography.

Regarding depression or depressive symptoms, the most used instrument was the Center for Epidemiologic Studies Depression Scale (CES-D), 13 articles, followed by the Depression, Anxiety and Stress Scale (DASS 21), 6 articles, then the Beck Depression Inventory (BDI-II), 5 articles, and other 5 articles used a variety of other scales: Hospital Anxiety and

Depression Scale, The Minnesota Multiphasic Personality Inventory, the Patient Depression Questionnaire, Self-Rating Depression Scale and the WHO-Five Well-being Index. Three articles used a combination of scales, CES-D with DASS 21 or the Hamilton Depression Rating Scale.

4. Discussion

The results in this systematic review reinforce the notion that there is a strong association between sleep quality, or sleep hygiene, and depression or depressive symptoms (Table 1). The literature is not consensual of the direction of association, and some authors hypothesize there is a bidirectional association between sleep and depression.(4-7) In terms of cross-sectional studies, there is a greater number of those which support that poor sleep quality predicts depression or depressive symptoms. The other longitudinal studies pointed to a greater risk of depression, or greater presence of depressive symptoms when sleep problems were present.(4, 5, 8-10) A quasi-experimental study comparing two different types of interventions for improving sleep quality showed reduction in depressive symptoms in both interventions.(11) One experimental study established a model in which poor sleep quality was associated with greater difficulty disengaging attention from negative stimuli, which in turn predicted upsurges in depressive symptoms.(12) There is also evidence that poor sleep quality is associated with greater struggle disengaging attention from negative stimuli, which will increase depressive symptoms. (12)

In the perspective of depression, students who are depressed are less likely to use adaptive coping strategies, therefore may experience diminished sleep quality or greater sleep start time variability.(10, 13) There is also evidence that as depression increases in severity, the odds ratio or risk of poor sleep quality also increase.(1, 14) Episodic and chronic depression are both linked to sleep quality. Despite episodic depressive symptoms being significant predictors of sleep complications, chronic depressive symptoms are particularly greater predictors.(4) Interestingly one of the longitudinal studies, which studied the transition from high-school to college, found that pre-transition depressive symptoms were associated with subsequent post-transition subjective and objective sleep problems, but not the reverse. However post-transition depressive symptoms were concurrently associated with greater sleep problems. (8) In an interesting opposite view, in the point of view from optimism, one longitudinal study found that optimism is a predictor of sleep quality. (15)

Regarding the studies' overall results, it's plausible to affirm that in the presence of a bidirectional relationship, self-reported sleep quality is being shown to be more consistent as a predictor of depression or depressive symptoms, despite depression itself also constitutes a predictor of sleep quality.

Table 1 – Interventions and characteristics of individual studies

Study	Design	Setting	Population	Interventions and Outcome measures	Results	Limitations
Moo-Estrella et al. (2004)	Cross-Sectional	University of Yucatan, Mexico	College Students from various schools	Questionnaires from events occurring the previous week; SHB and BDI-II	Students with depressive symptoms had more severe sleep alterations than those without symptoms. The proportion of subjects with depressive symptoms and who reported a nonrestorative sleep was five times higher than that of students with depressive symptoms who had good sleep quality (OR = 4.9, 2.03?12.13, p <0.0001)	Self-reported questionnaires; Cross-sectional study; selection bias
Carney et al. (2006)	Cross-Sectional	University in United States of America	College Students from various schools	Questionnaires after a 2-week Social Rhythm Metric evaluation. PSQI determined good and poor sleepers; PSQI and BDI-II	Results showed a significant group effect [F(12, 154) = 2.15, p < 0.017]. Follow-up ANOVAs revealed a statistically significant group effect for “Out of bed,” “Into bed.” An examination of the means revealed good sleepers were less variable than poor sleepers regarding their rise times, and their bed times. After controlling for depression, poor sleepers continued to have greater variability [F(4, 216) = 2.47, p < 0.046]; Lower rates of social engagement and regularity of social engagement seen in poor sleepers	Self-reported questionnaires; Cross-sectional study; the sample of participants was composed predominantly by female Caucasian subjects; the study focus only social zeitbergs

Study	Design	Setting	Population	Interventions and Outcome measures	Results	Limitations
Regestein et al. (2010)	Longitudinal	University in United States of America	Residents of a woman's college	A pilot study 6 week before end of the school year and a main study afterwards; CES-D, HAM-D3 and a Sleep Quality Score created by the authors including sleep debt, daytime sleepiness, rising time trouble sleeping; uses of alcohol, sedatives, tranquillizers, or anti-depressants	Diminished sleep apparently risked depressive symptoms. Top quartile Depressive Tendency scores. ($\chi^2 = 42.0$; $P=0.001$) associated with many sleep measures, including late bedtimes (OR=6.5; $P=0.01$);, greater sleep debt (OR=12.6; Wald $\chi^2 = 12.6$; $P=0.0004$);, greater sleepiness (OR=8.0; $P=0.005$; effect size=2.6) and more caffeine drinks daily (OR=5.1; $P=0.02$)	Self-reported sleep habits; the sample of participants was composed only by females
Augner et al. (2011)	Cross-Sectional	Three nurse's schools in Upper Austria and one technical school in Salzburg, Austria	Students from the mentioned colleges	Questionnaires from previous events; Self-created sleep quality scale (How often during the past 2 weeks did you feel drowsy or sleepy during the day? - 0 (never) to 4 (very often)); WHO-5	Subjective sleep quality is significantly associated with parameters of mental health (Depression score $r = -.57$; $p < 0.001$); Best predictors for poor subjective sleep quality were high depression score and long sleep onset latency; Results show significant increased risk of poor sleep experience for persons with high depression score (OR=3.90 [1,88-8,06] $p < 0.001$)	Self-reported questionnaires; Cross-sectional study; High female proportion sample

Study	Design	Setting	Population	Interventions and Outcome measures	Results	Limitations
Matsumoto et al. (2011)	Cross-sectional	Kurume University, Japan	Male students from the mentioned university	Questionnaires from previous events; TMIN-LHI, SDS and a adapted version of SF-36	Mental component of instruments subjects, poor sleep group das significantly poorer evaluation than the scores of any other sleep type. SDS scores in this group were >40 (p<0,001). as in SF-36 scores (p<001)	Self-reported questionnaires; Cross-sectional study; the sample of participants was small and composed only by males
Trockel et al. (2011)	Quasi-experimental	Private University in the United States of America	Students in the 2 first year residence halls	8-week cognitive behavioral therapy for insomnia via e-mail delivery; Questionnaires; PSQI and CES-D	Among students with lower PSQI scores at baseline, there were statistically significant differences between the intervention groups. Participation was associated with greater improvements in sleep quality and greater reductions in symptoms of depression among subjects with low sleep quality at baseline for the most successful intervention: PSQI 7.68 to 5.26 = -2.42 points; Cohen's d = 1.33 and; CES-D 19.69 to 13.75 = -5.94 points; Cohen's d = 0.57	Selection bias; Intervention group population had some degree of heterogeneity; Self-reported questionnaires

Study	Design	Setting	Population	Interventions and Outcome measures	Results	Limitations
Benitez et al. (2012)	Cross-Sectional	Midwestern University in United States of America	Students from the mentioned university	Questionnaires from previous events; PSQI and MMPI-2	Multivariate analysis of variance identified a significant difference between poor and good sleepers on the MMPI-2 RC, $F(9,57)=2.471$, $p<.019$, partial $\eta^2 =.281$. Post-hoc analyses with a Bonferroni correction identified differences on scales that were related to depression and anxiety, including: Demoralization, Low Positive Emotions, Dysfunctional Negative Emotions	Self-reported sleep habits; Cross-sectional study; modest sample size, composed mostly by females
Carskadon et al. (2012)	Longitudinal	Brown University, United States of America	First-year students from the mentioned university	Daily questionnaires and self-reported sleep diaries; DNA samples taken for genotyping; Sleep Diaries and CES-D	Total sleep time ($F 3,131 = 125.87$, $P < 0.001$) and CES-D ($F 3,131 = 69.48$, $P < 0.001$) showed significant group differences driven by the phenotype grouping parameter. A significant representation of the S'S' genotype for the 5-HTTLPR was seen for participants who reported shorter sleep and high CES-D scores relative to other groups (chi-square = 15.04; df = 6; $P = 0.02$), even after excluding clinically significant CES-D scored participants (chi-square = 12.90; df = 6; $P = 0.045$)	Small sample size; Self-reported; Unmeasured third variables, including the possibility of population stratification or linkage disequilibrium between measured and causal variants

Study	Design	Setting	Population	Interventions and Outcome measures	Results	Limitations
Lemma et al. (2012)	Cross-Sectional	Haramaya University and University of Gondar, Ethiopia	Undergraduate students from second year to final year	Subjects assessed by Bi-lingual questionnaires; PSQI and DASS 21	The level of perceived depression has shown significantly increasing odds of poor sleep quality across the quartiles as the level of depression increased from mild to extremely severe with the following odds ratio (95% CI) [1.36 (1.06, 1.75)], [1.64 (1.27, 2.11)], [1.64 (1.11, 2.42)], [2.65 (1.56, 4.49)] respectively	Self-reported sleep habits; Cross-sectional study
Adams et al. (2013)	Pilot Study	University in the United States of America	Junior and senior college students	Assessed by questionnaires and 7-day self-reported sleep diaries; PSQI and BDI-II	After controlling for age and gender, poor sleep quality significantly predicted depression $b=1.389$, $p<0.001$. Poor sleep quality, age, and gender explained 29.7 percent of variance in depression, $F(3,225)=31.256$, $p<0.001$. Sleep quality was a mediator of the relation between technology use and depression $b=1.526$, $p<0.001$	Self-report bias; Selection bias

Study	Design	Setting	Population	Interventions and Outcome measures	Results	Limitations
Kenney et al. (2013)	Cross-Sectional	West coast universities in the United States of America	Undergraduate students	Subjects assessed by questionnaires; PSQI and DASS 21	Poor mental health predicted poor sleep quality (Std. Coefficient= 0.41 p<0,001)	Self-reported bias; Cross-sectional study; Depression was integrated in mental health variable
Wong et al. (2013)	Longitudinal	Universities and colleges from Hong Cong and Macau	Undergraduate students	The study was conducted across three consecutive academic semesters; Subjects assessed by questionnaires; PSQI and DASS 21	Feeling of depression was predicted by daytime dysfunction (Standardized Regression Coefficient, β =.121, p =.024) even after we controlled for participants' depression at baseline (β =.621, p .001)	Self-report bias; Selection bias
Tavernier and Willoughby. (2014)	Longitudinal	University in Southern Ontario, Canada	First-year university students from the mentioned university	Two assessment times with questionnaires; Sleep routines questionnaire made by authors and CES-D	In terms of intrapersonal adjustment, subgroups classified by good sleep characteristics reported significantly better intrapersonal adjustment relative to subgroups characterized by poor sleep characteristics, among both morning-types and evening-types. $F(4, 775) = 3.645, p = 0.006$	Self-report bias; may not be generalizable; no bidirectional association addressed

Study	Design	Setting	Population	Interventions and Outcome measures	Results	Limitations
Lovell et al. (2014)	Cross-Sectional	University of the Sunshine, Australia	Students in the mentioned university	Subjects assessed by questionnaires; PSQI and DASS 21	Females and Males with depressive symptoms were more likely to have unhealthy sleep hours than were males or females without depressive symptoms; females (OR = 1.98, P = 0.007) males (OR = 4.24, P = 0.014)	Self-reported bias; Cross-sectional study; Time period bias; measures for health behaviors were brief
Matsushita et al. (2014)	Cross-Sectional	Preparatory schools for university entrance and 4 Japanese universities	Undergraduate students and "Ronin-sei" – young adults still trying to access university	Subjects assessed by questionnaires; Authors created sleep measurement questionnaire and used CES-D	Greater depressive symptoms if daily sleep duration <5 h (OR = 2.01, CI = 1.11–3.63, P = 0.022), 5to<6 h(OR = 1.43,CI = 1.05–1.95, P = 0.024),and 8 h or more (OR = 4.79, CI = 1.05–21.90, P = 0.043) vs. 6 to <7 h; Later bedtime was associated with increased depressive symptoms (OR = 1.47, 95% CI = 1.12–1.95, P = 0.006)	Self-report bias; Selection bias
Asaoka et al. (2014)	Cross-Sectional	Japan	Students, workers, and unemployed youth, aged 19–25 years	Web-based questionnaire survey; PSQI, CES-D and SF-8	New university graduates experienced ~1 h of sleep phase advancement which revealed that the interaction between current bedtime and that at one year before was significant for the scores of CESD [F (1, 113) = 5.52, P < 0.05], MCS [F(1, 113) = 10.09, P < 0.01], and PSQI [F (1, 113) = 4.10, P < 0.05]	Self-report bias; Recall bias; Cross-sectional study; no data about the weekend sleep phase

Study	Design	Setting	Population	Interventions and Outcome measures	Results	Limitations
Vanderlind et al. (2014)	Experimental	University of Texas, United States of America	Students in the mentioned university	Assessment with questionnaires; PSQI and CES-D; Actinography (Ambulatory Monitoring); participants also completed cognitive and affective measures both at baseline and after the 3-week period	One model ([df = 2] = 0.95, P = .62; RMSEA = .00, CFI = 1.00) reported Greater sleep difficulty and more sleep stability both significantly predicted greater difficulty disengaging attention (i.e., less cognitive control) from negative stimuli. indirect effects among associations in the initial model for which there was a possible intervening (or mediating) variable. For time 2 depressive symptoms, there was a marginally significant indirect effect for self-reported sleep difficulty (P = .07) and a significant indirect effect for sleep stability (P = .014). Cognitive control over negative stimuli was the intervening variable for these indirect effects	Self-reported bias; Sample size; Selection bias; study used correlational analyses and several of the associations tested in the path model were cross-sectional

Study	Design	Setting	Population	Interventions and Outcome measures	Results	Limitations
Wilson et al. (2014)	Cross-Sectional	Private women's liberal arts college, United States of America	Students in the mentioned college	Subjects assessed by questionnaires via internet survey; Authors created sleep measurement questionnaire and used CES-D and DASS-21	Poor sleep quality was significantly associated with both CES-D and DASS-21 measures of depression after adjusting for confounders. Students who reported poor/extremely poor sleep quality were more likely to also report prevalent CES-D scores ≥ 16 (AOR 2.8, 95% CI 1.3–5.8). Similarly, the odds of prevalent DASS-21 depression was 2.8 (95% CI 1.4–5.8) times that of students who self-reported sleep to be excellent, good, or okay	Self-reported bias; Cross-sectional study; the sample of participants was composed only by females
Demirci et al. (2015)	Cross-Sectional	Süleyman Demirel University, Turkey	Students in the mentioned university	Subjects assessed by questionnaires; PSQI, BDI-II and Smartphone Addiction Scale	Regression analyses indicated that higher levels of smartphone use and poor sleep quality predicted depression ($\beta = 0.226$, $t = 4.131$, $p < 0.01$; $\beta = 0.448$, $t = 8.173$, $p < 0.01$; respectively) In addition, depression predicted poor sleep quality. The findings suggest that depression acted as a mediator between smartphone overuse and sleep quality	Self-report bias; Selection bias; small sample size

Study	Design	Setting	Population	Interventions and Outcome measures	Results	Limitations
Doane et al. (2015)	Longitudinal	High school and a large southwestern university, United States of America	Students in the senior year in high school which then attended the mentioned university	At T1, participants completed a packet of self-report questionnaires (PSQI, CES-D and DASS 21) and portable Actinography. The same at T2 and T3. The average time between T1 and T2 assessments was 5.2 months (SD = .96), and the average time between T2 and T3 was 4.1 months	The direct paths examining depressive symptoms and sleep latency were significant across time. There was only one significant prospective effect, with depressive symptoms at T1 predicting increased sleep latency at T2 ($\beta = .24, p < .05$). The concurrent, and positive, association between depressive symptoms and latency at T3 was also significant ($\beta = .26, p < .05$), but no other concurrent correlations were significant. One crossover effect was significant with depressive symptoms at T1 predicting increased sleep start variability at T2 ($\beta = .25, P < .05$). The concurrent relationship at T2 was significant ($\beta = .28, p = .01$), with higher depressive symptoms being significantly associated within increased sleep start variability. There was one concurrent, positive, correlation between depressive symptoms and sleep efficiency at T2 ($\beta = .24, p < .05$) There was one predictive effect, such that depressive symptoms at T1 predicted increased subject sleep complaints at T2 ($b = .26, p \backslash .01$). There were also positive and significant concurrent association between depressive symptoms and subjective sleep problems at all three-time points (all p 's < .01)	Self-reported bias; small sample size

Study	Design	Setting	Population	Interventions and Outcome measures	Results	Limitations
Kono et al. (2015)	Cross-Sectional	Hokkaido University, Japan	International students in the mentioned college	Subjects assessed by questionnaires via internet survey; Authors created sleep measurement questionnaire and also used CES-D	The unadjusted odds ratios suggested that 12 variables had statistically significant associations with depressive symptoms: quality of sleep (adjusted OR 7.35; 95 % CI 3.87–14.0)	Self-reported bias; Cross-sectional study;
Peltzer and Pengpid (2015)	Cross-Sectional	Universities in different countries	Undergraduate students in the mentioned universities	Subjects assessed by questionnaires; Authors created sleep measurement questionnaire and used CES-D	Depression symptoms (moderate/severe) was associated with nocturnal sleep problems AOR (95% CI) 2.61 (2.29–2.99) p<0.001	Self-report bias; Cross-sectional study; recall bias; only those students reporting severe or extreme sleeping problems were included,
Fatima et al. (2016)	Cross-Sectional	Mater–University of Queensland Study of Pregnancy, Australia	Offspring with 21 years old	Subjects assessed by questionnaires; PSQI and CES-D	depression had a impact on poor sleep quality in males (OR 1.15; 95% CI 1.12-1.18) and females (OR 1.11; 95% CI 1.08-1.13)	Self-report bias; Cross-sectional study; Shortened version of PSQI used

Study	Design	Setting	Population	Interventions and Outcome measures	Results	Limitations
Kabrita and Hajjar-Muça. (2016)	Cross-Sectional	Six private and public universities, Lebanon	Undergraduate students in the mentioned universities	Subjects assessed by questionnaires; PSQI and CES-D	In females, wakeup time on weekdays, but not weekends, was negatively correlated with CES-D score ($r=-0.168$, $P<0.05$), as was the average weekday sleep duration ($r=-0.221$, $p<0.01$). No significant correlation between these sleep parameters and CES-D was observed in males.	Self-reported bias; Cross-sectional study
Pensuksan et al. (2016)	Cross-Sectional	One autonomous university, Thailand	Undergraduate students in the mentioned university	Subjects assessed by questionnaires; PSQI and DASS-21	Poor sleep quality was statistically significantly associated with symptoms of depression ($R=0.34$; $p < 0.001$)	Self-reported bias; Cross-sectional study
Supartini et al. (2016)	Cross-Sectional	Kyushu University, Japan	Students in the mentioned university	Subjects assessed by questionnaires; PSQI and CES-D	Depressive symptoms were significantly associated with bedtime ($p = 0.01$), sleep-onset latency ($p < 0.001$), and poor sleep quality ($p < 0.001$). No statistically significant association was found between depressive symptoms and either wake time or sleep duration; multivariate-adjusted OR of depression for bedtime of later than 01:30 versus 23:30 or earlier was 1.59 (95% CI: 1.04–2.44) Sleep-onset latency of more than 30 minutes was significantly associated with an increased prevalence of depressive symptoms (OR = 1.53; 95% CI: 1.22–1.93). Poor sleep quality was significantly associated with an increased prevalence of depressive symptoms in age and sex adjusted model (OR = 2.49; 95% CI: 2.02–3.06)	Self-report bias; Cross-sectional study; Shortened version of PSQI used

Study	Design	Setting	Population	Interventions and Outcome measures	Results	Limitations
Bhandari et al. (2015)	Cross-Sectional	Undergraduate campuses in Kathmandu and Chitwan, Nepal	Undergraduate students in the mentioned campuses	Subjects assessed by questionnaires; PSQI and PHQ-9	Mediation of association between sleep quality and depressive symptoms by internet addiction was statistically significant: 16.5% of the indirect effect of sleep quality on depressive symptoms	Self-reported bias; Cross-sectional study
Lau et al. (2015)	Longitudinal	University of Hong Kong, China	Undergraduate students in the mentioned university	Subjects assessed by questionnaires in 3 different period times; PSQI and DASS-21	There was no significant relationship between poor sleep quality at Wave 1 and any of the three mood variables at Wave 2 for the morning-type group. In both the evening-type and intermediate-type groups, on the contrary, poor sleep quality at Wave 1 positively predicted higher levels of depressive mood, at Wave 2 ($\beta = .424, p < .001 / \beta = .260, p < .001$)	Self-report bias; Adaptation bias in instruments

Study	Design	Setting	Population	Interventions and Outcome measures	Results	Limitations
Mokros et al. (2017)	Cross-Sectional	Faculty of Medicine at Medical University of Lodz, Poland	Students from the mentioned university	Subjects assessed by questionnaires; PSQI and BDI-II	Sleep quality predicted depressive symptoms independently of the investigated personal dispositions among students [GML model – BDI/PSQI: Step 1: $R^2 = 0.151$, $df = 3$, $F = 9.098$, $p < 0.001$ – $\beta = 0.393$ (0.237-0.549) $p < 0,001$; Step 2: $R^2 = 0.260$, $df = 5$, $F = 10.636$, $p < 0.001$ - $\beta = 0.396$ (0.245-0.547) $p < 0,001$; Step 3: $R^2 = 0.275$, $df = 6$, $F = 9.519$, $p < 0.001$ - $\beta = 0.380$ (0.228-0.531) $p < 0,001$]	Self-reported bias; Cross-sectional study; Small sample size; selection bias
Seun-Fadipe. (2017)	Cross-Sectional	Obafemi Awolowo University, Nigeria	Students from the mentioned university	Subjects assessed by questionnaires; PSQI and HADS	Depression has significant association with poor sleep quality ($p < .001$). Pearson correlation analysis showed a positive correlation between global sleep quality score and depression score on HADS ($r = 0.387$, $p < .001$)	Self-reported bias; Cross-sectional study; Recall bias
Tao et al. (2017)	Cross-Sectional	College in Anhui, China	Students in the mentioned college	Subjects assessed by questionnaires; PSQI, CES-D	Problematic mobile phone use and sleep and sleep quality are independently associated with mental health symptoms. Poor sleep quality was positively correlated with depressive symptoms (OR: 4.97, 95% CI: 3.99–6.19)	Self-reported bias; Cross-sectional study

Study	Design	Setting	Population	Interventions and Outcome measures	Results	Limitations
Wallace et al. (2017)	Longitudinal	Three colleges in Minnesota, United States of America	Students from the mentioned colleges	Subjects assessed by questionnaires; PSQI and CES-D (10 question version)	Analysis examining sleep deprivation, the final model shows that age ($b = -0.02$, $p < .05$), race ($b = 0.22$, $p < .05$), episodic and chronic depressive symptoms ($b L1 = 0.47$, $p < .0001$; $b L2 = 0.81$, $p < .0001$), and sleep quantity ($b L1 = -0.21$, $p < .0001$; $b L2 = -0.24$, $p < .0001$) were all significant predictors. Both the L1 and L2 effects for depressive symptoms were significantly predictive of sleep deprivation, suggesting that both episodic and chronic depressive symptoms were uniquely associated with sleep deprivation, although chronic depressive symptoms exhibited the larger effect. Fewer hours of sleep than usual (L1) and fewer hours of sleep across the 2 years (L2) were both associated with increased sleep deprivation.	Self-reported bias; Cross-sectional study; Small sample size; selection bias

SHQ Sleeps Habits Questionnaire; BDI-II Beck Depression Inventory; OR Odds Ratio; SD Standard Deviation; PSQI Pittsburgh Sleep Quality Score; CES-D Center for Epidemiologic Studies Depression Scale ; HAM-D3 Hamilton Depression Rating Scale; WHO-5 Self-Rating Depression Scale and the WHO-Five Well-being Index; TMIN-LHI Tokyo Metropolitan Institute for Neuroscience life habits inventory; SDS Self-Rating Depression Score; SF-36 Social Functioning 36; MMPI-2 Minnesota Multiphasic Personality Inventory; DASS-21 Depression, Anxiety and Stress Scale (DASS21); PHQ-9 Patient Health Questionnaire; HADS Hospital Anxiety and Depression Scale

4.1 Factors influencing sleep quality and depression

4.1.1 Sex

Females have been found to have higher risk of poor sleep quality, which may be associated by the gender-based differences in the biology of sleep (16) or contact to socio-economic pressures and cultural norms, reaction, and coping mechanisms to life stresses, in addition to biological factors.(17) Interestingly one study revealed that poor sleep quality is related to depressive symptoms in males, but stress in females.(13) Existing studies do not deliver information on whether gender difference remains significant after concurrently considering the impact of other socio-demographic, lifestyle factors, and affective disorders. The gender difference in sleep problems is mostly attributed to the physiologic differences, the importance of affective disorders and socioeconomic inequalities, indicating these may be the pathway variables through which gender disparity in poor sleep is displayed.(16)

4.1.2 Genetic Factors

A longitudinal study found that individuals carrying 2 alleles with the low-polymorphism of the serotonin transporter gene 5-HTTLPR reported more depressive mood in the presence of a persistent pattern of short nocturnal sleep. Thus, exposure to chronic levels of insufficient or disrupted sleep may manifest a preexisting vulnerability in genetically susceptible individuals.(5) Other longitudinal study found that the number of G alleles in rs11932595, from the CLOCK gene, is positively associated with self-reported sleep quality, suggesting that polymorphisms within this gene may be considered a factor.(12)

4.1.3 Year of study and College

Some authors found that increasing year of study in university or college predicted good sleep quality. It indicates that students in later years may have developed better coping strategies for their academic life.(1, 18, 19) There is also evidence that poor sleep quality and depression is common among medical students. (20) Thus the demands over specific students in different schools may also contribute to influence the first.

4.1.4 Use of technology

With the recent technological advancement, a new factor may have appeared as a potential influencer in the relationship between sleep and depression. This is evident when studies prior to the technological boom of electronic devices didn't account this variable.(21, 22) Some authors found that waking up to answer one's phone predicted lower sleep quality, which in turn predicted depression.(23) In another study, depression acted as a mediator between smartphone overuse and poor sleep quality. The mechanisms behind this phenomenon may be due to probably affecting sleep architecture, alteration of melatonin release by influence of bright light, displacement of sleep, influence of electromagnetic fields emitted from the devices on brain activity, physical discomfort during its usage or because of cognitive, emotional or physiological arousal. (24)

4.1.5 Chronotype

As individuals may be classified as either morning-types, intermediate types or evening-types, some studies report that circadian preferences may have an important role in the connection between sleep and depression. (10, 15, 20) The literature reveals mixed results concerning this factor. Several studies settle that an evening chronotype is linked to poor sleep

quality among young adults and college students. (20) Some authors support that the prevalence of depressive symptoms was significantly higher among students with a late bedtime and prolonged sleep-onset latency. (18, 25) One longitudinal study suggests that morning-type chronotype has a protective effect against the potential harm of poor sleep on optimism, yet not against the damage of pessimism on sleep quality(15). However, one study found that 40% of evening-types were classified as having good sleep characteristics and did not generally differ on the sleep characteristics relative to most of morning-types, who reported good sleep. The authors propose that perceived morningness–eveningness may not account to influence sleep quality. (10)

Regarding weekend, one study suggests that students tend to maintain similar sleep wake patterns across the week and weekend. (10) However, other study reported that students sleep significantly longer in weekends, mainly females. (17)

4.1.6 Sleep Quantity

The mechanisms underlying the association between bedtime and depression remain unknown as some authors haven't found a consistent relation between sleep duration and depression.(18) However, some authors support that short sleep duration is associated with increased depressive symptoms in a U-shaped relationship.(25) Despite these contradictory results, some authors found that there are significant improvements in several sleep patterns across the transition from high-school to college: increases in sleep efficiency, sleep minutes, and subjective sleep quality after this transition.(26)

4.2 Studies Limitations

Regarding the cross-sectional studies, the studies were limited mainly by their nature. In this case, causality is impossible to be established.(7) As the questionnaires used were mostly self-reported, being a subjective measure, recall bias must be accounted. Confirmation of these results using objective measurements, such as actigraphy monitoring throughout the day, would be desirable.(8, 25)

In the case of longitudinal studies which used actigraphy, sample size is one of the most significant limitations. The use of this instrument is yet difficult and expensive.(5)

The students who were able to participate in these studies were expected to maintain a healthy lifestyle and relatively good mental health. Consequently, the students who could not participate in the studies might be likely to have more severe problem and become socially withdrawn with problems such as sickness and absenteeism in the future. Such high-risk students should have been included in the studies.(25)

Other compensatory sleep habits haven't been accounted for, such as napping or nodding in the classroom.(25) Also, the weekend period was only considered in two of the studies (10, 17), which may have an important role in the characteristics in the subject's sleep phase(19)

The presence of mental health problems and poor sleep quality was assessed by standardized clinical questionnaires. However, these measures are not equivalent to clinical diagnoses(27), and these instruments have their own limitations. For example, PSQI may underestimate the outcomes of sleep quality(16) and both CES-D and DASS-21 instruments measure short- term depressive mood, therefore the outcome may not reproduce long-term associations that would result in clinical depression.(6)

5. Limitations

This systematic review has its own limitations. First, it only used one search engine and only searched for studies written in English. It is possible that a great quantity of relevant articles has been missed out and its results could bring a different perspective. Second, the focus on depression and sleep is not totally realistic, since depression and sleep quality are influenced by many other variants such as stress, anxiety or physical health, which in term may have an important role in their relationship (27-30). This indicates that the relationship between sleep and depression is a complex phenomenon and its approach by simplifying the variables may not represent the reality of the problem. Third, this review focused in a very specific population with habits and pressures that may not be seen in the general population, thus the variables may be influenced by other mechanisms not applied to college students. For example, university students have later sleep and wake times, higher rates of daytime sleepiness, and physical and mental health complaints, than the general population.(30) Fourth, the impact of socioeconomic and cultural factors was not deeply explored in this review as it includes studies from developed and under development countries.

6. Conclusion

It is becoming consistent that self-reported sleep quality is a predictor of depression or depressive symptoms, at least regarding the young adults that attend universities and colleges. However, it is important to acknowledge that sleep disturbances and depression have a bidirectional association, so depression also influences sleep, as a recent meta-analysis on older adults found.(31) Defended by many of the authors, and demonstrated by an experimental study, an intervention in sleep hygiene can have benefic effects on reducing depression among the students. The institutions should recognize the importance of sleep hygiene and educate its population of its consequences, having in consideration that this population is probably unaware of this problem and it's late night culture. As the understanding of the complexity of sleep is becoming clearer, also pragmatic interventions should result from its study. In this direction, some research has already been conducted, demonstrating that cognitive behavioral self-help programs can improve sleep and depressive symptoms.(11) The research on this field still has its challenges concerning the measurements. However, with the uprising in the miniaturization of technology, the actinography may become cheaper and easy to use and so giving the possibility of a profound study, among others, of the sleep habits in the human populations. Finally, there is a lack of evidence in how the different grades of depression affect sleep quality and vice versa. This could possibly demonstrate if sleep affects the progression of depression's severity, if there is a correlation between the worsening of the quality of sleep and depression stage, or how different types of depression relate to sleep.

7. Funding

None of the authors had any direct or indirect funding in support of this study.

8. Conflict of interest

The authors have no financial relationships, or conflicts of interest to disclose

9. Bibliography

1. Lemma S, Gelaye B, Berhane Y, Worku A, Williams MA. Sleep quality and its psychological correlates among university students in Ethiopia: a cross-sectional study. *BMC Psychiatry*. 2012;12:237.
2. Seun-Fadipe CT, Mosaku KS. Sleep quality and psychological distress among undergraduate students of a Nigerian university. *Sleep Health*. 2017;3(3):190-4.
3. Kenney SR, Lac A, Labrie JW, Hummer JF, Pham A. Mental health, sleep quality, drinking motives, and alcohol-related consequences: a path-analytic model. *J Stud Alcohol Drugs*. 2013;74(6):841-51.
4. Wallace DD, Boynton MH, Lytle LA. Multilevel analysis exploring the links between stress, depression, and sleep problems among two-year college students. *J Am Coll Health*. 2017;65(3):187-96.
5. Carskadon MA, Sharkey KM, Knopik VS, McGeary JE. Short sleep as an environmental exposure: a preliminary study associating 5-HTTLPR genotype to self-reported sleep duration and depressed mood in first-year university students. *Sleep*. 2012;35(6):791-6.
6. Wilson KT, Bohnert AE, Ambrose A, Davis DY, Jones DM, Magee MJ. Social, behavioral, and sleep characteristics associated with depression symptoms among undergraduate students at a women's college: a cross-sectional depression survey, 2012. *BMC Womens Health*. 2014;14:8.
7. Tao S, Wu X, Zhang Y, Zhang S, Tong S, Tao F. Effects of Sleep Quality on the Association between Problematic Mobile Phone Use and Mental Health Symptoms in Chinese College Students. *Int J Environ Res Public Health*. 2017;14(2).
8. Regestein Q, Natarajan V, Pavlova M, Kawasaki S, Gleason R, Koff E. Sleep debt and depression in female college students. *Psychiatry Res*. 2010;176(1):34-9.
9. Wong ML, Lau EY, Wan JH, Cheung SF, Hui CH, Mok DS. The interplay between sleep and mood in predicting academic functioning, physical health and psychological health: a longitudinal study. *J Psychosom Res*. 2013;74(4):271-7.
10. Tavernier R, Willoughby T. Are all evening-types doomed? Latent class analyses of perceived morningness-eveningness, sleep and psychosocial functioning among emerging adults. *Chronobiol Int*. 2014;31(2):232-42.
11. Trockel M, Manber R, Chang V, Thurston A, Taylor CB. An e-mail delivered CBT for sleep-health program for college students: effects on sleep quality and depression symptoms. *J Clin Sleep Med*. 2011;7(3):276-81.
12. Vanderlind WM, Beevers CG, Sherman SM, Trujillo LT, McGeary JE, Matthews MD, et al. Sleep and sadness: exploring the relation among sleep, cognitive control, and depressive symptoms in young adults. *Sleep Med*. 2014;15(1):144-9.
13. Lovell GP, Nash K, Sharman R, Lane BR. A cross-sectional investigation of depressive, anxiety, and stress symptoms and health-behavior participation in Australian university students. *Nurs Health Sci*. 2014.
14. Augner C. Associations of subjective sleep quality with depression score, anxiety, physical symptoms and sleep onset latency in students. *Cent Eur J Public Health*. 2011;19(2):115-7.
15. Lau EY, Hui CH, Lam J, Cheung SF. Sleep and optimism: A longitudinal study of bidirectional causal relationship and its mediating and moderating variables in a Chinese student sample. *Chronobiol Int*. 2017;34(3):360-72.
16. Fatima Y, Doi SA, Najman JM, Mamun AA. Exploring Gender Difference in Sleep Quality of Young Adults: Findings from a Large Population Study. *Clin Med Res*. 2016;14(3-4):138-44.
17. Kabrita CS, Hajjar-Muca TA. Sex-specific sleep patterns among university students in Lebanon: impact on depression and academic performance. *Nat Sci Sleep*. 2016;8:189-96.
18. Supartini A, Honda T, Basri NA, Haeuchi Y, Chen S, Ichimiya A, et al. The Impact of Sleep Timing, Sleep Duration, and Sleep Quality on Depressive Symptoms and Suicidal Ideation amongst Japanese Freshmen: The EQU SITE Study. *Sleep Disord*. 2016;2016:8737654.

19. Asaoka S, Komada Y, Aritake S, Morita Y, Fukuda K, Inoue Y. Effect of delayed sleep phase during university life on the daytime functioning in work life after graduation. *Sleep Med.* 2014;15(9):1155-8.
20. Mokros L, Witusik A, Michalska J, Lezak W, Panek M, Nowakowska-Domagala K, et al. Sleep quality, chronotype, temperament and bipolar features as predictors of depressive symptoms among medical students. *Chronobiol Int.* 2017;34(6):708-20.
21. Taub JM. Behavioral and psychophysiological correlates of irregularity in chronic sleep routines. *Biol Psychol.* 1978;7(1-2):37-53.
22. Pilcher JJ, Ginter DR, Sadowsky B. Sleep quality versus sleep quantity: relationships between sleep and measures of health, well-being and sleepiness in college students. *J Psychosom Res.* 1997;42(6):583-96.
23. Adams SK, Kisler TS. Sleep quality as a mediator between technology-related sleep quality, depression, and anxiety. *Cyberpsychol Behav Soc Netw.* 2013;16(1):25-30.
24. Demirci K, Akgonul M, Akpınar A. Relationship of smartphone use severity with sleep quality, depression, and anxiety in university students. *J Behav Addict.* 2015;4(2):85-92.
25. Matsushita M, Koyama A, Ushijima H, Mikami A, Katsumata Y, Kikuchi Y, et al. Sleep duration and its association with sleepiness and depression in "ronin-sei" preparatory school students. *Asian J Psychiatr.* 2014;9:61-6.
26. Doane LD, Gress-Smith JL, Breitenstein RS. Multi-method assessments of sleep over the transition to college and the associations with depression and anxiety symptoms. *J Youth Adolesc.* 2015;44(2):389-404.
27. Feng Q, Zhang QL, Du Y, Ye YL, He QQ. Associations of physical activity, screen time with depression, anxiety and sleep quality among Chinese college freshmen. *PLoS One.* 2014;9(6):e100914.
28. Wu X, Tao S, Zhang Y, Zhang S, Tao F. Low physical activity and high screen time can increase the risks of mental health problems and poor sleep quality among Chinese college students. *PLoS One.* 2015;10(3):e0119607.
29. Matsumoto Y, Toyomasu K, Uchimura N. Assessment of physical and mental health in male university students with varying sleep habits. *Kurume Med J.* 2011;58(4):105-15.
30. Samaranayake CB, Arroll B, Fernando AT, 3rd. Sleep disorders, depression, anxiety and satisfaction with life among young adults: a survey of university students in Auckland, New Zealand. *N Z Med J.* 2014;127(1399):13-22.
31. Bao YP, Han Y, Ma J, Wang RJ, Shi L, Wang TY, et al. Cooccurrence and bidirectional prediction of sleep disturbances and depression in older adults: Meta-analysis and systematic review. *Neurosci Biobehav Rev.* 2017;75:257-73.

Attachments

Information retrieved from: <http://www.sleepscience.org.br/instructions-for-authors>

Sleep Science

Instructions for Authors

SCOPE AND POLICY

The SLEEP SCIENCE journal (ISSN 1984-0659 print version) is the official organization of Associação Brasileira de Sono (ABS – Brazilian Association of Sleep) and Federação Latino-Americana de Sociedades de Sono (FLASS – Latin American Federation of Sleep Societies) for publication of scientific papers concerning sleep, chronobiology, and related topics. After being approved by the Editorial Board, all articles will be evaluated by two or more qualified Reviewers, in a blind process. Articles that fail to present merit, have significant errors in methodology or are not in accordance with the editorial policy of the journal will be directly rejected by the Editorial Board, with no recourse. Manuscripts that have not been published elsewhere except in abstract form, on any aspect of sleep and chronobiology and both in basic and clinical areas will be considered. The accuracy of all concepts presented in the manuscript is the exclusive responsibility of the authors. The journal reserves the right to make stylistic, grammatical and other alterations to the manuscript. Manuscripts must not be concurrently submitted to any other publication, print or electronic. Articles may be written in English (American or British usage is accepted, but not a mixture of these) and should state that the protocol has been approved by the Ethics Committee of the Institution where the research was carried out. All studies involving human subjects should inform that written consent has been obtained from all subjects (individually).

SUBMISSION CHECKLIST

For each manuscript submitted to Sleep Science, the authors should assure that a) the content of the manuscript has not been submitted for publication in another journal or book (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint) in English or in any other language, including electronically without the written consent of the copyright-holder; b) each potential conflict of interests of the authors are fully disclosed; c) the experimental procedures were properly approved by the ethics committee of the institution where the research was carried out, if the investigation involves experiments on humans or animals; d) every possible funding source was disclosed; e) all participants provided signed consent forms, in the case of medical research on humans; f) the copyright of the manuscript after its acceptance will be transferred to Sleep Science. To verify originality, your article may be checked by the originality detection service CrossCheck.

It is requested that the authors strictly follow the editorial guidelines of the journal. Failure to comply with the author instructions will result in the manuscript being returned to the authors. Abbreviations should be used sparingly and should be limited only to those that are widely accepted. All abbreviations should be defined at first use.

Sleep Science will accept clinical trials for publication only if they have received an identification number from the Clinical Trial Registry, or from a similar clinical trial database, validated by the criteria established by World Health Organization (WHO) and International Committee of Medical Journals Editors (ICMJE). For more information: <http://clinicaltrials.gov>.

CHANGES TO AUTHORSHIP

Authors are expected to consider carefully the list and order of authors before submitting their manuscript and provide the definitive list of authors at the time of the original submission. Any addition, deletion or rearrangement of author names in the authorship list should be made only before the manuscript has been accepted and only if approved by the journal Editor. To request such a change, the Editor must receive the following from the corresponding author: (a) the reason for the change in author list and (b) written confirmation (e-mail, letter) from all authors that they agree with the addition, removal or rearrangement. In the case of addition or removal of authors, this includes confirmation from the author being added or removed. Only in exceptional circumstances will the Editor consider the addition, deletion or rearrangement of authors after the manuscript has been accepted. While the Editor considers the request, publication of the manuscript will be suspended. If the manuscript has already been published in an online issue, any requests approved by the Editor will result in a corrigendum.

TYPES OF PAPER

This journal publishes contributions in the following categories:

Original Articles: This manuscript should be the result of original investigation, presenting unpublished data. Each manuscript should clearly state its objective or hypothesis; the design and methods used; the study setting and time period; the essential features of any interventions; the main outcome measures; the main results of the study; and a section placing the results in the context of published literature.

Short Communication: A Short Communication is related to concise reports on a single issue, preliminary findings on a given subject, or any other case in which the findings are not sufficient for a full original article. As in Original Articles, should be the result of original investigation, presenting unpublished data. The scope of this section is intended to be wide and to encompass methodology and experimental data on subjects of interest to the readers of the journal.

Review article: A Review article should provide a broad and critical analysis of a relevant area and should not be merely a chronological description of the literature or a data compilation. Illustrations summarizing the main concepts of the Reviews are highly recommended.

Overview: As a Review, An Overview does not contain unpublished data. It presents the point of view of the author(s) in a less rigorous form than in regular Reviews and is of interest to the general reader.

Case report: A case report should have at least one of the following characteristics to be published in Sleep Science: a case of special interest to the clinical research community; a rare case that is particularly useful to demonstrate a mechanism or a difficulty in diagnosis; new diagnostic method; new or modified treatment. If available, the authors could present a series of cases, rather than one individual case.

Theoretical essays: This is a very broad category of articles. Theoretical essays encompass several kinds of non-experimental articles, such as perspectives, hypotheses, and opinions. In these articles, the goal is to clearly report authors' point of view regarding a specific field of research clinical area.

Correspondences: Correspondences are intended to promote scientific debate. This specific and brief kind article will be an opportunity to comment, criticize, or highlight papers that were already published in Sleep Science.

	Maximum length*	Mandatory sections	Abstract	Keywords	Figures and tables	References
Original articles	Up to 6,000 words	Abstract; Introduction; Material and Methods; Results; Discussion; References	Maximum of 250 words, structured as Objective, Methods, Results, Discussion	Three to six	Limited to 8	Limited to 60
Short communications	Up to 3,000 words	Abstract; Introduction; Material and Methods; Results; Discussion; References	Maximum of 150 words, structured as Objective, Methods, Results, Discussion	Three to six	Limited to 4	Limited to 20
Reviews	Up to 8,000 words	Abstract; References	Maximum of 250 words; unstructured	Three to six	Limited to 8	Unlimited
Overviews	Up to 5,000 words	Abstract; References	Maximum of 250 words; unstructured	Three to six	Limited to 4	Limited to 40
Case reports	Up to 1,500 words	Abstract; References	Maximum of 150 words; unstructured	Three to six	Limited to 4	Limited to 20
Theoretical essays	Up to 4,000 words	Abstract; References	Maximum of 250 words; unstructured	Three to six	Limited to 4	Limited to 40
Correspondences	Up to 500 words	Not applicable	Not applicable	Not applicable	Not allowed	Limited to 10

MANUSCRIPT SUBMISSION

Submission to this journal proceeds totally online. Use the following guidelines to prepare your article. Via the homepage of this journal <http://sleepscience.gnpapers.com.br> you will be guided stepwise through the creation and uploading of the various files. All correspondence, including notification of the Editor's decision and requests for revision, takes place by e-mail and via the author's homepage, removing the need for a hard-copy paper trail. If you are unable to provide an electronic version, please contact the editorial office prior to submission (email: sleepscience@sleepscience.com.br).

Article structure

Divide your article into clearly defined and numbered sections. Subsections should be numbered 1.1 (then 1.1.1, 1.1.2, ...), 1.2, etc. (the abstract is not included in section numbering). Use this numbering also for internal cross-referencing: do not just refer to 'the text'. Any subsection may be given a brief heading. Each heading should appear on its own separate line.

Title page: The title page should include the title and a running title to be used as a page heading, which should not exceed 60 letters and spaces; and a declaration of any and all sources of funding.

Abstract and keywords: The abstract should present the information in such a way that the reader can easily understand without referring to the main text and should be followed by 3 to 6 keywords in English defining the main subjects of the study.

Tables and Figures: All tables and figures should be in black and white or in gray scales, on separate pages. Tables and figures should be included in the step 5, the figures should be included in step 6 of the submission process. Photographs depicting surgical procedures, as well as those showing the results of exams or biopsies, will be considered for publication in color, at no additional cost to the authors, provided that they are essential for the manuscript. Each table and figure should be numbered with an Arabic numeral corresponding to its citation in the text.

Legends: Table legends should accompany the respective tables in the main manuscript file. For figure legends they should be inserted on a separate page, after the references. In addition, all abbreviations, acronyms, and symbols should be defined below each table or figure in which they appear.

References: References should be listed in order of their appearance in the text and should be numbered consecutively with Arabic numerals. The presentation should follow the Vancouver Style, according to the examples below. The titles of the journals listed should be abbreviated according to the style presented by the List of Journals Indexed in the Index Medicus of the National Library of Medicine, available at: <http://www.ncbi.nlm.nih.gov/entrez/journals/loftext.noprov.html>. A total of six authors may be listed. For works with more than six authors, list the first six, followed by 'et al.'

Important notes

- **Corresponding author :** Clearly indicate who will handle correspondence at all stages of refereeing and publication, also post-publication. Ensure that the e-mail address is given and that contact details are kept up to date by the corresponding author.
- **Keywords:** Immediately after the abstract, provide a maximum of 6 keywords, using American spelling and avoiding general and plural terms and multiple concepts (avoid, for example, 'and', 'of'). Be sparing with abbreviations: only abbreviations firmly established in the field may be eligible. These keywords will be used for indexing purposes.
- **Acknowledgements :** Collate acknowledgements in a separate section at the end of the article before the references and do not, therefore, include them on the title page, as a footnote to the title or otherwise. List here those individuals who provided help during the research (e.g., providing language help, writing assistance or proof reading the article, etc.).