

A comparison of *Helicobacter pylori* and non - *Helicobacter pylori* *Helicobacter* spp. Binding to Canine Gastric Mucosa with Defined Gastric Glycophenotype

Irina Amorim,^{*,†} Daniela P. Freitas,[†] Ana Magalhães,[†] Fa_tima Faria,^{*} Celia Lopes,^{*} Augusto M. Faustino,^{*} Annemieke Smet,[‡] Freddy Haesebrouck,[‡] Celso A. Reis^{†,§,¶} and Fa_tima Gaertner^{*,†}

*Department of Pathology and Molecular Immunology of the Institute of Biomedical Sciences Abel Salazar (ICBAS), University of Porto, Rua Jorge Viterbo Ferreira nr. 228, Porto 4050-313, Portugal,

†Institute of Molecular Pathology and Immunology of the University of Porto (IPATIMUP), Rua Dr. Roberto Frias s/n, Porto 4200-465, Portugal,

‡Department of Pathology, Bacteriology and Avian Diseases, Faculty of Veterinary Medicine, Ghent University, Salisburylaan 133, Merelbeke B-9820, Belgium,

§Department of Molecular Biology of the Institute of Biomedical Sciences Abel Salazar (ICBAS), University of Porto, Rua Jorge Viterbo Ferreira nr. 228, Porto 4050-313, Portugal,

¶Department of Pathology and Oncology, Faculty of Medicine of Porto, Alameda Prof. Hern^ani Monteiro, Porto 4200-319, Portugal

Originally published in *Helicobacter*. 2014 Aug;19(4):249-59. doi: 10.1111/hel.12125. Epub 2014 Apr 2.

Keywords: Canine gastric mucosa; *Helicobacter pylori*; bacterial adhesion; histo blood group antigens; non-*Helicobacter pylori* helicobacters

ABSTRACT

Background

The gastric mucosa of dogs is often colonized by non - *Helicobacter pylori* helicobacters (NHPH), while *H. pylori* is the predominant gastric *Helicobacter* species in humans. The colonization of the human gastric mucosa by *H. pylori* is highly dependent on the recognition of host glycan receptors. Our goal was to define the canine gastric mucosa glyco-phenotype and to evaluate the capacity of different gastric *Helicobacter* species to adhere to the canine gastric mucosa.

Materials and Methods

The glycosylation profile in body and antral compartments of the canine gastric mucosa, with focus on the expression of histo - blood group antigens was evaluated. The in vitro binding capacity of FITC - labeled *H. pylori* and NHPH to the canine gastric mucosa was assessed in cases representative of the canine glycosylation pattern.

Results

The canine gastric mucosa lacks expression of type 1 Lewis antigens and presents a broad expression of type 2 structures and A antigen, both in the surface and glandular epithelium. Regarding the canine antral mucosa, *H. heilmannii* s.s. presented the highest adhesion score whereas in the body region the SabA - positive *H. pylori* strain was the strain that adhered more.

Conclusions

The canine gastric mucosa showed a glycosylation profile different from the human gastric mucosa suggesting that alternative glycan receptors may be involved in *Helicobacter* spp. binding. *Helicobacter pylori* and NHPH strains differ in their ability to adhere to canine gastric mucosa. Among the NHPH, *H. heilmannii* s.s. presented the highest adhesion capacity in agreement with its reported colonization of the canine stomach.

INTRODUCTION

The gastric mucosa of dogs is often colonized by helicobacters different from *Helicobacter pylori* (*H. pylori*). These non - *H. pylori* helicobacters (NHPH) are present in 67-86% of clinically healthy dogs, in 61-100% of animals presenting chronic vomiting 1-5 and in about 100% of laboratory dogs and dogs from local shelters 6-8. The predominant gastric *Helicobacter* species in dogs are *H. felis*, *H. bizzozeronii*, and *H. heilmannii sensu stricto* (s.s.), while *H. salomonis* is less often detected and the prevalence of *H. cynogastricus* and *H. baculiformis* has not yet been studied 9-12. Mixed infections with different species can also occur 13. The pathogenic significance of gastric NHPH in dogs is poorly understood and remains controversial; therefore veterinarians are facing the dilemma of either treating or ignoring spiral organisms observed in canine gastric biopsies. However, these NHPH are of zoonotic significance and have been associated with gastritis, peptic ulcers and mucosa associated lymphoid tissue lymphomas in human patients 11. Dogs may constitute a source of infection for their owners, although the prevalence of these canine - and feline - associated helicobacters in humans is much lower than that of *H. pylori*. On the other hand, *H. pylori* has occasionally been identified in the canine stomach 14, 15. Gastric *Helicobacter* species have a host species preference, but may occasionally cross this host species barrier. The molecular mechanisms underlying this partial host - adaptation are not known, but may be related to differences in glycosylation profile of the gastric mucosa.

The gastrointestinal tract glycosylation profile determines the colonization capability of various infectious agents. Bacterial binding occurs through recognition of specific glycan receptors expressed by the host epithelial cells 16-18. The biosynthesis of glycan chains is controlled by enzymatic activity of several glycosyltransferases, which are expressed in a cell, tissue and species - specific manner (for review see 19).

The expression of the terminal Lewis glycan structures that are recognized as receptors for *H. pylori* in human gastric tissue depends on the enzymatic activity of specific fucosyltransferases and sialyltransferases 20 and are schematically represented in Fig. 1A, B. Two backbone structures are the precursors for the Lewis antigens biosynthesis: the type 1 Gal β 1 - 3GlcNAc, and the type 2 Gal β 1 - 4GlcNAc chains. Addition of a fucose to terminal galactose on type 1 structures leads to H - type 1 structure, which can be further modified with a fucose on the GlcNAc residue resulting in Lewis b (Leb). Alternatively, the type 1 backbone may be fucosylated on the GlcNAc residue leading to Lewis a (Lea) structure. Furthermore, the action of a sialyltransferase toward type 1 chains can lead to the biosynthesis of the sialyl - Lea (sLea) antigen. The biosynthesis of type 2 based Lewis antigens depends on the addition of the same glycan units but with different linkages, originating the isomers Lewis x (Lex), Lewis y (Ley) and sialyl - Lex (sLex) antigens (Fig. 1B) (as reviewed by 20).

The *H. pylori* blood group antigen - binding adhesin (BabA) recognizes both H - type 1 and Leb fucosylated antigens 16 expressed on the surface of the gastric mucosa of secretor individuals 21, 22. The secretor status is determined by the activity of the fucosyltransferase 2 enzyme (FUT2) 23 and the capacity to synthesize H - type 1 and Leb antigens in body secretions. The secretor and Lewis status in humans are associated with the adhesion and infection of *H. pylori* strains expressing the BabA adhesin 21, 24, 25. In addition, a second *H. pylori* adhesin, the sialic acid - binding adhesin (SabA) has been described, which recognizes the sialylated antigens sLea and sLex 17. These sialylated structures are absent in normal human gastric mucosa but are induced upon *H. pylori* infection and gastric mucosa inflammation 17, 26, 27.

The canine secretory alloantigen alloantibody system (CSA) is closely related to the human ABH - Le system being equivalent to the major human ABO exocrine glycophenotype. The CSA was first described in 1966 by Zweibaum and colleagues 28. The genetic polymorphisms of the CSA were later analysed 29 and the structure of canine polymorphic antigens was further characterized in four phenotypes: A, X, Y, and AY 30 (Fig. 1C). However, limited information is available about the glycosylation profile of the canine gastric mucosa.

Here we describe, for the first time, the glycophenotype observed in the canine gastric mucosa, with focus on the expression of Lewis glycan antigens. Furthermore, the binding capacity of *H. pylori* and NHPH to canine gastric mucosa was evaluated. This study provides valuable information about the canine gastric glycosylation profile and about the *Helicobacter* capacity to adhere to the canine gastric mucosa, contributing to the understanding of the host colonization spectrum of the different *Helicobacter* species.

MATERIALS AND METHODS

Tissue Samples and Histology

Gastric samples were obtained during necropsy procedures from dogs that died from non-infectious causes (eight representative sections of the body and six of the antrum of the stomach).

Tissues were fixed in 10% buffered formalin and paraffin-embedded. Serial sections 3 μ m thick were made, one being stained with haematoxylin and eosin (HE) for histopathology and the others were used for the histochemical and immunohistochemical studies.

Sections stained with HE were examined by three pathologists and were considered as normal gastric tissues, according to the criteria proposed by Prachasilpchain et al. 31. Additionally, the presence of *Helicobacter* spp. was excluded based on the negative results of modified Giemsa stain and of anti-*H. pylori* immunohistochemistry, using a polyclonal antibody (RBK012; Zytomed, Berlin, Germany), which shows immunoreactivity with a wide range of bacteria belonging to the *Helicobacter* genus.

Immunohistochemistry

For the immunohistochemical study, sections were deparaffinized, hydrated and antigen retrieval was performed in a pressure cooker in 10 mmol/L sodium citrate buffer, pH 6.0, for 2 minutes. Slides were cooled for 10 minutes at room temperature and rinsed twice in triphosphate buffered saline (TBS) for 5 minutes. The Novolink™ Max - Polymer detection system (Novocastra, Newcastle, UK) was used for visualization, according to the manufacturer's instructions. After blocking endogenous peroxidase with 3% hydrogen peroxide in methanol for 10 minutes, sections were incubated, overnight at 4 °C, with the monoclonal antibodies specific for the carbohydrate antigens (Table 1). Sections were rinsed with TBS between each step of the procedure. Color was developed for up to 7 minutes at room temperature with 3,3'-diamino-benzidine (DAB; Sigma, St. Louis, MO, USA) and sections were then lightly counterstained with haematoxylin, dehydrated, and mounted.

Sections of human gastric mucosa with intestinal metaplasia were obtained from the pathology department archive of Hospital Santo Antonio – Porto and were used as positive controls. Negative controls were performed by replacing the primary antibody with an antibody of the same immunoglobulin isotype.

Lectin Staining

The biotinylated lectin *Dolichus biflorus* agglutinin (DBA), for detection of terminal alpha linked GalNAc, and the avidin-biotin-peroxidase complex (Vectastain Elite ABC kit) solution were purchased from Vector Laboratories (Burlingame, CA, USA). After deparaffination and rehydration, endogenous peroxidase activity was blocked with 3% hydrogen peroxide in methanol for 10 minutes at room temperature. After washing twice with phosphate buffered saline (PBS) for 5 minutes, sections were incubated with 10% bovine serum albumin (BSA) in PBS for 30 minutes at room temperature and then in the biotinylated lectin solution (1 : 100 in PBS) for 1 hour, at room temperature. The slides were subsequently washed in PBS and immersed in ABC solution for 30

minutes at room temperature, according to manufacturer's recommendations. After washing twice with PBS for 5 minutes, slides were immersed in a freshly prepared solution of DAB containing 0.02% hydrogen peroxide for 7 minutes at room temperature to visualize the lectin binding sites, counterstained with haematoxylin, dehydrated and mounted. Sections of canine normal intestine were used as positive controls. Negative controls were performed by replacing the lectin by PBS.

Evaluation of Gastric Tissue Glycosylation Profile

A microscopic evaluation was performed by analyzing the entire section of the gastric tissue. Positive immunoreactivity was recorded as a distinct brown labeling of cytoplasm and/or membrane of epithelial cells. Extracellular staining of the superficial mucus was excluded from our evaluation. The immunostaining was scored according to the intensity (0, absent; +, weakly positive; ++, moderately positive; or + + +, intense stain) and distribution (sparse, focal, multifocal, and generalized).

Bacterial Strains

The *H. pylori* strains 17875/Leb and 17875babA1::kan babA2::cam (17875babA1A2) were obtained from the Department of Medical Biochemistry and Biophysics, Umeå University, Sweden 17. These model strains were selected based on their capacity to adhere to human gastric mucosa tissue sections. The 17875/Leb strain is a spontaneous mutant which expresses both BabA and SabA adhesins, however, does not show sialylated dependent binding. The 17875babA1A2 is a BabA adhesin mutant strain which expresses a functional SabA adhesin 17.

Helicobacter pylori were grown in Pylori agar (BioMérieux, Marcy l'Étoile, France) at 37 ° C under microaerobic conditions (85% N₂, 10% CO₂, 5% O₂; 37 ° C). For strain babA1A2 - mutant, culture media included also 20 mg/L Chloramphenicol (Sigma) and 25 mg/L Kanamycin (Sigma).

The NHPH strains were provided by the Department of Pathology, Bacteriology and Avian Diseases, Faculty of Veterinary Medicine, Ghent University. The *H. felis* CS₁, *H. salomonis* R1053 and *H. bizzozeronii* R1051 strains were cultured biphasically on Brain Haert Infusion agar plates (BHI; Oxoid, Basingstoke, UK) supplemented with 10% horse blood, 20 µg/mL amphotericin B (Fungizone; Bristol - Myers Squibb, Eppernon, France), Vitox supplement (Oxoid), and Campylobacter selective supplement (Skirrow, containing 10 µg/mL vancomycin, 5 µg/mL trimethoprim lactate, and 2.5 U/mL; Oxoid). The *H. heilmannii* strain ASB1.4 was cultivated biphasically on Brucella agar plates (Oxoid) supplemented with 20% (v/v) fetal calf serum (HyClone, Logan, UT), 5 mg/L amphotericin, Skirrow and Vitox supplements (Oxoid), and 0.05% HCl (pH 5). All strains were incubated under microaerobic condition.

In vitro Binding Assay of *Helicobacter pylori* and NHPH to Canine Gastric Mucosa

For the in vitro *Helicobacter* binding assay we selected three representative cases (n = 3) of the general canine glycosylation pattern, belonging to three different dogs, in which body and antrum samples were available.

The different *Helicobacter* strains were labeled with fluorescein isothiocyanate (FITC) and concentration adjusted as previously described 32. Adhesion assays were performed as previously described 22, 32 with the following modifications. Briefly, paraffin embedded canine gastric sections were deparaffinized and rehydrated followed by overnight incubation at 4 °C and 1 hour at room temperature, with blocking buffer: 1% BSA in PBS containing 0.05% Tween 20. BSA was previously submitted to Periodate oxidation to destroy competitive carbohydrate receptors for *Helicobacter* binding 33. The FITC - labeled bacterial suspension was diluted 5 - fold in blocking buffer and 100 μ L was used to incubate each section for 2 hours at room temperature. Slides were subsequently washed five times with PBS containing 0.05–0.1% Tween 20 and stained with DAPI. Evaluation was estimated by the number of adhered bacteria under 100 \times magnification, and analyzing at least five different fields per case and quantified using the ImageJ software (NIH, Bethesda, MD, USA).

Statistical Analysis

The Student's t - test was used to compare the differences between the mean adhesion values of both stomach regions for the same NHPH strain. The significance of differences between mean values of each *Helicobacter* strain was assessed using a one - way ANOVA (GraphPad Prism, GraphPad Software, Inc., La Jolla, CA, USA). p values < 0.05 were considered as statistically significant.

RESULTS

Expression of Type 1 Lewis Antigens in Dog Gastric Mucosa

There was no Lea antigen expression in the body region of the stomach from seven of the eight canine cases studied (Fig. 2A, Table 2). Only in one case, a very weak and sparse Lea expression was detected in few superficial foveolar epithelial cells. In the antral region of the stomach, there was no Lea antigen expression in all six cases studied (Fig. 2B, Table 2). Lewisb and sLea expression were absent in both the body and antrum of the stomach (Fig. 2D,E,G,H, Table 2).

Expression of Type 2 Lewis Antigens in Dog Gastric Mucosa

In the body of the stomach, expression of Lex was seen in five of the eight canine cases (Table 2). The staining intensity varied from moderate to strong, always limited to the superficial foveolar epithelium (Fig. 2J). In all positive cases, the staining was multifocal and restricted to the cytoplasm. In the antrum of the stomach, Lex expression was found in all cases. The staining intensity varied from weak to strong and was limited to the superficial foveolar epithelium. The pattern of staining was multifocal in four cases and generalized in the other two cases (Fig. 2K, Table 2).

All cases evaluated presented Ley in the body and antrum. In the body, the level of immunostaining was weak in one case and intense in seven cases. The staining included superficial foveolar epithelium and parietal cells (Fig. 2M). The distribution of the staining was multifocal in four cases and generalized in the other four. The subcellular staining was mainly localized in the cytoplasm of the cells. In the antrum, the level of Ley immunostaining varied from moderate to intense, presenting stronger labeling in the cells of the foveolar epithelium and gastric pits. The pattern of staining was generalized in all cases (Fig. 2N, Table 2).

In the body of the stomach, sLex expression was noted in four cases. The level of immunostaining varied from weak to strong. The staining included superficial foveolar epithelium (Fig. 2P). The staining distribution was multifocal in all cases studied. The subcellular staining was mainly localized in the cytoplasm of the cells. In the antrum of the stomach, a sparse and moderate expression of sLex was observed in only one case. The staining was limited to the cytoplasm of the foveolar superficial epithelium and cells of the pyloric glands (Fig. 2Q, Table 2).

Expression of Terminal α - GalNAc in Dog Gastric Mucosa

In the body of the stomach, expression of terminal α - GalNAc was found in all canine cases, as detected by DBA lectin labeling. The staining was moderate in three cases and intense in the remaining five cases. The staining included superficial foveolar epithelium and parietal cells (Fig. 2S). The staining pattern was generalized in all cases. The subcellular staining was localized in the cytoplasm of the cells. In the antrum of the stomach, generalized expression of terminal α - GalNAc was detected in all cases. The level of staining was moderate in four cases (Fig. 2T) and intense in the other two. The staining included the foveolar epithelium, gastric pits and pyloric glands. Generally, the superficial pyloric glands showed an intense staining higher than that recorded in the deeper glands; only in one case the staining was identical in both areas. The subcellular staining was localized in the cytoplasm of the cells. (Table 2).

In vitro Binding Assay of *Helicobacter pylori* and NHPH to Canine Gastric Mucosa

The FITC labeled *Helicobacter* strains were verified for size and morphology in accordance with literature: *H. pylori* strains measured about 2.5–5 μm in length and 0.5–1 μm in width (Fig. 3A,B); The NHPH strains were larger and apparently more tightly coiled spiral - shaped bacteria: ASB1.4 was, approximately 3–6.5 μm long and 0.6–0.7 μm wide (Fig. 3C); CS1 measured about 5–7.5 μm in length and 0.4 in width (Fig. 3D) while R1051 ranged from 5 to 10 μm in length and 0.3 in width (Fig. 3E). R1053 was less spiral, measuring 5–7 μm in length and 0.8–1.2 μm wide (Fig. 3F) (as reviewed by 34). The length was variable depending on the state of contraction.

Regarding tissue sections from the body of canine stomach, we observed that the *H. pylori* strain 17875babA1A2 (SabA+) had the highest adhesion. In contrast, the *H. pylori* strain 17875/Leb (BabA+) showed low levels of adhesion. Among the NHPH strains evaluated, ASB1.4 presented the highest adhesion value, followed by CS1 and R1051. The R1053 was the strain which adhered less being possible to observe gastric regions completely devoid of bacterial adhesion (Fig. 4).

In the antral sections, the *H. pylori* strains 17875babA1A2 and 17875/Leb showed a similar binding behavior, exhibiting very closely means values of adhesion. The quantification revealed that ASB1.4 was the strain that presented highest adhesion values to the canine antrum tissue sections. Additionally, CS1 and R1051 strains adhered similarly to canine antral mucosa, whereas R1053 adhered less (Fig. 5).

The ASB1.4 adhesion value quantified in the antral mucosa was significantly higher from that obtained in the body region of the canine stomach ($p < .05$). No significant differences were found in CS1, R1051, and R1053 bacterial adhesion, concerning both canine stomach compartments.

DISCUSSION

In the present study, we have characterized the expression of histo - blood group antigens in the normal canine gastric mucosa to define the gastric glycosylation profile of dogs (Table 2). The expression pattern of type 1 and type 2 Lewis antigens in the canine gastric mucosa revealed a different distribution when compared with the human gastric mucosa. Normal canine mucosa showed minor expression of type 1 Lewis antigens, with absence of Leb and sLea expression and only a sparse expression of Lea in the body of one of the animals tested. In the human normal gastric mucosa, and depending on the secretor status of the individual, Lea and Leb are highly expressed in the superficial foveolar epithelium 23, 35, where MUC5AC mucin is co - expressed 36. The normal canine mucosa demonstrated expression of Lewis type 2 antigens characterized by expression of Lex and Ley, along with a minor expression of sLex. In the canine gastric mucosa this expression was mostly observed in the superficial and foveolar epithelium, with Ley expression extending to the deeper portions of the gastric gland. In the human gastric mucosa, the expression of Lewis type 2 antigens is generally restricted to the deeper part of the gastric glands, where MUC6 mucin is co - expressed 37, 38. Knowing that type 1 fucosylated structures are receptors for *H. pylori* in human stomach, the different expression patterns of type 1 and 2 Lewis antigens observed may play a role in differences in prevalence of gastric helicobacters in both mammalian species.

The pattern of Lewis antigen expression observed in canine gastric mucosa is in agreement with a previous study which reported a prominent representation of type 2 chains in dog intestinal secretions 39.

According to our results, the A - antigen is also extensively expressed in the normal canine gastric mucosa, representing an important component of the dog gastric mucosa glyco phenotype (Table 2). This evidence is in accordance with earlier observations documenting that this antigen is expressed in canine gastrointestinal tissues 30. Interestingly, the pattern of expression of the A - antigen in canine gastric mucosa resembles the reactivity of DBA lectin previously reported in human gastric tissue 40. However, its putative involvement in *Helicobacter* adhesion has never been addressed.

Lewis antigens biosynthesis is dependent on the cells repertoire of glycosyltransferases, namely fucosyltransferases. The absence of type 1 Lewis antigens in dog gastric mucosa may be explained by the lack of expression of enzymes with $\alpha_{1,4}$ fucosyltransferase activity. The human enzymes with this activity, FUT3 and FUT5, seem to have originated late in hominid evolution from duplication of an ancestor Lewis gene present in lower mammals 41. Moreover, it has been demonstrated that the acceptor substrate specificity of vertebrate $\alpha_{1,3/1,4}$ - fucosyltransferases is determined by a single amino acid in the hypervariable stem domain 42. Most vertebrate $\alpha_{1,3}$ fucosyltransferases present a conserved arginine in this domain while a tryptophan residue is present at the same position on human FUT3 and FUT5. Noteworthy, the dog gene orthologous of the human FUT3 and FUT5 gene presents at this position an Arg residue, compatible with $\alpha_{1,3}$ fucosyltransferase activity and therefore type 2 Lewis synthesis (Fig. S1).

The host glycan receptors are recognized by *Helicobacter* outer membrane proteins (OMPs) which mediate several important host - pathogen interactions 18. Recently, it has been shown that NHPH OMPs genes share sequence homology with *H. pylori*. Interestingly, it was found that *H. acinonychis*, a NHPH colonizing big cats, express BabA adhesin 43.

However, among the NHPH strains included in this study, none presented in their genomes orthologues of *H. pylori* adhesins. *H. bizzozeronii* and *H. mustelae* lack homologs of major *H. pylori* virulence genes BabA/B and SabA while *H. suis* lacks BabA/B and SabA/B 44-46. The genome of *H. heilmannii* s.s. showed that this strain encodes several OMPs but lacks the Bab and Sab adhesins 47.

The absence of homologues of the BabA in these NHPH may underlie the low rate of human infection by these organisms and suggests that other adhesins with different receptor specificity may play a role in the colonization of NHPH. Additionally, the lack of Lewis type 1 antigens expression in the canine gastric mucosa might explain why *H. pylori* rarely colonizes the stomach of dogs.

To determine if the glycosylation profile of canine gastric mucosa was related to the host - adaptation of the different NHPH, we evaluated the binding capacity of some *Helicobacter* species to canine gastric mucosa.

Our results show that *H. heilmannii* s.s. adhered most to the canine gastric mucosa. This result highlights the contribution of adhesion in the canine gastric colonization by this species.

In 2012, *H. heilmannii* was isolated and described as a novel species 34. The name *H. heilmannii* sensu stricto (s.s.) was proposed to refer to the novel *Helicobacter* species and the term *H. heilmannii* sensu lato (s.l.) to refer to the whole group NHPH 12. Since the discovery and description of *H. heilmannii* s.s. is still relatively recent, the majority of the epidemiological data available does not comprise the identification of this particular species in the stomach of dogs and cats. This impairs the comparison of the adhesion capacity achieved in this experiment with the spontaneous incidence of *H. heilmannii* s.s. in dogs.

We also found that the *H. heilmannii* s.s. adhesion levels were higher in the antrum than in the body region of canine stomach ($p < .05$). In agreement with our observations, Joosten et al. 48 recently studied bacterium - host interactions of 9 *H. heilmannii* s.s. strains in Mongolian gerbils and concluded that, in general, the colonization capacity in the fundus was lower than in the antrum for all strains tested. Additionally, a clear association was observed between the colonization capacity of the *H. heilmannii* s.s. strains and the gastric inflammation scores in the antrum of the stomach. In dogs, while many studies reported that the fundus and body present higher bacteria density and higher probability to find *Helicobacter* spp. 8, 49, 50 others found no significant differences between the density of NHPH in the fundus, body and antrum 31, 51-54. The discrepancies in the results regarding the spatial distribution of *Helicobacter* spp. in canine gastric mucosa can be attributed to the different laboratorial approaches and diagnostic methodologies used by the various research groups.

The other NHPH strains tested showed lower adhesion levels to the canine gastric mucosa. In comparison to *H. heilmannii* s.s., *H. felis* has a lower adhesion capacity, followed by *H. bizzozeronii* and *H. salomonis* and this trend was observed in the body as well as in the antrum.

The two *H. pylori* strains used in this study are model strains frequently used in adhesion studies due to their well - defined binding properties. In both stomach regions, the *H. pylori* strain 17875/Leb, which is a BabA positive strain, adhered less than *H. heilmannii* s.s. This can be explained by the fact that dogs do not express Lewis type 1 antigens, including the BabA - ligand Leb. The *H. pylori* 17875babA1A2, which is a SabA positive strain, adhered considerably in the body region. This result is in accordance with the concomitant observed expression of sLex, the ligand of the SabA adhesin. However, in vivo and probably due to competitive factors between bacteria and/or different immunological responses of the host, *H. pylori* often fails to colonize the canine gastric mucosa.

Our results support that the differences in gastric mucosa glycosylation profile play a role in the host adaptation of gastric *Helicobacter* species. The defined canine gastric mucosa glyco-phenotype indicates that adhesion of NHPH involves different adhesion mechanisms mediated by proteins with alternative receptor specificity. The knowledge of the adhesion abilities of different *Helicobacter* strains to canine gastric mucosa is of outmost relevance for the understanding of the physiopathology of *Helicobacter* infection in the dogs.

Acknowledgements and Disclosures

We kindly thank Prof. Thomas Borén from the Department of Medical Biochemistry and Biophysics, Umeå University, Sweden for providing the 17875/Leb and 17875babA1A2 *H. pylori* strains. The authors thank Dr. Fernando Rodrigues, Dr. Ana Laura Saraiva, and Cristina Bacelar who kindly provided technical support. I. Amorim (SFRH/BD/76237/2011) and A. Magalhães (SFRH/BPD/75871/2011) acknowledge FCT for financial support. This study was partially funded by the Portuguese Foundation for Science and Technology (PTDC/CTM - BPC/121149/2010; PTDC/CVT/117610/2010; PTDC/BBB - EBI/0786/2012). The Institute of Molecular Pathology and Immunology of the University of Porto (IPATIMUP) is an Associate Laboratory of the Portuguese Ministry of Science, Technology and Higher Education and is partially supported by FCT.

REFERENCES

- 1 Geyer C, Colbatzky F, Lechner J, Hermanns W. Occurrence of spiral-shaped bacteria in gastric biopsies of dogs and cats. *Vet Rec* 1993;133:18–9.
- 2 Hermanns W, Kregel K, Breuer W, Lechner J. *Helicobacter*-like organisms: Histopathological examination of gastric biopsies from dogs and cats. *J Comp Pathol* 1995;112:307–18.
- 3 Lecoindre P, Chevallier M, Peyrol S, Boude M, De Montclos H. Contribution à l'étude des *helicobacteries* de l'estomac du chien et de leur rôle pathogène. *Revue Médicine Veterinaire* 1995;146:671–80.

- 4 Hwang CY, Han HR, Youn HY. Prevalence and clinical characterization of gastric *Helicobacter* species infection in dogs and cats in Korea. *J Vet Sci* 2002;3:123–33.
- 5 Polanco R, Salazar V, Reyes N, Garcia-Amado MA, Michelan-geli F, Contreras M. High prevalence of DNA from non-*H. pylori* *Helicobacter* species in the gastric mucosa of Venezuelan pet dogs and its histological alterations. *Rev Inst Med Trop Sao Paulo* 2011;53:207–12.
- 6 Henry GA, Long PH, Burns JL, Charbonneau DL. Gastric spirilliosis in beagles. *Am J Vet Res* 1987;48:831–6.
- 7 Eaton KA, Dewhirst FE, Paster BJ, Tzellas N, Coleman BE, Paola J, Sherding R. Prevalence and varieties of *Helicobacter* species in dogs from random sources and pet dogs: animal and public health implications. *J Clin Microbiol* 1996;34:3165–70.
- 8 Simpson KW, Strauss-Ayali D, McDonough PL, Chang YF, Valentine BA. Gastric function in dogs with naturally acquired *Helicobacter* species infection. *J Vet Intern Med* 1999;13:507–15.
- 9 Van den Bulck K, Decostere A, Baele M, Vandamme P, Mast J, Ducatelle R, Haesebrouck F. *Helicobacter cynogastricus* sp. nov., isolated from the canine gastric mucosa. *Int J Syst Evol Microbiol* 2006;6:1559–64.
- 10 Baele M, Pasmans F, Flahou B, Chiers K, Ducatelle R, Haesebrouck F. Non-*Helicobacter pylori* *Helicobacter* species detected in the stomach of humans comprise several naturally occurring *Helicobacter* species in animals. *FEMS Immunol Med Microbiol* 2009;55:306–13.
- 11 Haesebrouck F, Pasmans F, Flahou B, Chiers K, Baele M, Meyns T, Decostere A, Ducatelle R. Gastric *Helicobacter* in domestic animals and nonhuman primates and their significance for human health. *Clin Microbiol Rev* 2009;22:202–23.
- 12 Haesebrouck F, Pasmans F, Flahou B, Smet A, Vandamme P, Ducatelle R. Non-*Helicobacter pylori* *Helicobacter* species in the human gastric mucosa: a proposal to introduce the term *H. heilmannii sensu lato* and *sensu stricto*. *Helicobacter* 2011;16:339–40.
- 13 Kman E, Fredriksson M, Trowald-Wigh G. *Helicobacter* spp. in the saliva, stomach, duodenum and faeces of colony dogs. *Vet J* 2013;195:127–9.
- 14 Taulescu M, Catoi C, Gal A, Bolfa P, Rus VI, Budugan M, Tabaran F, Nagy A. Diagnosis of spontaneous gastric infection with *Helicobacter* species in dogs using PCR method. *Lucrari stiintifice USAMV Iasi Med Vet* 2009;52:304–10.
- 15 Chung TH, Kim HD, Lee YS, Hwang CY. Determination of the prevalence of *Helicobacter heilmannii*-like organisms type 2 (HHLO-2) infection in humans and dogs using non-invasive genus/species-specific PCR in Korea. *J Vet Med Sci* 2013;76:73–9.
- 16 Ilver D, Arnqvist A, Ogren J, Frick IM, Kersulyte D, Incecik ET, Berg DE, Covacci A, Engstrand L, Boren T. *Helicobacter pylori* adhesin binding fucosylated histo-blood group antigens revealed by retagging. *Science* 1998;16:373–7.

- 17 Mahdavi J, Sonden B, Hurtig M, et al. Helicobacter pylori SabA adhesin in persistent infection and chronic inflammation. *Science* 2002; 297: 573–8.
- 18 Magalhães A, Reis CA. Helicobacter pylori adhesion to gastric epithelial cells is mediated by glycan receptors. *Braz J Med Biol Res* 2010; 43: 611–8.
- 19 Reis CA, Osorio H, Silva L, Gomes C, David L. Alterations in glycosylation as biomarkers for cancer detection. *J Clin Pathol* 2010; 6: 322–9.
- 20 Magalhães A, Ismail MN, Reis AC. Sweet receptors mediate the adhesion of the gastric pathogen Helicobacter pylori: glycoproteomic strategies. *Expert Rev Proteomics* 2010; 7: 307–10.
- 21 Azevedo M, Eriksson S, Mendes N, et al. Infection by Helicobacter pylori expressing the BabA adhesin is influenced by the secretor phenotype. *J Pathol* 2008; 215: 308–16.
- 22 Magalhães A, Gomes J, Ismail MN, et al. Fut2-null mice display an altered glycosylation profile and impaired BabA-mediated Helicobacter pylori adhesion to gastric mucosa. *Glycobiology* 2009; 19: 1532–6.
- 23 Mollicone R, Bara J, Le Pendu J, Oriol R. Immunohistologic pattern of type 1 (Lea, Leb) and type 2 (X, Y, H) blood group-related antigens in the human pyloric and duodenal mucosae. *Lab Invest* 1985; 53: 219–27.
- 24 Ikehara Y, Nishihara S, Yasutomi H, et al. Polymorphisms of two fucosyltransferase genes (Lewis and Secretor genes) involving type I Lewis antigens are associated with the presence of anti-Helicobacter pylori IgG antibody. *Cancer Epidemiol Biomarkers Prev* 2001; 10: 971–7.
- 25 Lee HS, Choe G, Kim WH, Kim HH, Song J, Park KU. Expression of Lewis antigens and their precursors in gastric mucosa: relationship with Helicobacter pylori infection and gastric carcinogenesis. *J Pathol* 2006; 209: 88–94.
- 26 Ota H, Nakayama J, Momose M, Hayama M, Akamatsu T, Ka-tsuyama T, Graham DY, Genta RM. Helicobacter pylori infection produces reversible glycosylation changes to gastric mucins. *Virchows Arch* 1998; 433: 419–26.
- 27 Marcos NT, Magalhães A, Ferreira B, et al. Helicobacter pylori induces beta3GnT5 in human gastric cell lines, modulating expression of the SabA ligand sialyl-Lewis x. *J Clin Invest* 2008; 118: 2325–36.
- 28 Zweibaum A, Oudea P, Halpern B, Veyre C. Presence in colic glandular cells of various mammalian species of an antigen cross-reacting with human blood group A substance. *Nature* 1966; 8: 159–61.
- 29 Zweibaum A, Oriol R, Feingold N, Dussaulx E, Rousset M, Chevalier G. Studies on canine secretory alloantigens (CSA). *Tissue Antigens* 1974; 4: 115–29.

- 30 Oriol R, Strecker G, Rousset M, Chevalier G, Dalix AM, Dussaulx E, Zweibaum A. Determination of the structure of the immunodominant groups of the canine secretory alloantigens (CSA) by radioimmunoassay. *Transplant Proc* 1975;7:523–8.
31. Prachasilpachai W, Nuanualsuwan S, Chatsuwana T, Techan-gamsuwan S, Wangnaitam S, Sailasuta A. Diagnosis of *Helicobacter* spp. infection in canine stomach. *J Vet Sci* 2007;8:139–45.
- 32 Falk P, Roth KA, Boroen T, Westblom TU, Gordon JI, Normark S. An in vitro adherence assay reveals that *Helicobacter pylori* exhibits cell lineage-specific tropism in the human gastric epithelium. *Proc Natl Acad Sci USA* 1993;90:2035–9.
- 33 Aspholm M, Olfat FO, Nordoen J, et al. SabA is the *H. pylori* hemagglutinin and is polymorphic in binding to sialylated glycans. *PLoS Pathog* 2006;2:e110.
- 34 Smet A, Flahou B, D’Herde K, Vandamme P, Cleenwerck I, Ducatelle R, Pasmans F, Haesebrouck F. *Helicobacter heilmanni* sp. nov., isolated from feline gastric mucosa. *Int J Syst Evol Microbiol* 2012;62(Pt 2):299–306.
- 35 Lopez-Ferrer A, De Bolos C, Barranco C, Garrido M, Isern J, Carlstedt I, Reis CA, Torrado J, Real FX. Role of fucosyltransferases in the association between apomucin and Lewis antigen expression in normal and malignant gastric epithelium. *Gut* 2000;47:349–56.
- 36 Teixeira A, David L, Reis CA, Costa J, Sobrinho-Simões M. Expression of mucins (MUC1, MUC2, MUC5AC, and MUC6) and type 1 Lewis antigens in cases with and without *Helicobacter pylori* colonization in metaplastic glands of the human stomach. *J Pathol* 2002;197:37–43.
- 37 De Bolos C, Garrido M, Real FX. MUC6 apomucin shows a distinct normal tissue distribution that correlates with Lewis antigen expression in the human stomach. *Gastroenterology* 1995;109:723–34.
- 38 Moran AP. Relevance of fucosylation and Lewis antigen expression in the bacterial gastroduodenal pathogen *Helicobacter pylori*. *Carbohydr Res* 2008;343:1952–65.
- 39 Smith EL, McKibbin JM, Karlsson KA, Pascher I, Samuelsson BE. Characterization by mass spectrometry of blood group A active glycolipids from human and dog small intestines. *Biochemistry* 1975;14:2120–4.
- 40 Kuhlmann WD, Peschke P, Wurster K. Lectin-peroxidase conjugates in histopathology of gastrointestinal mucosa. *Virchows Arch A Pathol Anat Histopathol* 1983;398:319–28.
- 41 Oriol R, Mollicone R, Cailleau A, Balanzino L, Breton C. Divergent evolution of fucosyltransferase genes from vertebrates, invertebrates, and bacteria. *Glycobiology* 1999;9:323–34.
- 42 Dupuy F, Petit JM, Mollicone R, Oriol R, Julien R, Maftah A. A single amino acid in the hypervariable stem domain of vertebrate α 1,3/1,4-fucosyltransferases determines the type 1/type 2 transfer. Characterization of acceptor substrate specificity of the Lewis enzyme by site-directed mutagenesis. *J Biol Chem* 1999;274:12257–62.

- 43 Tegtmeyer N, Rivas Traverso F, Rohde M, Oyarzabal OA, Lehn N, Schneider-Brachert W, Ferrero RL, Fox JG, Berg DE, Backert S. Electron microscopic, genetic and protein expression analyses of *Helicobacter acinonychis* strains from a Bengaltiger. *PLoS One* 2013;8:e71220.
- 44 O'Toole PW, Snelling WJ, Canchaya C, et al. Comparative genomics and proteomics of *Helicobacter mustelae*, an ulcerogenic and carcinogenic gastric pathogen. *BMC Genomics* 2010;11:164.
- 45 Schott T, Kondadi PK, Hanninen ML, Rossi M. Comparative genomics of *Helicobacter pylori* and the human-derived *Helicobacter bizzozeronii* CIII-1 strain reveal the molecular basis of the zoonotic nature of non-*pylori* gastric *Helicobacter* infections in humans. *BMC Genomics* 2011;12:534.
- 46 Vermoote M, Vandekerckhove TT, Flahou B, Pasmans F, Smet A, De Groot D, Van Criekinge W, Ducatelle R, Haesebrouck F. Genome sequence of *Helicobacter suis* supports its role in gastric pathology. *Vet Res* 2011;42:51.
- 47 Smet A, Van Nieuwerburgh F, Ledesma J, Flahou B, Deforce D, Ducatelle R, Haesebrouck F. Genome sequence of *Helicobacter heilmanni* sensu stricto ASB1 isolated from the gastric mucosa of a kitten with severe gastritis. *Genome Announc* 2013;1:1–2.
- 48 Joosten M, Blaecher C, Flahou B, Ducatelle R, Haesebrouck F, Smet A. Diversity in bacterium-host interactions within the species *Helicobacter heilmanni* sensu stricto. *Vet Res* 2013;44:65.
- 49 Yamasaki K, Suematsu H, Takahashi T. Comparison of gastric lesions in dogs and cats with and without gastric spiral organisms. *J Am Vet Med Assoc* 1998;212:529–33.
- 50 Anacleto TP, Lopes LR, Andreollo NA, Bernis Filho WO, Resck MC, Macedo A. Studies of distribution and recurrence of *Helicobacter* spp. gastric mucosa of dogs after triple therapy. *Acta Cir Bras* 2011;26:82–7.
- 51 Cattoli G, van Vugt R, Zanoni RG, Sanguinetti V, Chiochetti R, Gualtieri M, Vandembroucke-Grauls CM, Gaastra W, Kusters JG. Occurrence and characterization of gastric *Helicobacter* spp. in naturally infected dogs. *Vet Microbiol* 1999;70:239–50.
- 52 Buczolits S, Hirt R, Rosengarten R, Busse HJ. PCR-based genetic evidence for occurrence of *Helicobacter pylori* and novel *Helicobacter* species in the canine gastric mucosa. *Vet Microbiol* 2003;95:259–70.
- 53 Recordati C, Gualdi V, Craven M, Sala L, Luini M, Lanzoni A, Rishniw M, Simpson KW, Scanziani E. Spatial distribution of *Helicobacter* spp. in the gastrointestinal tract of dogs. *Helicobacter* 2009;14:180–91.
- 54 Gombac M, Svara T, Cerne M, Pogacnik M. Histological changes in stomachs of apparently healthy dogs infected with *Helicobacter*. *Acta Vet (Beogr)* 2010;60:173–82.
- 55 Young WW Jr, Portoukalian J, Hakomori S. Two monoclonal anticarbohydrate antibodies directed to glycosphingolipids with a laco-N-glycosyl type II chain. *J Biol Chem* 1981;256:10967–72.



56 Fukushi Y, Hakomori S, Nudelman E, Cochran N. Novel fucosylated oligosaccharides accumulating in human adenocarcinoma. II. Selective isolation of hybridoma antibodies that differentially recognize mono-, di-, and trifucosylated type 2 chain. *J Biol Chem* 1984;259:4681-5.

57 Abe K, McKibbin JM, Hakomori S. The monoclonal antibody directed to difucosylated type 2 chain (Fuc alpha 1 leads to 2Gal beta 1 leads to 4[Fuc alpha 1 leads to 3]GlcNAc; Y Determinant). *J Biol Chem* 1983;258:11793-7.

**INSTITUTO
DE INVESTIGAÇÃO
E INOVAÇÃO
EM SAÚDE**
UNIVERSIDADE
DO PORTO

Rua Alfredo Allen, 208
4200-135 Porto
Portugal
+351 220 408 800
info@i3s.up.pt
www.i3s.up.pt

Version: Postprint (identical content as published paper) This is a self-archived document from i3S – Instituto de Investigação e Inovação em Saúde in the University of Porto Open Repository For Open Access to more of our publications, please visit <http://repositorio-aberto.up.pt/>

FIGURES




Figure 1. Schematic representation of Lewis antigens. Type 1 chains (panel A) are characterized by the Gal β 1,3 linkage, while type 2 chains (panel B) display a Gal β 1,4 linkage. (Panel A) Addition of a fucose to terminal galactose on type 1 structures leads to H - type 1 structure, which can be further modified with a fucose on the GlcNAc residue resulting in Le^b antigen. The type 1 backbone may be just fucosylated on the GlcNAc residue leading to Le^a antigen biosynthesis. The action of a sialyltransferase towards the type 1 chain can lead to the biosynthesis of the sLe^a antigen. (Panel B) the biosynthesis of type 2 based Lewis antigens depends on the addition of the same glycan units but with different linkages, originating the isomers Le^x , Le^y and sLe^x antigens. Addition of a fucose to terminal galactose on type 2 structures leads to H - type 2, which can be further modified with a fucose on the GlcNAc residue resulting in Le^y antigen. The type 2 backbone may be just fucosylated on the GlcNAc residue leading to Le^x antigen biosynthesis. The action of a sialyltransferase toward the type 2 chain can lead to the biosynthesis of the sLe^x antigen. (Panel C) Schematic representation of canine secretory alloantigens (CSA) polymorphisms (adapted from Oriol et al. 30).




Figure 2. Expression of Lewis antigens and Dolichus biflorus agglutinin (DBA) lectin staining in gastric mucosa: (A) and (B) Normal canine gastric mucosa, IHC Lea, 100x and 40x respectively. There was no Lea antigen expression in both body and antrum. (D) and (E) Normal canine gastric mucosa, IHC Leb, 100x and 40x respectively. There was no Leb antigen expression in both body and antrum. (G) and (H) Normal canine gastric mucosa, IHC sLea, 100x and 40x, respectively. There was no sLea antigen expression in both body and antrum. (J) Normal canine gastric body, IHC Lex, 100x. Multifocal and moderate cytoplasm expression of Lex in the superficial foveolar epithelium. (K) Normal canine gastric antrum, IHC Lex, 40x. Generalized and moderate staining of the superficial foveolar epithelium. (M) Normal canine gastric body, IHC Ley, 100x. Generalized and intense immunostaining of superficial foveolar epithelium and parietal cells. (N) Normal canine gastric antrum, IHC Ley, 40x. Intense and generalized expression of Ley in foveolar epithelium and gastric pits. (P) Normal canine gastric body, IHC sLex, 100x. Multifocal and intense immunostaining of superficial foveolar epithelium. (Q) Normal canine gastric antrum, IHC sLex, 40x. Sparse and weak immunostaining of the foveolar superficial epithelial cells and in cells of the pyloric glands. (C), (F), (I), (L), (O), (R) Expression of Lewis antigens in human gastric mucosa with intestinal metaplasia used as positive controls, 100x. (S) Normal canine gastric body, Dolichus biflorus agglutinin (DBA), 100x. Generalized and intense staining of superficial foveolar epithelium and parietal cells. (T) Normal canine gastric antrum, DBA, 100x. Intense and generalized staining of foveolar epithelium and gastric pits, contrasting with lesser intensity in deeper pyloric glands. (U) DBA staining in normal canine intestinal mucosa used as positive control, DBA, 200x.




Figure 3. The panel shows the various morphologies of fluorescein - labeled *Helicobacter* spp. strains: (A) *H. pylori* 17875/Leb (B) *H. pylori* 17875babA1A2 (C) *H. heilmannii* s.s. ASB1.4 (D) *H. felis*CS1 (E) *H. bizzozeronii*R1051 (F) *H. salomonis*R1053. Magnification 1000 × .




Figure 4. Evaluation of Helicobacter adherence to the body region of the canine gastric mucosa. Adhesion of fluorescein - labeled *H. pylori* model strains: (A) 17875/Leb and (B) 17875babA1A2; Adhesion of fluorescein - labeled Non - *H. pylori* Helicobacters (NHPH) strains: (C) ASB1.4, (D) CS1, (E) R1051 and (F) R1053. Magnification 200x. (G) Quantification of adherence of the different Helicobacter spp. strains. Each symbol represents the mean value obtained by evaluating at least five different fields within the foveolar or the glandular regions, as defined in 2. The differences of the mean values between the groups were statistically significant using one - way ANOVA ($p < .0001$).




Figure 5. Evaluation of *Helicobacter* adherence to the antral region of the canine gastric mucosa. Adhesion of fluorescein - labeled *H. pylori* model strains: (A) 17875/Leb and (B) 17875babA1A2; Adhesion of fluorescein - labeled Non - *H. pylori* *Helicobacters* (NHPHs) strains: (C) ASB1.4, (D) CS1, (E) R1051 and (F) R1053. Magnification 200x. (G) Quantification of adhesion of the different *Helicobacter* spp. strains. Each symbol represents the mean value obtained by evaluating at least five different fields within the foveolar or glandular regions, as defined in 2. The differences of the mean values between the groups were statistically significant using one - way ANOVA ($p < .0001$).

TABLES

Table 1 Antibodies used in immunohistochemistry

Monoclonal antibody	Antigen	Supplier/References	Dilution
CA3F4	Le ^a	Young et al. [55]	1 : 5
BG6	Le ^b	Signet [®] (Dedham, MA, USA)	1 : 50
CA19-9	Sialyl-Le ^a	Santa Cruz [®] (Santa Cruz, CA, USA)	1 : 500
SH1	Le ^x	Fukushi et al. [56]	1 : 5
IM93	Sialyl-Le ^x	Calbiochem [®] (Darmstadt, Germany)	1 : 60
AH6	Le ^y	Abe et al. [57]	1 : 10

Table 2 Terminal α -GalNAc (DBA), type 1 and type 2 Lewis antigens expression in canine gastric mucosa

		DBA	Type 1			Type 2		
			Le ^a	Le ^b	sLe ^a	Le ^x	Le ^y	sLe ^x
Body (n = 8)	Positive immunostaining	8/8	1/8	0/8	0/8	5/8	8/8	4/8
	Immunostaining distribution							
	Sparse	0	1*	0	0	0	0	0
	Focal	0	0	0	0	0	0	0
	Multifocal	0	0	0	0	5 ⁺⁺⁺	4 ⁺⁺⁺	4 ⁺⁺⁺
	Generalized	8 ⁺⁺⁺	0	0	0	0	4 ⁺⁺⁺	0
Antrum (n = 6)	Positive immunostaining	6/6	0/6	0/6	0/6	6/6	6/6	1/6
	Immunostaining distribution							
	Sparse	0	0	0	0	0	0	1 ⁺⁺
	Focal	0	0	0	0	0	0	0
	Multifocal	0	0	0	0	4 ⁺⁺⁺	0	0
	Generalized	6 ⁺⁺⁺	0	0	0	2 ⁺⁺⁺	6 ⁺⁺⁺	0

DBA, *Dolichus biflorus* agglutinin.

Intensity variation observed: +weak; ++moderate; +++intense.