



Original Research

A preliminary study on the role of personal history of infectious and parasitic diseases on self-reported health across countries

Gerit Pfuhl^{a,*}, Filipe Prazeres^{b,c,d,do}, Marta Kowal^e, Toivo Aavik^f, Beatriz Abad-Villaverde^g, Reza Afhami^h, Leonardo Aguilarⁱ, Grace Akello^j, Laith Al-Shawaf^{k,l,m}, Jan Antfolkⁿ, Chiemezie S. Atama^o, Derya Atamturk Duyar^p, Roberto Baiocco^q, Sercan Balım^r, Carlota Batres^s, Yakhlef Belkacem^t, Théo Besson^u, Adam Bode^v, Merve Boğa^w, Jordane Boudesseul^{x,y}, Mahmoud Boussena^z, Hamdaoui Brahim^{aa}, Nana Burduli^{ab}, Ali R. Can^{ac}, Hakan Cetinkaya^{ad}, Antonio Chirumbolo^{ae}, Dimitri Chubinidze^{af}, Clément Cornec^{ag}, Bojana M. Dinić^{ah}, Seda Dural^{ai}, Izzet Duyar^p, Samuel O. Ebimngbo^{aj,ak}, Edgardo Etchezahar^{al}, Peter Fedor^{am}, Tomasz Frackowiak^{an}, David A. Frederick^{ao}, Katarzyna Galasinska^{ap}, Felipe E. García^{aq}, Talia Gomez Yepes^{al}, Dmitry Grigoryev^{ar}, Farida Guemaz^{as}, Ivana Hromatko^{at}, Gözde Ikizer^{au}, Steve M. J. Janssen^{av}, Julia A. Kamburidis^{aw}, Tina Kavčič^{ax,ay}, Nicolas Kervyn^{az}, Farah Khan^{ba}, Aleksander Kobylarek^{bb}, Mehmet Koyuncu^{bc}, Yoshihiko Kunisato^{bd}, David Lacko^{be}, Miguel Landa-Blanco^{bf}, Linda H. Lidborg^{bg}, Samuel Lins^{bh}, Tetyana Mandzyk^{bi}, Silvia Mari^{bj}, Tiago A. Marot^{bk}, Martha Martinez-Banfi^{bl}, Alan D. A. Mattiassi^{bm}, Marlon Mayorga-Lascano^{bn}, Moises Mebarak^{bo}, Norbert Meskó^{bp}, Maria Rosa Miccoli^{bq}, Vita Mikuličiūtė^{br}, Taciona L. Milfont^{bs}, Katarina Mišetić^{bt}, Mara Morelli^{bu}, Jean C. Natividade^{bv}, Izuchukwu L. G. Ndukaihe^{bw}, Felipe Novaes^{bx}, Salma S. Omar^{by}, Mohd Sofian Omar Fauzee^{bz}, Tobias Otterbring^{ca}, Barış Özener^p, Simon Ozer^{cb}, Ju Hee Park^{cc}, Irena Pavela Banai^{cd}, Farid Pazhoohi^{ce}, Mariia Perun^{bi}, Martin Pírko^{cf}, Ekaterine Pirtskhalava^{cg}, Katarzyna Pisanski^{ag,an,ch}, Nejc Plohl^{ci}, Koen Ponnet^{cj}, Pavol Prokop^{ck,cl}, Matheus F. F. Ribeiro^{cm}, Frederico Rosário^{cn}, Ayşegül Şahin^p, Fatima Zahra Sahli^{co}, Dušana Šakan^{cp}, Oksana Senyk^{cq,cr}, Henrik Siepelmeyer^{ca}, Diana Ribeiro da Silva^{cs}, Sangeeta Singh^{ct}, Çağlar Solak^{cu}, Sinem Söylemez^{cu}, Anna Studzinska^{cv}, Chee-Seng Tan^{cw}, Gulmira T. Topanova^{cx}, Merve Topcu Bulut^{cy}, Ezgi Toplu-Demirtaş^{cz}, Bastien Trémolière^{da,db}, Singha Tulyakul^{dc}, Joaquín Ungaretti^{dd}, Jaroslava V. Valentova^{de}, Marco A.C. Varella^{de}, Mona Vintila^{df}, Tatiana Volkodav^{dg}, Anna Włodarczyk^{dh}, Yao-Yuan Yeh^{di}, Gyesook Yoo^{dj}, Oulmann Zerhouni^{dk,dl}, Marcos Zumárraga-Espinosa^{dm}, Maja Zupančič^{dn}, Piotr Sorokowski^{an}

^a Department of Psychology, NTNU, Norway

^b Faculty of Health Sciences, University of Beira Interior, Portugal

^c Family Health Unit Beira Ria, Portugal

^d RISE-Health, MEDCIDS, Faculty of Medicine of the University of Porto, Portugal

* Corresponding author.

E-mail address: gerit.pfuhl@ntnu.no (G. Pfuhl).

<https://doi.org/10.1016/j.puhe.2025.02.030>

Received 12 August 2024; Received in revised form 29 January 2025; Accepted 20 February 2025

Available online 23 March 2025

0033-3506/© 2025 The Authors. Published by Elsevier Ltd on behalf of The Royal Society for Public Health. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

- ^e IDN Being Human Lab, Institute of Psychology, University of Wrocław, Poland
- ^f Institute of Psychology, University of Tartu, Estonia
- ^g Faculty of Education and Humanities, Universidad Nacional Pedro Henríquez Ureña, Dominican Republic
- ^h Department of Art Studies, Tarbiat Modares University, Iran
- ⁱ School of Psychology, Central University of Venezuela, Venezuela
- ^j Faculty of Medicine, Department of Mental Health, Gulu University, Uganda
- ^k Department of Psychology, University of Colorado, United States
- ^l Center for Cognitive Archeology, University of Colorado, United States
- ^m Institute for Advanced Study in Toulouse, France
- ⁿ Faculty of Arts, Psychology, and Theology, Åbo Akademi University, Finland
- ^o Department of Sociology & Anthropology, University of Nigeria, Nigeria
- ^p Department of Anthropology, İstanbul University, Turkey
- ^q Department of Developmental and Social Psychology, Sapienza University of Rome, Italy
- ^r Department of Psychology, Bursa Technical University, Turkey
- ^s Department of Psychology, Franklin and Marshall College, United States
- ^t Ecole Normale Supérieure, Assia DJEBBAR de Constantine Algérie, Algeria
- ^u Laboratoire de Psychologie Sociale, Université Paris Cité, France
- ^v School of Archaeology and Anthropology, ANU College of Arts and Social Sciences, Australian National University, Australia
- ^w Department of Psychology, Bursu Uludag University, Turkey
- ^x Laboratoire Parisien de Psychologie Sociale, Psychologie Sociale des Comportements et des Cognitions, Univ. Paris Nanterre, France
- ^y Instituto de Investigación Científica, Universidad de Lima, Peru
- ^z Department of Psychology and Educational Sciences, University of Mohamed Lamine Debaghine, Setif2, Algeria
- ^{aa} Département de Sociologie, Ibn Tofail University, Morocco
- ^{ab} Department of Psychology, University of Georgia, Georgia
- ^{ac} Department of Anthropology, Hatay Mustafa Kemal University, Turkey
- ^{ad} Department of Psychology, Yaşar University, Turkey
- ^{ae} Department of Psychology, Sapienza University of Rome, Italy
- ^{af} Psychological Medicine, King's College London, United Kingdom
- ^{ag} ENES Bioacoustics Research Lab, Jean Monnet University of Saint Etienne, France
- ^{ah} Department of Psychology, Faculty of Philosophy, University of Novi Sad, Serbia
- ^{ai} Department of Psychology, İzmir University of Economics, Turkey
- ^{aj} Department of Social Work, University of Nigeria, Nigeria
- ^{ak} Faculty of Catholic Theology, Julius-Maximilians University, Germany
- ^{al} Faculty of Teacher Training and Education, Autonomous University of Madrid, Spain
- ^{am} Department of Environmental Ecology, Comenius University, Faculty of Natural Sciences, Slovakia
- ^{an} Institute of Psychology, University of Wrocław, Poland
- ^{ao} Crean College of Health and Behavioral Sciences, Chapman University, United States
- ^{ap} Department of Psychology, SWPS University, Poland
- ^{aq} Departamento de Psiquiatría y Salud Mental, Universidad de Concepción, Chile
- ^{ar} Center for Sociocultural Research, HSE University, Russia
- ^{as} Department of Psychology and Educational Sciences, University of Setif2, Algeria
- ^{at} Department of Psychology, University of Zagreb, Croatia
- ^{au} Department of Psychology, TOBB University of Economics and Technology, Turkey
- ^{av} School of Psychology, University of Nottingham Malaysia, Malaysia
- ^{aw} Department of Social, Organizational, Clinical and Pedagogical Psychology, Sofia University “St.Kliment Ohridsky”, Bulgaria
- ^{ax} Department of Psychology, Faculty of Arts, University of Ljubljana, Slovenia
- ^{ay} Faculty of Health Sciences, University of Ljubljana, Slovenia
- ^{az} Louvain School of Management, Université Catholique de Louvain, Belgium
- ^{ba} Department of Education, Women University Mardan KP, Pakistan
- ^{bb} Institute of Pedagogy, University of Wrocław, Poland
- ^{bc} Department of Psychology, Izmir Bakircay University, Turkey
- ^{bd} Department of Psychology, Senshu University, Japan
- ^{be} Interdisciplinary Research Team on Internet and Society, Faculty of Social Studies, Masaryk University, Brno, Czech Republic
- ^{bf} School of Psychological Sciences, National Autonomous University of Honduras, Honduras
- ^{bg} Department of Psychology, Durham University, Durham, United Kingdom
- ^{bh} Department of Psychology, University of Porto, Portugal
- ^{bi} Department of Psychology, Ivan Franko National University of Lviv, Ukraine
- ^{bj} Department of Psychology, University of Milano-Bicocca, Italy
- ^{bk} Department of Administration, Fundação Getúlio Vargas, Brazil
- ^{bl} Facultad de Ciencias Jurídicas y Sociales, Centro de Investigaciones en Ciencias de la Vida-CICV, Universidad Simón Bolívar, Colombia
- ^{bm} Department of Education, Literatures, Intercultural Studies, Languages and Psychology, University of Florence, Italy
- ^{bn} Escuela de Psicología, Pontificia Universidad Católica del Ecuador-Ambato, Ecuador
- ^{bo} Department of Psychology, Universidad del Norte, Colombia
- ^{bp} Institute of Psychology, Department of Cognitive and Evolutionary Psychology, University of Pécs, Hungary
- ^{bq} Department of Psychology, University of Konstanz, Germany
- ^{br} Department of Psychology, Vilnius University, Lithuania
- ^{bs} School of Psychological and Social Sciences, University of Waikato, Tauranga, New Zealand
- ^{bt} Department of Psychology, University of Sarajevo, Bosnia and Herzegovina
- ^{bu} Department of Dynamic and Clinical Psychology, and Health Studies, Sapienza University of Rome, Italy
- ^{bv} Graduate Program of Psychology, Pontifical Catholic University of Campinas, Brazil
- ^{bw} Department of Psychology, Alex Ekwueme Federal University, Nigeria
- ^{bx} Department of Psychology, Pontifical Catholic University of Rio de Janeiro, Brazil
- ^{by} Department of Dermatology, Venereology & Andrology, Alexandria University, Egypt
- ^{bz} Sport Academy UPM, Faculty of Education and Liberal Arts, INTI International University, Malaysia
- ^{ca} Department of Management and Innovation, University of Agder, Norway
- ^{cb} Department of Psychology and Behavioural Sciences, Aarhus University, Denmark
- ^{cc} Department of Child and Family Studies, Yonsei University, Republic of Korea
- ^{cd} Department of Psychology, Faculty of Humanities and Social Sciences, J. J. Strossmayer University of Osijek, Croatia
- ^{ce} School of Psychology, University of Plymouth, United Kingdom
- ^{cf} Institute of Lifelong Learning, Mendel University in Brno, Czech Republic

- ^{c8} Department of Psychology, Ivane Javakhishvili Tbilisi State University, Georgia
- ^{ch} French National Centre for Scientific Research (CNRS), Language Dynamics Laboratory, University of Lyon, France
- ^{ci} Department of Psychology, University of Maribor, Slovenia
- ^{cj} Faculty of Social Sciences, imec-mict-Ghent University, Belgium
- ^{ck} Department of Environmental Ecology and Landscape Management, Faculty of Natural Sciences, Comenius University, Slovakia
- ^{cl} Department of Animal Ecology, Slovak Academy of Sciences, Slovakia
- ^{cm} Department of Psychology, University of Uberaba, Brazil
- ^{cn} USF Tondela, ACES Dão Lafões, Viseu, Portugal
- ^{co} IMS, Ibn Tofail University, Morocco
- ^{cp} Department of Psychology, Faculty of Legal and Business Studies Dr Lazar Vrkatić, Union University, Serbia
- ^{cq} Department of Humanities and Social Sciences, WSB Merito University in Gdansk, Poland
- ^{cr} Department of Psychology and Psychotherapy, Ukrainian Catholic University, Ukraine
- ^{cs} Center for Research in Neuropsychology and Cognitive Behavioral Intervention (CINEICC), Faculty of Psychology and Education Sciences, University of Coimbra, Portugal
- ^{ct} Department of Strategy and Leadership, University of Agder, Norway
- ^{cw} Department of Psychology, Manisa Celal Bayar University, Turkey
- ^{cx} Humanities Department, Icam School of Engineering, Toulouse Campus, France
- ^{cw} School of Psychology, Wenzhou-Kean University, China
- ^{cx} Department of Theoretical and Practical Psychology, Kazakh National Women's Pedagogical University, Kazakhstan
- ^{c7} Clinical Addiction Research Unit, Lund University, Sweden
- ^{cz} Department of Psychological Counseling and Guidance, MEF University, Turkey
- ^{da} Department of Psychology, University of Toulouse Jean Jaurès, France
- ^{db} Department of Psychology, Université du Québec à Trois-Rivières, Canada
- ^{dc} Department of Health and Physical Education, Thaksin University, Thailand
- ^{dd} Faculty of Education, Universidad Internacional de Valencia, Spain
- ^{de} Department of Experimental Psychology, Institute of Psychology, University of Sao Paulo, Brazil
- ^{df} Department of Psychology, West University of Timisoara, Romania
- ^{dg} Department of Pedagogy and Psychology, Kuban State University, Russia
- ^{dh} Escuela de Psicología, Universidad Católica del Norte, Chile
- ^{di} Department of International Studies & Modern Languages, University of St. Thomas, Houston, United States
- ^{dj} Department of Child and Family Studies, Kyung Hee University, Republic of Korea
- ^{dk} Centre de Recherche sur les Fonctionnements et Dysfonctionnements Psychologiques, University of Rouen Normandy, France
- ^{dl} Laboratoire Parisien de Psychologie Sociale, Université Paris Nanterre, France
- ^{dm} Universidad Politécnica Salesiana, Ecuador
- ^{dn} Faculty of Arts, University of Ljubljana, Slovenia

ARTICLE INFO

Keywords:

Mental health
Global health
Multiverse analysis
Communicable diseases

ABSTRACT

Objectives: Infectious diseases are often associated with decline in quality of life. The aim of this study is to analyze the relationship between personal history of communicable, i.e., infectious and parasitic diseases and self-rated health.

Study design: Secondary analysis of a large dataset multi-country observational study.

Methods: We used a four-pronged analysis approach to investigate whether personal history of infectious and parasitic diseases is related to self-reported health, measured with a single item.

Results: Three of the four analyses found a small positive effect on self-reported health among those reporting a history of pathogen exposure. The meta-analysis found no support but large heterogeneity that was not reduced by two classifications of countries.

Conclusion: Personal history of infectious and parasitic diseases does not reduce self-reported health across a global sample.

1. Introduction

Numerous people live with chronic diseases and conditions, as life expectancy is increasing, and the numbers of older persons also rise.^{1,2} For example, between 1985 and 2005, the prevalence of chronic diseases doubled.³ In 2017, chronic diseases represented more than 60 % of the global burden of disease.⁴ Nonetheless, when focusing on low-income countries, communicable (infectious and parasitic) diseases still contribute to the high disease burden.⁴

In 2019 alone, tuberculosis was responsible for 1.18 million deaths, malaria for 643,000 deaths, typhoid fever for 110,000 deaths, dengue for 36,100 deaths, schistosomiasis for 11,500 deaths, leishmaniasis for 5,710 deaths, and trypanosomiasis for 1,360 deaths.⁵ It is widely agreed that these numbers will – if at all – only slowly decrease given drug resistances,⁶ global climate change⁷ and the limitations of health education programs (e.g., personal hygiene) and sanitation.⁸ Besides the loss of human lives, infectious diseases have other consequences, such as

decline in quality of life, expenses, medication, incapacity to work, or strain on caregivers.²

Higher infectious disease rates and their impact on mortality and morbidity are more commonly found in individuals with lower social status.⁹ The prevalence of infectious diseases has been linked to cross-cultural differences between regions,^{10,11} which span from gastronomic variations¹² to distinctions of personality traits.¹³ Of relevance for this secondary analysis, poor self-reported health was found in individuals not only with chronic diseases^{14,15} but also among those with infectious diseases.¹⁶

Self-reported health is a general health indicator widely used in public health studies.^{9,17} Self-reported health usually consists of a single question asking people to rate their general health. Despite its simplicity, this single self-report item is considered a relatively reliable tool for assessing general health and has been used worldwide in an array of research settings and conditions.^{18,19} Its widespread use as an index of health may be tied to lessening the burden for the respondent, lower costs, and easier interpretation compared to multi-item measures.¹⁷

Across European countries, self-reported health status is positively

^{do} Gerit Pfuhl and Filipe Prazeres share the first co-authorship.

associated with life satisfaction²⁰. A meta-analysis of studies assessing self-reported health with a single item and all-cause mortality as outcome found that persons with poor self-reported health, although subjective, had a 2-fold higher mortality risk compared with persons reporting excellent self-rated health.²¹ This aligns well with a cohort study from Sweden, finding that self-reported health correlates negatively with more objective measures, such as mortality risk (hazard ratio .89), use of social insurance facilities (disability pension, hazard ratio .77), and utilization of health care services (hazard ratio .96).²²

Some authors have stated that even though communicable diseases have an important public health weight, they are known as neglected tropical diseases because they are not frequently researched^{23,24} and vaccine development has been slow.^{25–29} For instances, there is a lack of studies that consider both distinct adult age groups and individuals from different countries with infectious diseases and multiple parasitic infections.³⁰ The aim of this study, therefore, is to analyze the relationship between personal history of communicable, i.e., infectious and parasitic, diseases and self-rated health, through a secondary analysis of a large, cross-cultural, and cross-cohort database.³¹

2. Methods

The present study is a cross-sectional secondary analysis of a dataset from 187 countries. Data were collected through an online questionnaire administered between April and August 2021 in 45 different languages, with two countries (Algeria and Morocco) employing written questionnaires. Participants were recruited from diverse sample pools, considering factors such as sex, age, residence (including both small and large cities), and education. The survey link was shared on various social media platforms, while approximately 6 % of the data were collected through outsourcing platforms. Detailed information on data collection methodology can be found elsewhere.^{31,32} The original collection of data was approved by the Institutional Review Board (IRB) at the Institute of Psychology, University of Wrocław. All team members adhered to the principles of the Declaration of Helsinki³³ and the ethical guidelines of their respective IRBs, either based on the ethical approval of the Principal Investigator's IRB or their local IRB. Prior to participating in the survey, all participants provided informed consent, and the anonymity of their data was ensured. 117,289 participants (women: 71,325, men: 34,862) aged 18–90 years ($M = 30.35$, $SD = 12.55$) took part.

For the present study, participants that did not answer the self-perceived health item were excluded ($N = 26,486$). Participants not indicating their sex ($n = 9,527$, 8.1 %) or age ($n = 9,754$, 8.1 %) were removed. Also, given the low prevalence of non-binary participants ($n = 1,089$, .9 %) in the dataset (this option was not available in all countries during data collection), respondents indicating non-binary as gender were excluded from analysis. Participants not responding to any of the nine pathogen items were also removed ($n = 19,232$). This reduced the original number of observations from 117,289 to 87,673 (multiple causes possible). See the supplementary material for a comparison among responders and non-responders.

Note, the final number of participants included in the analysis varies for specific analysis (e.g., removing those raised in a different country than currently living, $N = 6,001$).

Self-perceived health was assessed by an 11-point Likert scale to the question: How healthy are you? (1 = Extremely unhealthy; 11 = Extremely healthy). This question has been used earlier in health, social, and psychology surveys.^{34–37}

Demographic variables were sex (female, male), age, country-raised, and country lived.

Personal history of infectious and parasitic diseases was assessed by the nine-item Pathogen Prevalence Index¹⁰ through the question “Have you ever contracted (been sick with) any of the following diseases?” (response alternatives were: Never (0), Once (1), More than once (2)) concerning each of nine infectious and parasitic diseases (see [supplementary material](#)). Multiple answers were possible, 269 participants

indicated to have been exposed to all pathogens at least once, and 30 participants indicated that they had all nine diseases multiple times. Overall, 10.06 % of the participants reported an exposure to any of the nine pathogens at least once (pathogen(s) once: $n = 6,345$, one pathogen multiple times: $n = 2,480$, more than one pathogen/disease: $n = 2103$. $N = 269$ indicated to have contracted all diseases at least once, whereof $n = 166$ live in the USA, see [supplementary material, table 1](#)). The nine pathogens significantly differ with respect to self-rated health ($F(9, 87671) = 153.33$, $p < .0001$, Pillai's trace = .015, with dengue, filariae and typhus not yielding any difference to those not exposed to any pathogen whereas participants who have been exposed to any of the other six pathogens once report higher health, see [supplementary material, Fig. S1](#)).

We had no prediction about differences by type of pathogen/disease, especially since their occurrence naturally varies by country. We therefore coded having had any of the pathogens at least once as a prevalence. It is worth noting that for participants who selected ‘More than once’, we do not know whether this “multiple times” refers to twice, thrice, or any other multiple exposure. We therefore used three groups, never exposed, exposed to one pathogen only once, exposed to either the same pathogen multiple times or exposed more than one pathogen. Thus, multiple exposure refers to either the same pathogen more than once or having contracted an infection with two or more pathogens during one's lifetime.

Regarding data analysis, we performed a four-pronged approach.³⁸ We firstly investigated the relationship between pathogen prevalence and self-rated health with linear regression, including age and sex as additional predictors. We secondly used linear mixed modelling nesting participants within countries (random effect) and pathogen prevalence, age and gender as fixed effects. For this analysis we excluded participants that reported living in a different country than born ($n = 6,001$) reducing the sample size to $N = 81,672$. Thirdly, we performed propensity scoring to limit the selection bias in our retrospective observational study.³⁹ Propensity scoring will pseudo-randomize participants who reported a prevalence with a participant who reported no prevalence. Since propensity scores analyze effects at the sample level, not the individual level as regression analysis does, we excluded countries with less than 30 responses;³⁹ this exclusion left 90 countries (from 187 countries). Propensity scoring by age, sex and country (using exact matching for country and sex, allowing nearest for age to increase the number of matches) yielded 5,212 participants per group ($N = 10,424$).

Fourthly, we performed a random effect meta-analysis. For this analysis we removed all countries with less than 30 respondents, and the countries also needed to have at least ten respondents reporting a prevalence. This ensured that the statistics per country were reliable (mean similarity difference⁴⁰). The meta-analysis is based on $n = 79$, 563.

We followed this up with two subgroup analyses, one by the UN economic categorization of countries into developed, developing, in transition and least developed countries, and another one by categorizing countries into temperate and tropical.

3. Results

Self-reported health on a scale from 1 to 11, with higher values representing better health, was $M = 7.43$ ($SD = 2.08$).

Among participants without pathogen prevalence self-perceived health rating was generally good, $M = 7.39$, $SD = 2.05$, 95 % CI [7.38, 7.40], participants with one pathogen prevalence reported a slightly higher health, $M = 7.65$, $SD = 2.14$, 95 % CI [7.59; 7.71] and participants with multiple exposures to either the same or different pathogens reported the highest health score, $M = 8.35$, $SD = 2.26$, 95 % CI [8.29; 8.42]. This difference was statistically significant, $F(2, 87670) = 414.7$, $p < .001$, $R^2_{adj} = .009$. Post-hoc tests yielded a significant difference between the “never being exposed” group and the “once exposed” group ($t = -8.19$, $p < .001$, $\beta = -.261$) and the “never being

exposed” group and the “multiple times” group ($t = -27.98, p < .001, \beta = -.963$), and between the “once exposed” and “multiple times” group ($t = -15.35, p = .262, \beta = -.702$). Note, both the “once exposed” and the “multiple times” group reported higher health. Given the low proportion of participants with exposure we decided to pool the two groups and contrast them with the never exposed group in all subsequent analyses. Applying this dichotomy we find that self-perceived health was statistically lower among those reporting no history of infectious and parasitic diseases ($M = 7.39, SD = 2.05, 95\% \text{ CI } [7.38; 7.4]$) than among those reporting pathogen exposure at least once ($M = 7.97, SD = 2.26, 95\% \text{ CI } [7.93; 8.02]$), Welch’s $t(9663.4) = -22.71, p < .001$, Cohen’s $d = .28$.

Supplementary Table 2 provides the self-rated health per country and separately for participants reporting an exposure or reporting no exposure to any of the nine diseases.

3.1. Regression analyses

A simple linear regression for the complete dataset ($N = 87,672$) with self-rated health as outcome and sex, age, and personal history of infectious and parasitic diseases as predictors, explained 1.1 % of the variance in self-perceived health ($F(3, 87,669) = 333.5, R_{\text{adj}}^2 = .011, p < .001$). Regarding the predictors; a) for every decade reported self-rated health reduced with .11 points on the scale, $t(87,669) = -20.125, \beta = -.11, p < .001$; b) for reporting exposure at least once to any pathogen listed increased self-rated health by over half a point, $t(87,669) = 23.4, \beta = .56, p < .001$, and c) being a woman statistically significantly reduced self-rated health (for men: $M = 7.47, SD = 2.14$, for women: $M = 7.43, SD = 2.04$), $t(87,669) = 2.35, \beta = .035, p = .019$. Pathogen prevalence was not detrimental to self-perceived health. We next conducted a sensitivity analysis using being female as a benchmark covariate, and pathogen prevalence (binary) as treatment. In explaining pathogen prevalence and self-rated health we investigate the maximum strength of a confounder once, twice, or three times as strong as female. We used the *sensemkr* package.⁴¹ The robustness value for testing the null hypothesis was 6.98 %, i.e., unobserved confounders not explaining at least 6.98 % of the residual variance are not sufficiently strong to change the association between pathogen prevalence and self-rated health (see supplementary Material). Unobserved confounders would need to explain at least .6 % of the residual variance of the treatment to fully explain away the observed effect. One possible confounder is country; hence we next tested a linear mixed model.

A linear mixed model, nesting participants within a country, age and sex as covariates, yielded results similar to the results of the simple linear regression, that is, self-rated health decreased with age ($t(81,390) = -7.19, p < .001, \beta = -.005$), women reported a lower self-rated health ($t(81,660) = 3.86, p = .0001, \beta = .059$), and exposure at least once to any pathogen increased self-rated health ($t(76,080) = 3.53, p = .0004, \beta = .011$). Note, the regression models explained very little of the variance for self-rated health. We next assessed robustness by running a model after excluding participants from countries with fewer than 30 respondents. This analysis replicated the above findings, i.e., self-rated health decreased with age ($t(81,110) = -7.25, p < .001, \beta = -.005$), women reported a lower self-rated health ($t(81,420) = 3.95, p < .001, \beta = .06$), and exposure at least once to any pathogen increased self-rated health ($t(75,930) = 3.66, p = .0003, \beta = .011$). An analysis of deviance yielded that in both models all three predictors were significant and similar in size (see supplementary material).

3.2. Propensity scoring

We performed propensity scoring with the *match-it* package in R.⁴² Country and sex were set as exact, age was set to nearest. The distance was calculated using generalized linear models with a logit link function, and a 1:1 approach used. There were 10,424 matched cases. Self-rated health was slightly higher in the pathogen group ($M = 7.52$,

$SD = 2.19$) than the non-pathogen group ($M = 7.4, SD = 2.14$). Welch’s t -test yielded a significant difference, $t(10,361) = -2.79, p = .005$, Cohen’s $d = .055$. Overall, participants with pathogen prevalence reported on average a slightly better perceived health (Supplementary Fig. S2), but note the negligible effect size.

3.3. Random-effect meta-analysis

For each country with more than 30 participants and at least 10 participants (55 countries, $n = 71,397$) that reported a personal history of infectious and parasitic diseases, we calculated the weighted effect size for the relation between self-rated health and personal history of infectious and parasitic diseases. We used a random effect model in the *meta* package. Across the 55 countries, we found no evidence of a significant difference in self-rated health between participants with or without a personal history of infectious and parasitic diseases (estimated difference = $-.006, 95\% \text{ CI } [-.008; .067], p = .876$). There was considerable heterogeneity ($I^2 = 82.9\%, 95\% \text{ CI } [78.4\%; 86.5\%]$). Fig. 1 presents the forest plot. To address the heterogeneity of our sample, we categorized countries either by their economy—as countries with better economy tend to have a better health system, easily offering adequate treatment for their citizen—and by their climate—as this might capture the disease load with tropical countries having a higher load than temperate countries.

A subgroup analysis by UN categorization by economy reduced heterogeneity in the least developed group (38.3 %) but not in the developing and developed group (81.1 % and 85.6 %, respectively). A subgroup analysis classifying countries into temperate and tropical reduced heterogeneity for tropical (49.2 %) but not for temperate regions (88.1 %). In both subgroup analyses there were no significant differences in the weighted effect size (see Supplementary Figs. 3 and 4).

4. Discussion

The present study was designed to investigate the impact of personal history of infectious and parasitic diseases on self-rated health, and its variation across age, sex, and countries. Using four different statistical approaches, we found no detrimental reduction on self-rated health among those exposed at least once to one of the nine pathogens we studied. Three analyses showed a lower rating of health among those never exposed to any pathogen, but with negligible effect sizes. The random-effects meta-analysis found no difference in self-reported health across 55 countries. However, there was large heterogeneity, particularly among developed and developing countries in temperate regions. The heterogeneity analyses (common effects model) suggest a beneficial relationship between pathogen prevalence and self-rated health in developed countries, and in temperate but not tropical regions, likely due to lower prevalence and better health and education systems.

The four-pronged approach started from treating the sample as being a random draw from the world population (linear regression), to acknowledging differences by country (linear mixed model, propensity scoring, meta-analysis). The analyses complement each other, and importantly, do not find a systematic reduction in self-perceived health among those having been exposed to pathogens.

A possible explanation for this might be that in the face of recurring acute health challenges, individuals develop better coping mechanisms, including palliative coping strategies,⁴³ that may lead to an improved perception of their health over time, and embracing the challenges that life provides.⁴⁴ Experiencing infectious and parasitic diseases may also led to increased appreciation of health and therefore higher satisfaction with one’s health.⁴⁵ On the other hand, lower self-perceived health is typically associated with the presence of chronic health conditions in individuals,⁴⁶ suggesting that participants in this study did not suffer from long-term consequences of the acute infection. A conceptual model⁴⁷ considers a comparison to other people’s health as a factor for how one rates one’s own health. As a mechanism of resilience,




Fig. 1. Forest plot random-effect model.

downward comparison but not upward comparison increases subjective well-being.⁴⁸ Future studies could examine if, indeed, those with exposure to pathogen(s) engaged more in downward comparison whereas those without exposure engaged more in upward comparison.

Older age^{49,50} and being a woman⁵⁰ are known predictors of poor self-perceived health and this aligns with the findings of the present research, supporting its trustworthiness.

Interpreting the findings regarding the association between personal history of infectious and parasitic diseases and self-reported health is challenging due to unaccounted interactions and confounding variables in the analysis. These factors, such as mental health problems, cultural beliefs, values, life satisfaction,²⁰ demographic, geographical and

ecological factors, differences in the distribution of the considered diseases in the participant countries, norms within the sampled countries, and disparities in access to healthcare services, may all contribute to the self-reported health levels, but we are not able to make any further conclusion on this complex set of interactions in this secondary analysis study. We note though, that subgroup analyses did not remove heterogeneity for all subgroups.

This study has some important limitations. Because it is a secondary analysis, the study is limited to the variables and data already collected, which may restrict the exploration of the self-reported health in individuals with personal history of infectious and parasitic diseases more thoroughly. The results of the present study therefore need to be

interpreted with caution. Future studies could contrast the types of infectious diseases, their severity and chronicity, and how these factors affect perceived health.^{51,52}

Further, our primary outcome measure, self-rated health, is a subjective rating. Still, health is a multi-faceted concept. The absence of somatic diseases or mental disorder, as assessed by clinicians, does not necessarily indicate excellent health. Note also that perceived health, as assessed here, is a strong predictor of morbidity and mortality.⁵³

Additionally, due to its cross-sectional nature, it cannot establish causality, but only associations. Therefore, it is essential to conduct further research, including qualitative studies, to gain a better understanding of the factors influencing self-perceived health in populations that have experienced multiple infections.

Nonetheless, this study offers valuable insights, particularly through its use of data from a large sample spanning numerous countries.

Our results can contribute to more targeted health support. Health education, including explanations of the non-detrimental findings, may reduce fear and stigma,^{54,55} thereby increasing the well-being and encouraging more people with a parasitic disease to seek health services. Further, given that the relationship between illnesses and self-perceived health varies across countries, one practical implication is the need to tailor public health policies to local contexts. Public health strategies should not only consider general health patterns but also the specific social, cultural, and economic factors of each population. A higher prevalence of poor or very poor self-perceived health is observed among participants living in sparsely populated areas or older populations.⁵⁰

This suggests that public policies should address local challenges, such as improving healthcare access, addressing cultural perceptions of illness, and reducing economic disparities. Tailoring interventions to these specific needs can improve their effectiveness and help reduce health inequalities. Public policies could focus on enhancing healthcare infrastructure, promoting education on managing health conditions, and ensuring that health services are culturally appropriate for each community.

Finally, to improve public health and manage the infectious diseases worldwide, vaccine development is crucial.⁵⁶ As a conclusion, individuals who have experienced infectious and parasitic diseases rate their health as similar good or somewhat higher than those never been exposed to a pathogen.

Author statements

Ethical approval

The original collection of data was approved by the Institutional Review Board (IRB) at the Institute of Psychology, University of Wrocław. All team members adhered to the principles of the Declaration of Helsinki³³ and the ethical guidelines of their respective IRBs, either based on the ethical approval of the Principal Investigator's IRB or their local IRB. All participants provided informed consent.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. Marta Kowal was supported by the Foundation for Polish Science (FNP) START scholarship.

Competing interests

All authors declare no competing interests.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.puhe.2025.02.030>.

References

1. Ansah JP, Chiu CT. Projecting the chronic disease burden among the adult population in the United States using a multi-state population model. *Front Public Health*. 2022;10, 1082183. <https://doi.org/10.3389/fpubh.2022.1082183>.
2. Hajat C, Stein E. The global burden of multiple chronic conditions: a narrative review. *Preventive Medicine Reports*. 2018;12/01/2018;12:284–293. <https://doi.org/10.1016/j.pmedr.2018.10.008>.
3. Uijen AA, van de Lisdonk EH. Multimorbidity in primary care: prevalence and trend over the last 20 years. *Eur J Gen Pract*. 2008;14(Suppl 1):28–32. <https://doi.org/10.1080/13814780802436093>.
4. Roser M, Ritchie H, Spooner F. *Burden of disease*. Our World in Data. 2021.
5. GBD 2019 Diseases and Injuries Collaborators. Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet*. Oct 17 2020;396(10258):1204–1222. [https://doi.org/10.1016/S0140-6736\(20\)30925-9](https://doi.org/10.1016/S0140-6736(20)30925-9).
6. Varo R, Chacour C, Bassat Q. Update on malaria. *Med Clin*. Nov 13 2020;155(9):395–402. <https://doi.org/10.1016/j.medcli.2020.05.010>.
7. Khasnis AA, Nettleman MD. Global warming and infectious disease. *Arch Med Res*. Nov-Dec 2005;36(6):689–696. <https://doi.org/10.1016/j.arcmed.2005.03.041>.
8. Asaolu SO, Ofoezie IE. The role of health education and sanitation in the control of helminth infections. *Acta Trop*. May 2003;86(2-3):283–294. [https://doi.org/10.1016/S0001-706X\(03\)00060-3](https://doi.org/10.1016/S0001-706X(03)00060-3).
9. Lee E, Hoskin AW, Ratnasingam M. *Handbook of Social Status Correlates*. Academic Press; 2018.
10. Murray DR, Schaller M. Historical prevalence of infectious diseases within 230 geopolitical regions: a tool for investigating origins of culture. *J Cross Cult Psychol*. 2009;41(1):99–108. <https://doi.org/10.1177/0022022109349510>.
11. Fincher CL, Thornhill R, Murray DR, Schaller M. Pathogen prevalence predicts human cross-cultural variability in individualism/collectivism. *Proc Biol Sci*. Jun 7 2008;275(1640):1279–1285. <https://doi.org/10.1098/rspb.2008.0094>.
12. Sherman PW, Billing J. Darwinian gastronomy: why we use spices. *Bioscience*. 1999;49(6):453–463. <https://doi.org/10.2307/1313553>.
13. Schaller M, Murray DR. Pathogens, personality, and culture: disease prevalence predicts worldwide variability in sociosexuality, extraversion, and openness to experience. *J Pers Soc Psychol*. Jul 2008;95(1):212–221. <https://doi.org/10.1037/0022-3514.95.1.212>.
14. Mavaddat N, Valderas JM, van der Linde R, Khaw KT, Kinmonth AL. Association of self-rated health with multimorbidity, chronic disease and psychosocial factors in a large middle-aged and older cohort from general practice: a cross-sectional study. *BMC Fam Pract*. Nov 25 2014;15:185. <https://doi.org/10.1186/s12875-014-0185-6>.
15. Palladino R, Tayu Lee J, Ashworth M, Triassi M, Millett C. Associations between multimorbidity, healthcare utilisation and health status: evidence from 16 European countries. *Age Ageing*. May 2016;45(3):431–435. <https://doi.org/10.1093/ageing/afw044>.
16. Samper-Ternent R, Michaels-Obregon A, Wong R, Palloni A. Older adults under a mixed regime of infectious and chronic diseases. *Salud Publica Mex*. Oct 2012;54(5):487–495. <https://doi.org/10.1590/s0036-36342012000500005>.
17. Bowling A. Just one question: if one question works, why ask several? *J Epidemiol Community Health*. May 2005;59(5):342–345. <https://doi.org/10.1136/jech.2004.021204>.
18. Bohlbro AS, Patsche CB, Mendes AM, et al. A self-rated health score predicts severe disease and high mortality in patients with pulmonary TB. *Int J Tuberc Lung Dis*. Feb 1 2022;26(2):158–165. <https://doi.org/10.5588/ijtld.21.0545>.
19. Cullati S, Mukhopadhyay S, Sieber S, Chakraborty A, Burton-Jeangros C. Is the single self-rated health item reliable in India? A construct validity study. *BMJ Glob Health*. 2018;3(6), e000856. <https://doi.org/10.1136/bmjgh-2018-000856>.
20. Kööts-Ausmees L, Realo A. The association between life satisfaction and self-reported health status in Europe. *Eur J Pers*. 2015;29(6):647–657. <https://doi.org/10.1002/per.2037>.
21. DeSalvo KB, Bloser N, Reynolds K, He J, Muntner P. Mortality prediction with a single general self-rated health question. A meta-analysis. *J Gen Intern Med*. Mar 2006;21(3):267–275. <https://doi.org/10.1111/j.1525-1497.2005.00291.x>.
22. Halford C, Wallman T, Welin L, et al. Effects of self-rated health on sick leave, disability pension, hospital admissions and mortality. A population-based longitudinal study of nearly 15,000 observations among Swedish women and men. *BMC Public Health*. Dec 22 2012;12:1103. <https://doi.org/10.1186/1471-2458-12-1103>.
23. Feasey N, Wansbrough-Jones M, Mabey DCW, Solomon AW. Neglected tropical diseases. *Br Med Bull*. 2009;93(1):179–200. <https://doi.org/10.1093/bmb/ldp046>.
24. Hotez PJ. *Introduction to the Neglected Tropical Diseases: The Ancient Afflictions of Stigma and Poverty*. Forgotten People, Forgotten Diseases; 2021:1–16.
25. Huang CH, Tsai YT, Wang SF, Wang WH, Chen YH. Dengue vaccine: an update. *Expert Rev Anti Infect Ther*. Dec 2021;19(12):1495–1502. <https://doi.org/10.1080/14787210.2021.1949983>.
26. Molehin AJ. Schistosomiasis vaccine development: update on human clinical trials. *J Biomed Sci*. Jan 22 2020;27(1):28. <https://doi.org/10.1186/s12929-020-0621-y>.
27. Sahastrabudhe S, Saluja T. Overview of the typhoid conjugate vaccine pipeline: current status and future plans. *Clin Infect Dis*. Feb 15 2019;68(Suppl 1):S22–S26. <https://doi.org/10.1093/cid/ciy884>.
28. Schragger LK, Vekemens J, Dräger N, Lewinsohn DM, Olesen OF. The status of tuberculosis vaccine development. *Lancet Infect Dis*. Mar 2020;20(3):e28–e37. [https://doi.org/10.1016/S1473-3099\(19\)30625-5](https://doi.org/10.1016/S1473-3099(19)30625-5).
29. Stanisic DI, Good MF. Malaria vaccines: progress to date. *BioDrugs*. Nov 2023;37(6):737–756. <https://doi.org/10.1007/s40259-023-00623-4>.

30. Raso G, Lugimbuhl A, Adjoua CA, et al. Multiple parasite infections and their relationship to self-reported morbidity in a community of rural Cote d'Ivoire. *Int J Epidemiol*. Oct 2004;33(5):1092–1102. <https://doi.org/10.1093/ije/dyh241>.
31. Kowal M, Sorokowski P, Pisanski K, et al. Predictors of enhancing human physical attractiveness: data from 93 countries. *Evol Hum Behav*. 2022;43(6):455–474. <https://doi.org/10.1016/j.evolhumbehav.2022.08.003>.
32. Kowal M, Sorokowski P, Dinic BM, et al. Validation of the short version (TLS-15) of the triangular love scale (TLS-45) across 37 languages. *Arch Sex Behav*. Feb 2024;53(2):839–857. <https://doi.org/10.1007/s10508-023-02702-7>.
33. World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA*. Nov 27 2013;310(20):2191–2194. <https://doi.org/10.1001/jama.2013.281053>.
34. Barari S, Caria S, Davola A, et al. Evaluating COVID-19 public health messaging in Italy: self-reported compliance and growing mental health concerns. *medRxiv*. 2020. <https://doi.org/10.1101/2020.03.27.20042820>.
35. Gozgor G. Global evidence on the determinants of public trust in governments during the COVID-19. *Appl Res Qual Life*. 2022;17(2):559–578. <https://doi.org/10.1007/s11482-020-09902-6>.
36. Kapoor H, Tagat A. Worry much? Preventive health behaviours related to worry across countries amid COVID-19. *J Health Psychol*. Apr 2022;27(5):1125–1136. <https://doi.org/10.1177/1359105321999090>.
37. Liu N, Zhong Q. The impact of sports participation on individuals' subjective well-being: the mediating role of class identity and health. *Humanities and Social Sciences Communications*. 2023;10(1). <https://doi.org/10.1057/s41599-023-02064-4>.
38. Steegen S, Tuerlinckx F, Gelman A, Vanpaemel W. Increasing transparency through a multiverse analysis. *Perspect Psychol Sci*. 2016;11(5):702–712. <https://doi.org/10.1177/1745691616658637>.
39. Chen JW, Maldonado DR, Kowalski BL, et al. Best practice guidelines for propensity score methods in medical research: consideration on theory, implementation, and reporting. A review. *Arthrosc J Arthrosc Relat Surg*. 2022/02/01/2022;38(2):632–642. <https://doi.org/10.1016/j.arthro.2021.06.037>.
40. Balduzzi S, Rücker G, Schwarzer G. How to perform a meta-analysis with R: a practical tutorial. *Evid Base Ment Health*. Nov 2019;22(4):153–160. <https://doi.org/10.1136/ebmental-2019-300117>.
41. Cinelli C, Hazlett C. Making sense of sensitivity: extending omitted variable bias. *J Roy Stat Soc B Stat Methodol*. 2019;82(1):39–67. <https://doi.org/10.1111/rssb.12348>.
42. Ho DE, Imai K, King G, Stuart EA. MatchIt: nonparametric preprocessing for parametric causal inference. *J Stat Software*. 2011;42(8). <https://doi.org/10.18637/jss.v042.i08>.
43. Endler NS, Kocovski NL, Macrodimitris SD. Coping, efficacy, and perceived control in acute vs chronic illnesses. *Pers Indiv Differ*. 2001;30(4):617–625. [https://doi.org/10.1016/s0191-8869\(00\)00060-x](https://doi.org/10.1016/s0191-8869(00)00060-x).
44. Anderson RL. Friedrich nietzsche. In: ENZaU Nodelman, ed. *The Stanford Encyclopedia of Philosophy*. Spring 2024. Metaphysics Research Lab, Stanford University; 2024.
45. Luhmann M, Hofmann W, Eid M, Lucas RE. Subjective well-being and adaptation to life events: a meta-analysis. *J Pers Soc Psychol*. Mar 2012;102(3):592–615. <https://doi.org/10.1037/a0025948>.
46. Guma J. What influences individual perception of health? Using machine learning to disentangle self-perceived health. *SSM Popul Health*. Dec 2021;16, 100996. <https://doi.org/10.1016/j.ssmph.2021.100996>.
47. Jylhä M. What is self-rated health and why does it predict mortality? Towards a unified conceptual model. *Soc Sci Med*. Aug 2009;69(3):307–316. <https://doi.org/10.1016/j.socscimed.2009.05.013>.
48. Wood JV, Taylor SE, Lichtman RR. Social comparison in adjustment to breast cancer. *J Pers Soc Psychol*. Nov 1985;49(5):1169–1183. <https://doi.org/10.1037//0022-3514.49.5.1169>.
49. Giacomozzi C, Palmieri L, Gargiulo L, et al. The perceived health status from young adults to elderly: results of the MEHM questionnaire within the CUORE project survey 2008–2012. *Int J Environ Res Publ Health*. Aug 25 2020;17(17). <https://doi.org/10.3390/ijerph17176160>.
50. Shaaban AN, Martins MRO, Peleteiro B. Factors associated with self-perceived health status in Portugal: results from the national health survey 2014. *Front Public Health*. 2022;10, 879432. <https://doi.org/10.3389/fpubh.2022.879432>.
51. Álvarez Cabo D, Alemany A, Martínez Sesmero JM, Moreno Guillén S. Healthcare and economic impact. *Enferm Infecc Microbiol Clín*. Sep 2018;36(Suppl 1):19–25. [https://doi.org/10.1016/s0213-005x\(18\)30242-8](https://doi.org/10.1016/s0213-005x(18)30242-8). Atención sanitaria e impacto económico.
52. Ayuso García B, Pérez López A, Besteiro Balado Y, et al. [Health-related quality of life in patients recovered from COVID-19]. *J Healthc Qual Res*. Jul-Aug 2022;37(4):208–215. <https://doi.org/10.1016/j.jhqr.2022.01.001>. Calidad de vida relacionada con la salud en pacientes recuperados de COVID-19.
53. Cleary PD. Subjective and objective measures of health: which is better when? *J Health Serv Res Policy*. Jan 1997;2(1):3–4. <https://doi.org/10.1177/135581969700200102>.
54. Asilimwe JB, Nagendrappa PB, Atukunda EC, et al. The meaning of caring for patients with cancer among traditional medicine practitioners in Uganda: a grounded theory approach. *PLOS Glob Public Health*. 2023;3(7), e0001764. <https://doi.org/10.1371/journal.pgph.0001764>.
55. Chandra M, Rana P, Chandra K, Arora VK. Tuberculosis - depression syndemic: a public health challenge. *Indian J Tubercul*. Jan 2019;66(1):197–202. <https://doi.org/10.1016/j.ijtb.2019.02.007>.
56. Allen T, Castellanos ME, Giacomini P, et al. Next-generation vaccines for tropical infectious diseases. *Int J Infect Dis*. 2024/06/01/2024;143, 107014. <https://doi.org/10.1016/j.ijid.2024.107014>.